

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

THIS PAGE BLANK (USPTO)



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07K 14/00	A2	(11) International Publication Number: WO 00/24768 (43) International Publication Date: 4 May 2000 (04.05.00)						
<p>(21) International Application Number: PCT/US99/24826</p> <p>(22) International Filing Date: 22 October 1999 (22.10.99)</p> <p>(30) Priority Data:</p> <table border="0"> <tr> <td>60/105,507</td> <td>23 October 1998 (23.10.98)</td> <td>US</td> </tr> <tr> <td>60/108,685</td> <td>16 November 1998 (16.11.98)</td> <td>US</td> </tr> </table> <p>(71) Applicant (<i>for all designated States except US</i>): MASSACHUSETTS INSTITUTE OF TECHNOLOGY [US/US]; 77 Massachusetts Avenue, Cambridge, MA 02138 (US).</p> <p>(72) Inventors; and</p> <p>(75) Inventors/Applicants (<i>for US only</i>): KAWASAKI, Hiroaki [JP/JP]; 3-20-2, Aoba, Higashi-ku, Fukuoka 813-0025 (JP). GRAYBIEL, Ann [US/US]; Boyce Farm Road, Lincoln, MA 01773 (US). HOUSMAN, David [US/US]; 64 Homer Street, Newton, MA 02158 (US).</p> <p>(74) Agent: CAMACHO, Jennifer, A.; Testa, Hurwitz & Thibault, LLP, High Street Tower, 125 High Street, Boston, MA 02110 (US).</p>		60/105,507	23 October 1998 (23.10.98)	US	60/108,685	16 November 1998 (16.11.98)	US	<p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published</p> <p><i>Without international search report and to be republished upon receipt of that report.</i></p>
60/105,507	23 October 1998 (23.10.98)	US						
60/108,685	16 November 1998 (16.11.98)	US						
<p>(54) Title: GENES INTEGRATING SIGNAL TRANSDUCTION PATHWAYS</p> <p>(57) Abstract</p> <p>The present invention describes the identification, isolation, sequencing and characterization of two human CalDAG-GEF, and two human cAMP-GEF genes, which are associated with the Ras pathway. Also identified are CalDAG-GEF gene homologues in mice and cAMP-GEF gene homologues in rats. Nucleic acids and proteins comprising or derived from the CalDAG-GEFs and/or cAMP-GEFs are useful in screening and diagnosing certain Ras-associated cancers, in identifying and developing therapeutics for treatment of certain Ras-associated cancers, and in producing cell lines and transgenic animals useful as models of Ras-associated cancers.</p>								

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

GENES INTEGRATING SIGNAL TRANSDUCTION PATHWAYS

Related Applications

This application claims the benefit of U.S. Application Nos. 60/105,507, filed on October 23, 1998, and 60/108,685, filed on November 16, 1998.

Field of the Invention

The present invention relates generally to novel polynucleotides and proteins encoded by such polynucleotides, along with therapeutic, diagnostic, and research utilities for these polynucleotides and proteins.

Background of the Invention

Ras proteins are key regulators of growth, differentiation and malignant transformation. In addition, these proteins are implicated in synaptic function and region-specific learning and memory functions in the brain.

As shown schematically in Figure 1, Ras proteins cycle between inactive GDP-complexed and active GTP-complexed states. GTPase-activating proteins (GAPs) inactivate Ras proteins by stimulating hydrolysis of the bound GTP to GDP, whereas guanine nucleotide exchange factors (GEFs) activate Ras proteins by stimulating release of GDP and the uptake of GTP. So essential are GEFs to Ras action, that genetic loss of GEF function has similar effects to those induced by loss of the Ras proteins themselves. Loss of GEF function can be circumvented by mutations that constitutively activate the Ras proteins, such as an oncogene mutation, or, in some cases, through loss of GAP activity. Activated Ras proteins, which are localized at the plasma membrane, transmit signals from tyrosine kinases to a cascade of serine/threonine kinases, which delivers the signals to the cell nucleus.

Activation of Ras can result in the activation of the mitogen-activated protein (MAP) kinase (also known as extracellular-signal regulated kinase, or ERK) pathway. For example, a receptor tyrosine kinase is activated by a peptide mitogen such as epidermal growth factor (EGF). The EGF-stimulated receptor undergoes autophosphorylation of specific tyrosine residues in its cytoplasmic domain which creates phosphotyrosyl binding sites for the Src homology 2 (SH2) and/or phosphotyrosyl binding (PTB) domains of certain adapter proteins. The adapter protein becomes autophosphorylated on association with activated receptor tyrosine kinases. The GEF is stably associated with the adapter protein which, upon autophosphorylation, mediates translocation of the GEF to the plasma membrane. The GEF then activates the Ras protein. Activated Ras relays its signal downstream through a cascade of cytoplasmic proteins, including

- 2 -

Raf-1 serine/threonine kinase. The Ras:Raf association promotes translocation of the normally cytoplasmic Raf protein to the plasma membrane, where subsequent events lead to the activation of its kinase function. Upon activation, Raf phosphorylates and activates two MAP kinases (also known as MEKs). MEKs directly associate with the catalytic domain of Raf-1 and are
5 phosphorylated by Raf. Activated MEKs function as dual-specificity kinases and phosphorylate tandem threonine and tyrosine residues in the MAP kinases to activate them. Once activated, the MAP kinases translocate to the nucleus where they phosphorylate and activate a variety of substrates.

Rap proteins, members of the Ras small GTPase superfamily, can inhibit Ras signaling of
10 the Ras/Raf-1(a serine/threonine kinase)/MAP kinase pathway or, through B-Raf, can activate MAP kinase. Rap1 consists of two isoforms, Rap1A and Rap1B, which differ mainly at the C-terminus. Characteristic features of Rap1 are its geranylgeranyl modification at the C-terminus, which is responsible for membrane attachments, and a threonine residue at position 61. In most other GTPases, the corresponding residue is a glutamine. Rap proteins, like Ras proteins, cycle
15 between inactive GDP-complexed and active GTP-complexed states. GEFs are required to activate Rap proteins by stimulating the release of GDP and the uptake of GTP.

Constitutive activation of the Ras pathway contributes to malignant transformation. In fact, the Ras gene has been implicated in many human cancers, including lung cancer, breast cancer, colorectal cancer, exocrine pancreatic cancer, and myeloid leukemia. Biological and
20 biochemical studies of Ras action indicate that Ras functions like a G-regulatory protein since Ras must be localized in the plasma membrane and must bind with GTP in order to transform cells. Gibbs et al., 53 MICROBIOL. REV. 171-286 (1989).

Targeting components of the Ras signaling pathways has been proposed as one approach for the development of anti-Ras drugs for cancer treatment. One potential approach for targeting
25 Ras for cancer treatment involves the use of farnesyltransferase inhibitors (FTIs). Inhibition of farnesyl-protein transferase, and thereby of farnesylation of the Ras protein, blocks the ability of Ras to transform normal cells to cancer cells. Certain inhibitors of Ras farnesylation cause an increase in soluble Ras which can act as a dominant negative inhibitor of Ras function. While soluble Ras in cancer cells can become a dominant negative inhibitor, soluble Ras in normal cells
30 would not be an inhibitor. A cytosol-localized and activated form of Ras acts as a dominant negative Ras inhibitor of membrane-bound Ras function. Gibbs et al., 86 PROC. NAT'L ACAD.

- 3 -

Sci. USA 6630-34 (1989). FTIs block Ras function by preventing its post-translational modification by the farnesyl isoprenoid.

Intervention of Ras signaling at multiple or various points can significantly impact the ability of Ras to cause cellular transformation. Since Ras protein function is believed to be crucial to so many cellular processes, targeting only a subset of Ras functions by downstream
5 intervention may provide significant advantages. Thus, there remains a need for identifying additional means for disrupting the Ras pathway. Applicants have discovered four new targets, namely GEFs specific for Rap1A, for disrupting the Ras pathway.

Summary of the Invention

10 Applicants have discovered four mammalian genes which have been designated CalDAG-GEFI, CalDAG-GEFII, cAMP-GEFI, and cAMP-GEFII, which encode proteins having a substrate specificity for Rap1A. The proteins encoded by CalDAG-GEFI and CalDAG-GEFII, referred to herein generally as "CalDAG-GEF," have dual binding domains for calcium and diacylglycerol. The proteins cAMP-GEFI and cAMP-GEFII, referred to herein generally as
15 "cAMP-GEF," have a binding domain for cyclic adenosine 3', 5'-monophosphate. The present disclosure provides polypeptide and polynucleotide sequences for Mus musculus CalDAG-GEFI, Homo sapiens CalDAG-GEFI, Rattus norvegicus CalDAG-GEFII, Homo sapiens CalDAG-GEFII, Rattus norvegicus cAMP-GEFI, Homo sapiens cAMP-GEFI, Homo sapiens alternatively spliced cAMP-GEFI, Rattus norvegicus cAMP-GEFII, and Homo sapiens cAMP-GEFII. See
20 Kawasaki et al., 95 Proc. Natl. Acad. Sci. USA 13278-83 (1998), and Kawasaki et al., 282 Sci. 2275-79 (1998), the disclosures of both of which are incorporated by reference herein.

Thus, in one series of embodiments, the present invention provides isolated nucleic acids including nucleotide sequences comprising or derived from CalDAG-GEF or cAMP-GEF, or encoding polypeptides comprising or derived from CalDAG-GEF or cAMP-GEF proteins. The
25 sequences of the invention include the specifically disclosed sequences, splice variants of these sequences, allelic variants of these sequences, synonymous sequences, and homologous or orthologous variants of these sequences. Thus, for example, the invention provides nucleic acid sequences from the Mus musculus CalDAG-GEFI, Homo sapiens CalDAG-GEFI, Rattus norvegicus CalDAG-GEFII, Homo sapiens CalDAG-GEFII, Rattus norvegicus cAMP-GEFI,
30 Homo sapiens cAMP-GEFI, Homo sapiens alternatively spliced cAMP-GEFI, Rattus norvegicus cAMP-GEFII, and Homo sapiens cAMP-GEFII. The present invention also provides allelic variants and homologous or orthologous sequences by providing methods by which such variants

- 4 -

may be routinely obtained. Because the nucleic acids of the invention may be used in a variety of diagnostic, therapeutic and recombinant applications, various subsets of the CalDAG-GEF and cAMP-GEF sequences are also provided. For example, for use in allele specific hybridization screening or PCR amplification techniques, subsets of the CalDAG-GEF and cAMP-GEF sequences, including both sense and antisense sequences, and both normal and mutant sequences, as well as intronic, exonic and untranslated sequences, are provided. Such sequences may comprise a small number of consecutive nucleotides from the sequences which are disclosed or otherwise enabled herein, but preferably include at least 8-10, more preferably 10-15, and most preferably 15-25, consecutive nucleotides from a CalDAG-GEF or cAMP-GEF sequence. In another embodiment, such sequences include at least 25-500 consecutive nucleotides from CalDAG-GEF or cAMP-GEF sequence. Other preferred subsets of a CalDAG-GEF or cAMP-GEF sequence include those encoding one or more of the functional domains or antigenic determinants of the CalDAG-GEF or cAMP-GEF protein and, in particular, may include either normal (wild-type) or mutant sequences. The invention also provides for various nucleic acid constructs in which CalDAG-GEF or cAMP-GEF sequences, either complete or subsets, are operably joined to exogenous sequences to form cloning vectors, expression vectors, fusion vectors, transgenic constructs, and the like. Thus, in accordance with another aspect of the invention, a recombinant vector for transforming a mammalian or invertebrate tissue cell to express a normal or mutant CalDAG-GEF and/or cAMP-GEF sequence in the cells is provided.

In another series of embodiments, the present invention provides for host cells which have been transfected or otherwise transformed with one of the nucleic acids of the invention. The cells may be transformed merely for purposes of propagating the nucleic acid constructs of the invention, or may be transformed so as to express the CalDAG-GEF and/or cAMP-GEF sequences. The transformed cells of the invention may be used in assays to identify proteins and/or other compounds which affect normal or mutant CalDAG-GEF and/or cAMP-GEF expression, which interact with the normal or mutant CalDAG-GEF and/or cAMP-GEF proteins, and/or which modulate the function or effects of the normal or mutant proteins, or to produce the CalDAG-GEF and/or cAMP-GEF proteins, fusion proteins, functional domains, antigenic determinants, and/or antibodies of the invention. Transformed cells may also be implanted into hosts, including humans, for therapeutic or other reasons. Preferred host cells include mammalian cells, including pure or mixed cell cultures, as well as bacterial, yeast, nematode,

- 5 -

insect and other invertebrate cells. For uses as described below, preferred cells also include embryonic stem cells, zygotes, gametes, and germ line cells.

In another series of embodiments, the present invention provides transgenic animal models of diseases or disorders associated with mutations in the CalDAG-GEF and/or cAMP-GEF genes. The animal may be essentially any non-human mammal, including rats, mice, hamsters, guinea pigs, rabbits, dogs, cats, goats, sheep, pigs, and non-human primates. In addition, invertebrate models, including nematodes and insects, may be used for certain applications. The animal models are produced by standard transgenic methods including microinjection, electroporation, transfection, or other forms of transformation of embryonic stem cells, zygotes, gametes, and germ line cells with vectors including genomic or cDNA fragments, minigenes, homologous recombination vectors, viral insertion vectors and the like. Suitable vectors include vaccinia virus, adenovirus, adeno-associated virus, retrovirus, liposome transport, neuraltropic viruses, and Herpes simplex virus. The animal models may include transgenic sequences comprising or derived from the CalDAG-GEF and/or cAMP-GEF genes, including normal and mutant sequences, intronic, exonic and untranslated sequences, and sequences encoding subsets of the CalDAG-GEF and/or cAMP-GEF proteins, such as functional domains. The major types of animal models provided include: (1) Animals in which a normal human CalDAG-GEF and/or cAMP-GEF gene has been recombinantly introduced into the genome of the animal as an additional gene, under the regulation of either an exogenous or an endogenous promoter element, and as either a recombinant gene or a large genomic fragment; in which a normal human CalDAG-GEF and/or cAMP-GEF gene has been recombinantly substituted for one or both copies of the animal's homologous CalDAG-GEF and/or cAMP-GEF gene by homologous recombination or gene targeting; and/or in which one or both copies of one of the animal's homologous CalDAG-GEF and/or cAMP-GEF genes have been recombinantly "humanized" by the partial substitution of sequences encoding the human homologue by homologous recombination or gene targeting; (2) Animals in which a mutant human CalDAG-GEF and/or cAMP-GEF gene has been recombinantly introduced into the genome of the animal as an additional gene, under the regulation of either an exogenous or an endogenous promoter element, and as either a recombinant gene or a large genomic fragment; in which a mutant human CalDAG-GEF and/or cAMP-GEF gene has been recombinantly substituted for one or both copies of the animal's homologous CalDAG-GEF and/or cAMP-GEF gene by homologous recombination or gene targeting; and/or in which one or both copies of one of the animal's

- 6 -

homologous CalDAG-GEF and/or cAMP-GEF gene have been recombinantly "humanized" by the partial substitution of sequences encoding a mutant human homologue by homologous recombination or gene targeting; (3) Animals in which a mutant version of one of that animal's CalDAG-GEF or cAMP-GEF gene has been recombinantly introduced into the genome of the animal as an additional gene, under the regulation of either an exogenous or an endogenous promoter element, and as either a recombinant gene or a large genomic fragment; and/or in which a mutant version of one of that animal's CalDAG-GEF or cAMP-GEF gene has been recombinantly substituted for one or both copies of the animal's homologous CalDAG-GEF or cAMP-GEF gene by homologous recombination or gene targeting; and (4) "Knock-out" animals in which one or both copies of one of the animal's CalDAG-GEF or cAMP-GEF genes have been partially or completely deleted by homologous recombination or gene targeting, or have been inactivated by the insertion or substitution by homologous recombination or gene targeting of exogenous sequences.

In another series of embodiments, the present invention provides for substantially pure protein preparations including polypeptides comprising or derived from the CalDAG-GEF and/or cAMP-GEF proteins. The CalDAG-GEF and cAMP-GEF protein sequences of the invention include the specifically disclosed sequences, variants of these sequences resulting from alternative mRNA splicing, allelic variants of these sequences, and homologous or orthologous variants of these sequences. Thus, for example, the invention provides amino acid sequences from the *Mus musculus* CalDAG-GEFI protein, *Homo sapiens* CalDAG-GEFI protein, *Rattus norvegicus* CalDAG-GEFII protein, *Homo sapiens* CalDAG-GEFII protein, *Rattus norvegicus* cAMP-GEFI protein, *Homo sapiens* cAMP-GEFI protein, *Homo sapiens* alternatively spliced cAMP-GEFI protein, *Rattus norvegicus* cAMP-GEFII protein, and *Homo sapiens* cAMP-GEFII protein. The present invention also provides allelic variants and homologous or orthologous proteins by providing methods by which such variants may be routinely obtained. The present invention also specifically provides for mutant or disease-causing variants of CalDAG-GEF and cAMP-GEF by providing methods by which such variants may be routinely obtained. Because the proteins of the invention may be used in a variety of diagnostic, therapeutic and recombinant applications, various subsets of the CalDAG-GEF and cAMP-GEF protein sequences and combinations of the CalDAG-GEF and cAMP-GEF protein sequences with heterologous sequences are also provided. For example, for use as immunogens or in binding assays, subsets of the CalDAG-GEF and cAMP-GEF protein sequences, including both normal and mutant

- 7 -

sequences, are provided. Such protein sequences may comprise a small number of consecutive amino acid residues from the sequences which are disclosed or otherwise enabled herein, but preferably include at least 4-8, and preferably at least 9-15 consecutive amino acid residues from a CalDAG-GEF or cAMP-GEF sequence. In another embodiment, such sequences comprise at least 15-200 consecutive amino acid residues from a CalDAG-GEF or cAMP-GEF sequence. Other preferred subsets of the CalDAG-GEF and cAMP-GEF protein sequences include those corresponding to one or more of the functional domains or antigenic determinants of the CalDAG-GEF and cAMP-GEF proteins and, in particular, may include either normal (wild-type) or mutant sequences. The invention also provides for various protein constructs in which a CalDAG-GEF and/or cAMP-GEF sequences, either complete or subsets thereof, are joined to exogenous sequences to form fusion proteins and the like. In accordance with these embodiments, the present invention also provides for methods of producing all of the above described proteins which comprise, or are derived from, CalDAG-GEF and/or cAMP-GEF.

In another series of embodiments, the present invention provides for the production and use of polyclonal and monoclonal antibodies, including antibody fragments, including Fab fragments, F(ab')₂, and single chain antibody fragments, which selectively bind to CalDAG-GEF or cAMP-GEF, or to specific antigenic determinants of CalDAG-GEF or cAMP-GEF. The antibodies may be raised in mouse, rabbit, goat or other suitable animals, or may be produced recombinantly in cultured cells such as hybridoma cell lines. Preferably, the antibodies selectively bind to a sequence comprising at least 4-8, and preferably at least 9-15, consecutive amino acid residues from a CalDAG-GEF or cAMP-GEF sequence. The antibodies of the invention may be used in the various diagnostic, therapeutic and technical applications described herein.

In another series of embodiments, the present invention provides methods of screening or identifying proteins, small molecules or other compounds which are capable of inducing or inhibiting the expression and/or function of the CalDAG-GEF and/or cAMP-GEF genes or proteins. The assays may be performed *in vitro* using non-transformed cells, immortalized cell lines, or recombinant cell lines, or *in vivo* using the transgenic animal models enabled herein. In particular, the assays may detect the presence of increased or decreased expression of CalDAG-GEF and/or cAMP-GEF-related genes or proteins on the basis of increased or decreased mRNA expression, increased or decreased levels of CalDAG-GEF and/or cAMP-GEF-related protein products, or increased or decreased levels of expression of a marker gene (*e.g.*, β -galactosidase,

- 8 -

green fluorescent protein, alkaline phosphatase or luciferase) operably joined to a 5' regulatory region in a recombinant construct. Cells known to express CalDAG-GEF or cAMP-GEF, or transformed to express CalDAG-GEF or cAMP-GEF, are incubated and one or more test compounds are added to the medium. After allowing a sufficient period of time (*e.g.*, 0-72 hours) for the compound to induce or inhibit the expression of the CalDAG-GEF or cAMP-GEF, any change in levels of expression from an established baseline may be detected using any of the techniques described above. In particularly preferred embodiments, the cells are from an immortalized cell line such as a human neuroblastoma, glioblastoma or a hybridoma cell line, or are transformed cells of the invention.

In another series of embodiments, the present invention provides methods for identifying proteins and other compounds which bind to, or otherwise directly interact with, CalDAG-GEF and/or cAMP-GEF. The proteins and compounds will include endogenous cellular components which interact with the CalDAG-GEF and/or cAMP-GEF *in vivo* and which, therefore, provide new targets for pharmaceutical and therapeutic interventions, as well as recombinant, synthetic, and otherwise exogenous compounds which may have CalDAG-GEF and/or cAMP-GEF binding capacity and, therefore, may be candidates for pharmaceutical agents. Thus, in one series of embodiments, cell lysates or tissue homogenates (*e.g.*, human brain homogenates, lymphocyte lysates) may be screened for proteins or other compounds which bind to one of the normal or mutant CalDAG-GEF or cAMP-GEF proteins. Alternatively, any of a variety of exogenous compounds, both naturally occurring and/or synthetic (*e.g.*, libraries of small molecules or peptides), may be screened for CalDAG-GEF or cAMP-GEF binding capacity. In each of these embodiments, an assay is conducted to detect binding between a "CalDAG-GEF component" or a "cAMP-GEF component" and some other moiety. In one embodiment, a CalDAG-GEF component comprises a CalDAG-GEF SRC1, SRC2, SRC3, EF hand or a DAG-binding domain. In another embodiment, a cAMP-GEF component comprises a cAMP-GEF SRC1, SRC2, SRC3, or a cAMP-binding domain. The "CalDAG-GEF component" or the "cAMP-GEF component" in these assays may be any polypeptide comprising or derived from a normal or mutant CalDAG-GEF or cAMP-GEF protein, including functional domains or antigenic determinants of CalDAG-GEF or cAMP-GEF, or CalDAG-GEF or cAMP-GEF fusion proteins. Binding may be detected by non-specific measures (*e.g.*, changes in intracellular Ca^{2+} , GTP/GDP ratio) or by specific measures (*e.g.*, changes in the expression of downstream genes which can be monitored by differential display, 2D gel electrophoresis, differential hybridization, or SAGE methods). The

- 9 -

preferred methods involve variations on the following techniques: (1) direct extraction by affinity chromatography; (2) co-isolation of CalDAG-GEF or cAMP-GEF components and bound proteins or other compounds by immunoprecipitation; (3) the Biomolecular Interaction Assay (BIAcore); and (4) the yeast two-hybrid systems.

5 In another series of embodiments, the present invention provides for methods of identifying proteins, small molecules and other compounds capable of modulating the activity of normal or mutant CalDAG-GEF or cAMP-GEF. Using normal cells or animals, the transformed cells and transgenic animal models of the present invention, or cells obtained from subjects bearing normal or mutant CalDAG-GEF or cAMP-GEF genes, the present invention provides
10 methods of identifying such compounds on the basis of their ability to affect the expression of CalDAG-GEF and/or cAMP-GEF, the intracellular localization of the CalDAG-GEF and/or cAMP-GEF, or other biochemical, histological, or physiological markers which distinguish cells bearing normal and mutant CalDAG-GEF and/or cAMP-GEF sequences. Using the transgenic animals of the invention, methods of identifying such compounds are also provided on the basis
15 of the ability of the compounds to affect behavioral, physiological or histological phenotypes associated with mutations in CalDAG-GEF and/or cAMP-GEF.

In another series of embodiments, the present invention provides methods and reagents for the screening and diagnosis of diseases or disorders associated with mutations in the CalDAG-GEF and/or cAMP-GEF genes. Screening and/or diagnosis can be accomplished by
20 methods based upon the nucleic acids (including genomic and mRNA/cDNA sequences), proteins, and/or antibodies disclosed and enabled herein, including functional assays designed to detect failure or augmentation of the normal CalDAG-GEF and/or cAMP-GEF activity and/or the presence of specific new activities conferred by the mutant CalDAG-GEF and/or cAMP-GEF. Thus, for example, screens and diagnostics based upon CalDAG-GEF and/or cAMP-GEF
25 proteins are provided which detect differences between mutant and normal CalDAG-GEF or cAMP-GEF in electrophoretic mobility, in proteolytic cleavage patterns, in molar ratios of the various amino acid residues, or in ability to bind specific antibodies. In addition, screens and diagnostics based upon nucleic acids (gDNA, cDNA or mRNA) are provided which detect differences in nucleotide sequences by direct nucleotide sequencing, hybridization using allele
30 specific oligonucleotides, restriction enzyme digest and mapping (*e.g.*, RFLP, REF-SSCP), electrophoretic mobility (*e.g.*, SSCP, DGGE), PCR mapping, RNase protection, chemical mismatch cleavage, ligase-mediated detection, and various other methods. Other methods are

- 10 -

also provided which detect abnormal processing of CalDAG-GEF and/or cAMP-GEF or proteins reacting with CalDAG-GEF and/or cAMP-GEF, alterations in CalDAG-GEF and/or cAMP-GEF transcription, translation, and post-translational modification; alterations in the intracellular and extracellular trafficking of CalDAG-GEF and/or cAMP-GEF gene products; or abnormal
5 intracellular localization of CalDAG-GEF and/or cAMP-GEF. Such methods and reagents are also useful in the analysis of neoplasias and mammalian immune system function, as well as functional *in vivo* imaging of mammalian organ systems. In accordance with these embodiments, diagnostic kits are also provided which will include the reagents necessary for the above-described diagnostic screens.

10 In another series of embodiments, the present invention provides methods and therapeutic agents for use in the treatment of conditions such as neurological and neuropsychiatric disorders such as Huntington's disease, Parkinson's disease, Alzheimer's disease, dystonia, Tourette's syndrome, obsessive-compulsive disorder, attention deficit/hyperactive disorder, depression, schizophrenia, and stroke; neoplasias such as solid tumors including colon, breast, lung, prostate,
15 and hematopoietic tumors such as leukemia, Hodgkins, and non-Hodgkins lymphomas; and autoimmune diseases, allergies, and asthma; as well as for the enhancement of the immune response in normal and immunocompromised individuals. These methods and therapeutic agents may be based upon (1) administration of normal CalDAG-GEF and/or cAMP-GEF proteins; (2) gene therapy with normal CalDAG-GEF and/or cAMP-GEF genes to compensate for or replace
20 the mutant genes; (3) gene therapy based upon antisense sequences to mutant CalDAG-GEF and/or cAMP-GEF genes or upon sequences which "knock-out" the mutant genes; (4) gene therapy based upon sequences which encode a protein which blocks or corrects the deleterious effects of CalDAG-GEF and/or cAMP-GEF mutants; (5) immunotherapy based upon antibodies to normal and/or mutant CalDAG-GEF and/or cAMP-GEF proteins; or (6) small molecules
25 (drugs) which alter CalDAG-GEF and/or cAMP-GEF expression, block interactions between (normal or mutant) forms of CalDAG-GEF and/or cAMP-GEF and other proteins or ligands, or which otherwise block the function of (normal or mutant) CalDAG-GEF and/or cAMP-GEF proteins by altering the structure of the proteins, by enhancing their metabolic clearance, or by inhibiting their function.

30 In accordance with another aspect of the invention, the proteins of the invention can be used as starting points for rational drug design to provide ligands, therapeutic drugs or other types of small chemical molecules. Alternatively, small molecules or other compounds

identified by the above-described screening assays may serve as "lead compounds" in rational drug design.

Brief Description of the Drawings

Figure 1 is a partial schematic diagram of a Ras pathway.

5 Figure 2A shows human (h) and mouse (m) CalDAG-GEFI, human (h) and rat (r) CalDAG-GEFII, and *C. elegans* (cel) (F25B3.3, GenBank accession number: 1262950) CalDAG-GEF. Figure 2B shows a computer-generated phylogenetic tree analysis of the GEF domains of CalDAG-GEFI and CalDAG-GEFII in relation to other Ras-superfamily GEFs. Figure 2C shows multiple alignment of GEF structurally conserved regions (SCRs) of CalDAG-
10 GEFs and several other GEFs of the Ras superfamily. Figure 2D shows the full-length amino acid sequences of human (h) and mouse (m) CalDAG-GEFI (box indicates amino acid differences). Figure 2E shows the sequence similarity (black indicates identity) of EF-hand domains in CalDAG-GEFs and other calcium binding proteins. Figure 2F shows the sequence similarity of DAG-binding domains of CalDAG-GEFs and PKC (protein kinase C) family
15 proteins.

Figure 3A is a schematic representation of cAMP-GEF family proteins, including human (h) and rat (r) cAMP-GEFI, human (h) cAMP-GEFII and *C. elegans* (cel) (T2OG5.5, GenBank accession number: 458480) cAMP-GEF. Figure 3B is a phylogenetic tree analysis of cAMP binding domains of cAMP-GEFI and II and other cyclic nucleotide binding proteins. Figure 3C
20 is a phylogenetic tree analysis of GEF domains of cAMP-GEFI and II and other Ras superfamily GEFs. Figure 3D shows the amino acid sequences of the structurally conserved regions (SCRs) of cAMP-GEFs and other Ras superfamily GEFs (black indicates identity). Figure 3E shows the amino acid sequences of the cAMP binding pockets of cAMP-GEFI and II and other cyclic nucleotide-binding proteins. The positions of invariant amino acid residues are shown by black
25 diamonds. The open diamond indicates the amino acid that determines the binding specificity for cAMP or cGMP. The arrow indicates the position of amino acid substitutions specific to cAMP-GEFs. Figure 3F is the full-length amino acid sequences of human cAMP-GEFI and II (boxes indicate amino acid identity).

Detailed Description of the Invention

30 The present invention is based, in part, upon the discovery of a family of mammalian genes which are associated with the Ras pathway. The discovery of these genes, designated

- 12 -

CalDAG-GEFI, CalDAG-GEFII, cAMP-GEFI, and cAMP-GEFII, as well as the characterization of these genes, their protein products, mutants, and possible functional roles, are described below.

I. Definitions

5 In order to facilitate review of the various embodiments of the invention, and an understanding of the various elements and constituents used in making and using the invention, the following definitions are provided for particular terms used in the description and the claims which follow:

CalDAG-GEF. As used without further modification herein, the terms "CalDAG-GEF" or
10 or "CalDAG-GEFs" refer to the CalDAG-GEFI and/or the CalDAG-GEFII genes/proteins. In particular, the unmodified terms "CalDAG-GEF" or "CalDAG-GEFs" refer to the mammalian genes/proteins and, preferably, the human genes/proteins.

cAMP-GEF. As used without further modification herein, the terms "cAMP-GEF" or
15 "cAMP-GEFs" refer to the cAMP-GEFI and/or the cAMP-GEFII genes/proteins. In particular, the unmodified terms "cAMP-GEF" or "cAMP-GEFs" refer to the mammalian genes/proteins and, preferably, the human genes/proteins.

CalDAG-GEF gene. As used herein, the term "CalDAG-GEF gene" means the
mammalian genes represented by SEQ ID NOS: 1, 3, 5, and 7, as well as any allelic variants and heterospecific mammalian homologues. A murine CalDAG-GEFI cDNA sequence is disclosed
20 herein as SEQ ID NO: 1, and a human CalDAG-GEFI cDNA sequence is disclosed herein as SEQ ID NO: 3. A rat CalDAG-GEFII cDNA sequence is disclosed herein as SEQ ID NO: 5, and a human CalDAG-GEFII cDNA sequence is disclosed herein as SEQ ID NO: 7. The term "CalDAG-GEF gene" primarily relates to a coding sequence, but can also include some or all of the flanking regulatory regions and/or introns. The term "CalDAG-GEF gene" specifically
25 includes artificial or recombinant genes created from cDNA or genomic DNA, including recombinant genes based upon splice variants.

CalDAG-GEF protein. As used herein, the term "CalDAG-GEF protein" means a protein encoded by a CalDAG-GEF gene, including allelic variants and heterospecific mammalian homologues. A murine CalDAG-GEFI protein sequence is disclosed herein as SEQ ID NO: 2,
30 and a human CalDAG-GEFI protein sequence is disclosed herein as SEQ ID NO: 4. A rat CalDAG-GEFII protein sequence is disclosed herein as SEQ ID NO: 6, and a human CalDAG-GEFII protein sequence is disclosed herein as SEQ ID NO: 8. Splice variants are also embraced

- 13 -

by the term CalDAG-GEF protein as used herein. The protein may be produced by recombinant cells or organisms, may be substantially purified from natural tissues or cell lines, or may be synthesized chemically or enzymatically. Therefore, the term "CalDAG-GEF protein" is intended to include the protein in glycosylated, partially glycosylated, or unglycosylated forms, as well as in phosphorylated, partially phosphorylated, unphosphorylated, sulphated, partially sulphated, or unsulphated forms. The term also includes allelic variants and other functional equivalents of the CalDAG-GEF amino acid sequences, including biologically active proteolytic or other fragments.

hCalDAG-GEF gene and/or protein. As used herein, the abbreviation "hCalDAG-GEF" refers to the human homologue and human allelic variants of the CalDAG-GEF genes and/or proteins. Two cDNA sequences of the human CalDAG-GEF genes are disclosed herein as SEQ ID NOS: 3 and 7. The corresponding hCalDAG-GEF protein sequences are disclosed herein as SEQ ID NOS: 4 and 8. Allelic variants, including deleterious mutants, are enabled in the description which follows.

mCalDAG-GEF gene and/or protein. As used herein, the abbreviation "mCalDAG-GEF" refers to the murine homologues and murine allelic variants of the CalDAG-GEF gene and/or protein. A cDNA sequence of one murine CalDAG-GEF gene is disclosed herein as SEQ ID NO: 16. The corresponding mCalDAG-GEF protein sequence is disclosed herein as SEQ ID NO: 17. Allelic variants, including deleterious mutants, are enabled in the description which follows.

rCalDAG-GEF gene and/or protein. As used herein, the abbreviation "rCalDAG-GEF" refers to the rat homologue and rat allelic variants of the CalDAG-GEF genes and/or proteins. A cDNA sequence of one rat CalDAG-GEF gene is disclosed herein as SEQ ID NO: 5. The corresponding rCalDAG-GEF protein sequence is disclosed herein as SEQ ID NO: 6. Allelic variants, including deleterious mutants, are enabled in the description which follows.

cAMP-GEF gene. As used herein, the term "cAMP-GEF gene" means the mammalian genes represented by SEQ ID NOS: 9, 11, 13, 15, and 17, as well as any allelic variants and heterospecific mammalian homologues. A rat cAMP-GEFI cDNA sequence is disclosed herein as SEQ ID NO: 9, and a human cAMP-GEFI cDNA sequence is disclosed as SEQ ID NO: 11.

Another human cAMP-GEFI cDNA sequence, resulting from alternative splicing of the mRNA transcript, is disclosed as SEQ ID NO: 13. A rat cAMP-GEFII cDNA sequence is disclosed as SEQ ID NO: 15, and a human cAMP-GEFII cDNA sequence is disclosed as SEQ ID NO: 17.

- 14 -

The term "cAMP-GEF gene" primarily relates to a coding sequence, but can also include some or all of the flanking regulatory regions and/or introns. The term cAMP-GEF gene specifically includes artificial or recombinant genes created from cDNA or genomic DNA, including recombinant genes based upon splice variants.

5 cAMP-GEF protein. As used herein, the term "cAMP-GEF protein" means a protein encoded by a cAMP-GEF gene, including allelic variants and heterospecific mammalian homologues. A rat cAMP-GEFI protein sequence is disclosed herein as SEQ ID NO: 10, and a human cAMP-GEFI protein sequence is disclosed as SEQ ID NO: 12. Another human cAMP-GEFI protein sequence, resulting from alternative splicing of the mRNA transcript, is disclosed
10 as SEQ ID NO: 14. A rat cAMP-GEFII protein sequence is disclosed as SEQ ID NO: 16, and a human cAMP-GEFII protein sequence is disclosed as SEQ ID NO: 18. Splice variants are also embraced by the term cAMP-GEF protein as used herein. The protein may be produced by recombinant cells or organisms, may be substantially purified from natural tissues or cell lines, or may be synthesized chemically or enzymatically. Therefore, the term "cAMP-GEF protein" is
15 intended to include the protein in glycosylated, partially glycosylated, or unglycosylated forms, as well as in phosphorylated, partially phosphorylated, unphosphorylated, sulphated, partially sulphated, or unsulphated forms. The term also includes allelic variants and other functional equivalents of the cAMP-GEF amino acid sequences, including biologically active proteolytic or other fragments.

20 hcAMP-GEF gene and/or protein. As used herein, the abbreviation "hcAMP-GEF" refers to the human homologue and human allelic variants of the cAMP-GEF gene and/or protein. One cDNA sequences of the human cAMP-GEF gene is disclosed herein as SEQ ID NO: 18. The corresponding hcAMP-GEF protein sequence is disclosed herein as SEQ ID NO: 19. Numerous allelic variants, including deleterious mutants, are disclosed and enabled throughout the
25 description which follows.

rcAMP-GEF gene and/or protein. As used herein, the abbreviation "rcAMP-GEF" refers to the rat homologue and rat allelic variants of the cAMP-GEF gene and/or protein. Two cDNA sequences of rat cAMP-GEF genes are disclosed herein as SEQ ID NOS: 9 and 15. The corresponding rcAMP-GEF protein sequences are disclosed herein as SEQ ID NOS: 10 and 16..
30 Numerous allelic variants, including deleterious mutants, are disclosed and enabled throughout the description which follows.

- 15 -

Normal. As used herein with respect to genes, the term "normal" refers to a gene which encodes and expresses a normal protein. As used herein with respect to proteins, the term "normal" means a protein which performs its usual or normal physiological role and which is not associated with, or causative of, a pathogenic condition or state. Therefore, as used herein, the term "normal" is essentially synonymous with the usual meaning of the phrase "wild type." For any given gene, or corresponding protein, a multiplicity of normal allelic variants may exist, none of which is associated with the development of a pathogenic condition or state. Such normal allelic variants include, but are not limited to, variants in which one or more nucleotide substitutions do not result in a change in the encoded amino acid sequence.

Mutant. As used herein with respect to genes, the term "mutant" refers to a gene which encodes a mutant protein and/or fails to express a normal protein. As used herein with respect to proteins, the term "mutant" means a protein which does not perform its usual or normal physiological role and which is associated with, or causative of, a pathogenic condition or state. Therefore, as used herein, the term "mutant" is essentially synonymous with the terms "dysfunctional," "pathogenic," "disease-causing," and "deleterious." With respect to the CalDAG-GEF and cAMP-GEF genes and proteins of the present invention, the term "mutant" refers to CalDAG-GEF and cAMP-GEF genes/proteins bearing one or more nucleotide/amino acid substitutions, insertions and/or deletions which cause the genes/proteins to be dysfunctional, pathogenic, disease-causing or otherwise deleterious. This definition is understood to include the various mutations that naturally exist, including but not limited to those disclosed herein, as well as synthetic or recombinant mutations produced by human intervention. The term "mutant," as applied to the CalDAG-GEF and cAMP-GEF genes, is not intended to embrace sequence variants which, due to the degeneracy of the genetic code, encode proteins identical to the normal sequences disclosed or otherwise enabled herein; nor is it intended to embrace sequence variants which, although they encode different proteins, encode proteins which are functionally equivalent to normal CalDAG-GEF and/or cAMP-GEF proteins.

Functional equivalent. As used herein in describing gene sequences and amino acid sequences, the term "functional equivalent" means that a recited sequence need not be identical to a particularly disclosed sequence of the SEQ ID NOs but need only provide a sequence which functions biologically and/or chemically as the equivalent of the disclosed sequence.

Substantially pure. As used herein with respect to protein preparations, the term "substantially pure" means a preparation which contains at least 60% (by dry weight) the protein

- 16 -

of interest, exclusive of the weight of other intentionally included compounds. Preferably the preparation is at least 75%, more preferably at least 90%, and most preferably at least 99%, by dry weight the protein of interest, exclusive of the weight of other intentionally included compounds. Purity can be measured by any appropriate method, e.g., column chromatography, gel electrophoresis, or HPLC analysis. If a preparation intentionally includes two or more different proteins of the invention, a "substantially pure" preparation means a preparation in which the total dry weight of the proteins of the invention is at least 60% of the total dry weight, exclusive of the weight of other intentionally included compounds. Preferably, for such preparations containing two or more proteins of the invention, the total weight of the proteins of the invention be at least 75%, more preferably at least 90%, and most preferably at least 99%, of the total dry weight of the preparation, exclusive of the weight of other intentionally included compounds. Thus, if the proteins of the invention are mixed with one or more other proteins (e.g., serum albumin, 6-OST) or compounds (e.g., diluents, detergents, excipients, salts, polysaccharides, sugars, lipids) for purposes of administration, stability, storage, and the like, the weight of such other proteins or compounds is ignored in the calculation of the purity of the preparation.

Isolated nucleic acid. As used herein, an "isolated nucleic acid" is a ribonucleic acid, deoxyribonucleic acid, or nucleic acid analog comprising a polynucleotide sequence that has been isolated or separated from sequences that are immediately contiguous (one on the 5' end and one on the 3' end) in the naturally occurring genome of the organism from which it is derived. The term therefore includes, for example, a recombinant nucleic acid which is incorporated into a vector, into an autonomously replicating plasmid or virus, or into the genomic DNA of a prokaryote or eukaryote; or which exists as a separate molecule (e.g., a cDNA or a genomic DNA fragment produced by PCR or restriction endonuclease treatment) independent of other sequences. It also includes a recombinant DNA which is part of a hybrid gene encoding additional polypeptide sequences and/or including exogenous regulatory elements.

Transformed cell. As used herein, a "transformed cell" is a cell into which (or into an ancestor of which) has been introduced, by means of recombinant DNA techniques, a nucleic acid molecule of interest. The nucleic acid of interest will typically encode a peptide or protein. The transformed cell may express the sequence of interest or may be used only to propagate the sequence. The term "transformed" may be used herein to embrace any method of introducing

- 17 -

exogenous nucleic acids including, but not limited to, transformation, transfection, electroporation, microinjection, viral-mediated transfection, and the like.

Operably joined. As used herein, a coding sequence and a regulatory region are said to be "operably joined" when they are covalently linked in such a way as to place the expression or transcription of the coding sequence under the influence or control of the regulatory region. If it is desired that the coding sequences be translated into a functional protein, two DNA sequences are said to be operably joined if induction of promoter function results in the transcription of the coding sequence and if the nature of the linkage between the two DNA sequences does not (1) result in the introduction of a frame-shift mutation, (2) interfere with the ability of the regulatory region to direct the transcription of the coding sequences, or (3) interfere with the ability of the corresponding RNA transcript to be translated into a protein. Thus, a regulatory region would be operably joined to a coding sequence if the regulatory region were capable of effecting transcription of that DNA sequence such that the resulting transcript might be translated into the desired protein or polypeptide.

Stringent hybridization conditions. Stringent hybridization conditions is a term of art understood by those of ordinary skill in the art. For any given nucleic acid sequence, stringent hybridization conditions are those conditions of temperature, chaotropic acids, buffer, and ionic strength which will permit hybridization of that nucleic acid sequence to its complementary sequence and not to substantially different sequences. The exact conditions which constitute "stringent" conditions, depend upon the nature of the nucleic acid sequence, the length of the sequence, and the frequency of occurrence of subsets of that sequence within other non-identical sequences. By varying hybridization conditions from a level of stringency at which non-specific hybridization occurs to a level at which only specific hybridization is observed, one of ordinary skill in the art can, without undue experimentation, determine conditions which will allow a given sequence to hybridize only with complementary sequences. Suitable ranges of such stringency conditions are described in KRAUSE ET AL., METHODS IN ENZYMOLOGY, 200: 546-56 (1991). Stringent hybridization conditions, depending upon the length and commonality of a sequence, may include temperatures of 20°C-65°C and ionic strengths from 5x to 0.1x SSC. Highly stringent hybridization conditions may include temperatures as low as 40-42°C (when denaturants such as formamide are included) or up to 60-65°C in ionic strengths as low as 0.1x SSC. These ranges, however, are only illustrative and, depending upon the nature of the target sequence, and possible future technological developments, may be more stringent than necessary.

- 18 -

Less than stringent conditions are employed to isolate nucleic acid sequences which are substantially similar, allelic or homologous to any given sequence.

Selectively bind. As used herein with respect to antibodies, an antibody is said to "selectively bind" to a target if the antibody recognizes and binds the target of interest but does not substantially recognize and bind other molecules in a sample, *e.g.*, a biological sample, which includes the target of interest.

CalDAG-GEF- or cAMP-GEF-associated disorder, condition, or disease. As used herein, the term "CalDAG-GEF or cAMP-GEF associated disorder, condition, or disease" means any disorder, condition, or disease to which a normal or mutant CalDAG-GEF and/or cAMP-GEF is related in any manner, such as in the causation, prevention, exacerbation, alleviation of the disorder. Thus, as used herein, a CalDAG-GEF- or cAMP-GEF-associated disorder, condition, or disease includes disorders related to the Ras-pathway, such as Ras-related cancers.

Adapter protein. As used herein, the term "adapter protein" means any protein that binds or is bound to a CalDAG-GEF or a cAMP-GEF protein, and facilitates localization of the bound CalDAG-GEF or cAMP-GEF at the plasma membrane, thereby facilitating Ras activation.

Variant. As used herein a "variant" sequence has, or will result in having, a sufficient amino acid similarity to have a reasonable expectation of success in the methods of the present invention. In order to produce variants of the disclosed sequences that may also functionally serve as a CalDAG-GEF or cAMP-GEF protein, any one or more of the naturally-occurring CalDAG-GEF or cAMP-GEF sequences disclosed herein may be used as a reference sequence to determine whether a candidate sequence possesses sufficient amino acid similarity to have a reasonable expectation of success in the methods of the present invention. Preferably, variant sequences are at least 70% similar or 60% identical, more preferably at least 75% similar or 65% identical, and most preferably 80% similar or 70% identical to one of the disclosed, naturally-occurring sequences.

To determine whether a candidate peptide region has the requisite percentage similarity or identity to a reference polypeptide or peptide oligomer, the candidate amino acid sequence and the reference amino acid sequence are first aligned using the dynamic programming algorithm described in Smith and Waterman (1981), *J. Mol. Biol.* 147:195-197, in combination with the BLOSUM62 substitution matrix described in Figure 2 of Henikoff and Henikoff (1992), "Amino acid substitution matrices from protein blocks", *PNAS* (1992 Nov), 89:10915-10919. For the present invention, an appropriate value for the gap insertion penalty is -12, and an appropriate

- 19 -

value for the gap extension penalty is -4. Computer programs performing alignments using the algorithm of Smith-Waterman and the BLOSUM62 matrix, such as the GCG program suite (Oxford Molecular Group, Oxford, England), are commercially available and widely used by those skilled in the art.

5 Once the alignment between the candidate and reference sequence is made, a percent similarity score may be calculated. The individual amino acids of each sequence are compared sequentially according to their similarity to each other. If the value in the BLOSUM62 matrix corresponding to the two aligned amino acids is zero or a negative number, the pairwise similarity score is zero; otherwise the pairwise similarity score is 1.0. The raw similarity score is
10 the sum of the pairwise similarity scores of the aligned amino acids. The raw score is then normalized by dividing it by the number of amino acids in the smaller of the candidate or reference sequences. The normalized raw score is the percent similarity. Alternatively, to calculate a percent identity, the aligned amino acids of each sequence are again compared sequentially. If the amino acids are non-identical, the pairwise identity score is zero; otherwise
15 the pairwise identity score is 1.0. The raw identity score is the sum of the identical aligned amino acids. The raw score is then normalized by dividing it by the number of amino acids in the smaller of the candidate or reference sequences. The normalized raw score is the percent identity. Insertions and deletions are ignored for the purposes of calculating percent similarity and identity. Accordingly, gap penalties are not used in this calculation, although they are used
20 in the initial alignment.

In all instances, variants of the naturally-occurring CalDAG-GEF or cAMP-GEF proteins, as described above, must be tested for biological activity as described below. Specifically, the proteins must exhibit guanine nucleotide exchange factor activity, and, preferably, they have the ability to inhibit Ras signaling of the Ras/Raf-1/MAP kinase pathway.

25 II. The CalDAG-GEFs

CalDAG-GEFI has a substrate specificity for Rap1A, dual binding domains for calcium (Ca^{2+}) and diacylglycerol (DAG), and enriched expression in brain basal ganglia pathways and their axon-terminal regions. Expression of CalDAG-GEFI activates Rap1A and inhibits Ras-dependent activation of the Erk/MAP kinase cascade in 293T cells. Ca^{2+} ionophore and phorbol
30 ester strongly and additively enhance this Rap1A activation. By contrast, CalDAG-GEFII exhibits a different brain expression pattern and fails to activate Rap1A, but activates H-Ras, R-Ras and the Erk/MAP kinase cascade under Ca^{2+} and DAG modulation. The CalDAG-GEF

- 20 -

proteins have a critical neuronal function in determining the relative activation of Ras and Rap1 signaling induced by Ca^{2+} and DAG mobilization. The expression of CalDAG-GEFI and CalDAG-GEFII in hematopoietic organs indicates that such control has broad significance in Ras/Rap regulation of normal and malignant states.

5 The basal ganglia are centrally implicated in movement control and in forms of procedural learning related to habit formation. It is not yet known whether particular neurochemical specializations of the basal ganglia contribute to these behavioral functions. The basal ganglia do, however, have a unique double-inhibitory pathway design combined with abundant expression of neuromodulators in striatal neurons. A number of genes with
10 differentially high expression in the striatum have also been identified. These include genes coding for proteins with signaling functions, such as adenylate cyclase V (Glatt et al., 361 NATURE (LONDON), 536-38 (1993)) and DARPP-32 (Hemmings et al., 310 NATURE (LONDON) 502-05 (1984)). To identify other cellular signaling molecules that could contribute to basal ganglia functions, a search for striatum-enriched transcripts was performed by a differential
15 display method, as discussed in Example 1. Among the transcripts identified in this search were a family of genes characterized by the presence of a Ras superfamily (GEF) domain.

Specific domains identified include structurally conserved GEF regions SCR1, SCR2, and SCR3, as shown in Figures 2C and 3D, and as shown in the following table.

- 21 -

TABLE 1

Gene	SCR1	SCR2	SCR3
hCalDAG-GEFI	SEQ ID NO.3: 605-677 SEQ ID NO.4: 149-173	SEQ ID NO.3: 817-946 SEQ ID NO.4: 219-262	SEQ ID NO.3: 1053-1185 SEQ ID NO.4: 298-320
hCalDAG-GEFII	SEQ ID NO.7: 728-800 SEQ ID NO.8: 205-229	SEQ ID NO.7: 913-1042 SEQ ID NO.8: 270-313	SEQ ID NO.7: 1084-1216 SEQ ID NO.8: 348-371
hcAMP-GEFI	SEQ ID NO.11: 2058-2130 SEQ ID NO.12: 205-229	SEQ ID NO.11: 2276-2405 SEQ ID NO.12: 688-731	SEQ ID NO.11: 2516-2582 SEQ ID NO.12: 767-789
rcAMP-GEFI	SEQ ID NO.9: 2050-2122 SEQ ID NO.10: 618-642	SEQ ID NO.9: 2267-2396 SEQ ID NO.10: 691-734	SEQ ID NO.9: 2502-2568 SEQ ID NO.10: 770-792
hcAMP-GEFII	SEQ ID NO.17: 2707-2779 SEQ ID NO.18: 767-791		
rcAMP-GEFII	SEQ ID NO.15: 576-648 SEQ ID NO.16: 192-216		

In addition, the EF hand and DAG-binding domains were identified as shown in Figures 2E and 2F, and as shown in the following table:

- 22 -

Table 2

Gene	EF Hand Domain	DAG-Binding Domain
hCalDaG-GEFI	SEQ ID NO.3: 1456-1516 SEQ ID NO.4: 432-452	SEQ ID NO.3: 1652-1804 SEQ ID NO.4: 498-548
hCalDAG-GEFII	SEQ ID NO.7: 1384-1444 SEQ ID NO.8: 427-447	SEQ ID NO.7: 1579-1729 SEQ ID NO.8: 492-542

Finally, the cAMP-binding domains were identified as shown in Figure 3E, and as shown in the following table:

Table 3

Gene	cAMP-Binding Domain
hcAMP-GEFI	SEQ ID NO.11: 2012-2255 SEQ ID NO.12: 219-300
rcAMP-GEFI	SEQ ID NO.9: 853-1096 SEQ ID NO.10: 219-300
rcAMP-GEFII	SEQ ID NO.17: 1522-1765 SEQ ID NO.18: 372-453

III. The cAMP-GEFs

Cyclic adenosine 3', 5'-monophosphate (cAMP) is a universal second messenger that induces a variety of physiological responses in eukaryotic cells ranging from growth, differentiation, and gene expression to secretion and neurotransmission. The cAMP second messenger system has also been centrally implicated in modulating synaptic function, neuroplasticity and learning and memory. Most of these effects have been attributed to the binding of cAMP to cAMP-dependent protein kinase (PKA), leading in turn to the activation of

- 23 -

intracellular phosphorylation cascades. Reported herein is the identification of a new family of cAMP binding proteins that are differentially distributed in the brain and body organs and that are characterized by the presence of both a cAMP binding domain and a guanine nucleotide exchange factor (GEF) domain. These proteins, cAMP-GEFs, bind cAMP and selectively
5 activate the Ras superfamily small G protein, Rap1A, in a cAMP-dependent but PKA-independent manner.

The general concept of cAMP signaling involves the sequential activation (or inhibition) of cAMP production by G proteins, the binding of cAMP to PKA, and the triggering of a series of downstream serine-threonine phosphorylation cascades. Viewed as the nearly exclusive
10 target of cAMP binding in eukaryotic cells, PKA has been considered the essential effector molecule mediating a wide range of physiological effects of G protein/cAMP-triggered phosphorylation cascades. As the main cAMP effector, PKA has also been shown to function in the indirect coupling of the cAMP signal transduction system to other intracellular signaling cascades. The cAMP signaling system has also been strongly implicated in neuronal functions
15 ranging from neurotransmitter-initiated signaling to neuroplasticity underlying development and memory, but PKA has not been clearly linked to all of these neuronal functions, and region-specific neuronal effects have been observed as well. The cAMP-GEF gene has a Ras superfamily GEF motif. Thus, the gene codes for a novel cAMP binding protein that directly couples the cAMP signal transduction system to Ras superfamily cascades.

20 IV. Preferred Embodiments

Based, in part, upon the discoveries disclosed and described herein, the following preferred embodiments of the present invention are provided.

1. Isolated Nucleic Acids

In one series of embodiments, the present invention provides isolated nucleic acids
25 corresponding to, or relating to, the CalDAG-GEF or cAMP-GEF nucleic acid sequences disclosed herein. As described more fully below, these sequences include normal CalDAG-GEF and cAMP-GEF sequences from humans and other mammalian species, mutant CalDAG-GEF and cAMP-GEF sequences from humans and other mammalian species, homologous sequences from non-mammalian species such as *Drosophila* and *C. elegans*, subsets of these sequences
30 useful as probes and PCR primers, subsets of these sequences encoding fragments of the CalDAG-GEF or cAMP-GEF proteins or corresponding to particular domains or polymorphic

- 24 -

regions, complementary or antisense sequences corresponding to fragments of the CalDAG-GEF or cAMP-GEF genes, sequences in which the CalDAG-GEF and/or cAMP-GEF coding regions have been operably joined to exogenous regulatory regions, and sequences encoding fusion proteins of the portions of the CalDAG-GEF or cAMP-GEF proteins fused to other proteins useful as markers of expression, as "tags" for purification, or in screens and assays for proteins interacting with the CalDAG-GEFs and/or cAMP-GEFs.

Thus, in a first series of embodiments, isolated nucleic acid sequences are provided which encode normal versions of the CalDAG-GEF and cAMP-GEF proteins. Examples of such nucleic acid sequences are disclosed herein. These nucleic acids may be genomic sequences or may be cDNA sequences (*e.g.*, SEQ ID NOs: 1, 3, 5, 7, 9, 11, 13, 15, and 17). Thus, for example, the invention provides nucleic acid sequences in which the alternative splice variants described herein are incorporated at the DNA level, thereby, enabling cells including these sequences to express only one of the alternative splice variants at each splice position. For example, a recombinant gene may be produced in which one of the splice variants of cAMP-GEF is incorporated into DNA such that cells having this recombinant gene can express only one of these variants. For purposes of reducing the size of a recombinant CalDAG-GEF or cAMP-GEF gene, a cDNA gene may be employed or various combinations of the introns and untranslated exons may be removed from a DNA construct. Such constructs may be particularly useful, as described below, in identifying compounds which can induce or repress the expression of the CalDAG-GEF or cAMP-GEF genes.

In addition to the disclosed CalDAG-GEF and/or cAMP-GEF sequences, one of ordinary skill in the art is now enabled to identify and isolate nucleic acids corresponding to CalDAG-GEF or cAMP-GEF genes or cDNAs which are allelic to the disclosed sequences or which are heterospecific homologues. Thus, the present invention provides isolated nucleic acids corresponding to these alleles and homologues, as well as various recombinant constructs derived from these sequences, by means which are well known in the art. Briefly, one of ordinary skill in the art may now screen preparations of genomic or cDNA, including samples prepared from individual organisms (*e.g.*, human cancer patients or their family members) as well as bacterial, viral, yeast or other libraries of genomic or cDNA, using probes or PCR primers to identify allelic or homologous sequences. Because it is desirable to identify additional CalDAG-GEF and/or cAMP-GEF gene mutations which may contribute to the development of Ras-related cancers, because it is desirable to identify additional CalDAG-GEF and/or cAMP-

- 25 -

GEF polymorphisms which are not mutant or have antitumorigenic effects, and because it is also desired to create a variety of animal models which may be used to study Ras-related cancers and screen for potential therapeutics, it is particularly contemplated that additional CalDAG-GEF and/or cAMP-GEF sequences will be isolated from other preparations or libraries of human
5 nucleic acids and from preparations or libraries from animals including rats, mice, hamsters, guinea pigs, rabbits, dogs, cats, goats, sheep, pigs, and non-human primates. Furthermore, CalDAG-GEF and/or cAMP-GEF homologues from yeast or invertebrate species, including *C. elegans* and other nematodes, as well as *Drosophila* and other insects, may have particular utility for drug screening. For example, invertebrates bearing mutant CalDAG-GEF and/or cAMP-GEF
10 homologues (or mammalian CalDAG-GEF and/or cAMP-GEF transgenes) which cause a rapidly occurring and easily scored phenotype (*e.g.*, abnormal eye development after several days) can be used as screens for drugs which block the effect of the mutant gene. Such invertebrates may prove far more rapid and efficient for mass screenings than larger vertebrate animals. Once lead compounds are found through such screens, they may be tested in higher animals.

15 Depending upon the intended use, the present invention provides nucleic acid subsequences of the CalDAG-GEF and/or cAMP-GEF genes which may have lengths varying from 8-10 nucleotides (*e.g.*, for use as PCR primers) to nearly the full size of the CalDAG-GEF and/or cAMP-GEF genes. Thus, the present invention provides isolated nucleic acids comprising sequences corresponding to at least 8, preferably at least 10, and more preferably at least 15
20 consecutive nucleotides of the CalDAG-GEF and/or cAMP-GEF genes, as disclosed or otherwise enabled herein, or to their complements.

In another series of embodiments, the present invention provides for isolated nucleic acids encoding all or a portion of the CalDAG-GEF and/or cAMP-GEF proteins in the form of a fusion protein. In these embodiments, a nucleic acid regulatory region (endogenous or
25 exogenous) is operably joined to a first coding region which is covalently joined in-frame to a second coding region. The CalDAG-GEF and/or cAMP-GEF sequences of the fusion protein may represent the first, second, or any additional coding regions. The CalDAG-GEF and/or cAMP-GEF sequences may be conserved or non-conserved domains and can be placed in any coding region for the fusion protein.

30 In another series of embodiments, the present invention provides isolated nucleic acids in the form of recombinant DNA constructs in which a marker or reporter gene (*e.g.*, β -galactosidase, luciferase) is operably joined to the 5' regulatory region of a CalDAG-GEF and/or

- 26 -

cAMP-GEF gene such that expression of the marker gene is under the control of the CalDAG-GEF and/or cAMP-GEF regulatory sequences. Such isolated nucleic acids may be used to produce cells, cell lines or transgenic animals which are useful in the identification of compounds which can, directly or indirectly, differentially affect the expression of the CalDAG-GEFs and/or cAMP-GEFs.

Finally, the isolated nucleic acids of the present invention include any of the above described sequences when included in vectors. Appropriate vectors include cloning vectors and expression vectors of all types, including plasmids, phagemids, cosmids, episomes, and the like, as well as integration vectors. The vectors may also include various marker genes (*e.g.*, antibiotic resistance or susceptibility genes) which are useful in identifying cells successfully transformed therewith. In addition, the vectors may include regulatory sequences to which the nucleic acids of the invention are operably joined, and/or may also include coding regions such that the nucleic acids of the invention, when appropriately ligated into the vector, are expressed as fusion proteins. Such vectors may also include vectors for use in yeast "two hybrid," baculovirus, and phage-display systems.

2. Substantially Pure Proteins

The present invention provides for substantially pure preparations of the CalDAG-GEF and/or cAMP-GEF proteins, fragments of the CalDAG-GEF and/or cAMP-GEF proteins, and fusion proteins including the CalDAG-GEFs and/or cAMP-GEFs or fragments thereof. The proteins, fragments and fusions have utility, as described herein, in the generation of antibodies to normal and mutant CalDAG-GEFs and/or cAMP-GEFs, in the identification of CalDAG-GEF and/or cAMP-GEF binding proteins, and in diagnostic and therapeutic methods. Therefore, depending upon the intended use, the present invention provides substantially pure proteins or peptides comprising amino acid sequences which are subsequences of the complete CalDAG-GEF and/or cAMP-GEF proteins and which may have lengths varying from 4-8 amino acids (*e.g.*, for use as immunogens), or 9-15 amino acids (*e.g.*, for use in binding assays), to the complete CalDAG-GEF and/or cAMP-GEF proteins. Thus, the present invention provides substantially pure proteins or peptides comprising sequences corresponding to at least 4, preferably at least 9, more preferably at least 15 consecutive amino acids of the CalDAG-GEF and/or cAMP-GEF proteins, as disclosed or otherwise enabled herein.

Purification can be achieved using standard protein purification procedures including, but not limited to, gel-filtration chromatography, ion-exchange chromatography, high-performance

- 27 -

liquid chromatography (RP-HPLC, ion-exchange HPLC, size-exclusion HPLC, high-performance chromatofocusing chromatography, hydrophobic interaction chromatography, immunoprecipitation, or immunoaffinity purification. Gel electrophoresis (*e.g.*, PAGE, SDS-PAGE) can also be used to isolate a protein or peptide based on its molecular weight, charge properties, and hydrophobicity.

A CalDAG-GEF or cAMP-GEF protein, or a fragment thereof, may also be conveniently purified by creating a fusion protein including the desired CalDAG-GEF or cAMP-GEF sequence fused to another peptide such as an antigenic determinant or poly-His tag (*e.g.*, QIAexpress vectors, (QIAGEN Corp., Chatsworth, CA)), or a larger protein (*e.g.*, GST using the pGEX-27 vector (Amrad, USA) or green fluorescent protein using the Green Lantern vector (GIBCO/BRL, Gaithersburg, MD)).

3. Antibodies to the CalDAG-GEF and/or cAMP-GEFs

The present invention also provides antibodies, and methods of making antibodies, which selectively bind to the CalDAG-GEF and/or cAMP-GEF proteins or fragments thereof. The antibodies of the invention have utility as laboratory reagents for, *inter alia*, immunoaffinity purification of the CalDAG-GEFs and/or cAMP-GEFs, Western blotting to identify cells or tissues expressing the CalDAG-GEFs and/or cAMP-GEFs, and immunocytochemistry or immunofluorescence techniques to establish the subcellular location of the protein.

The antibodies of the invention may be generated in a host using the entire CalDAG-GEF and/or cAMP-GEF proteins of the invention or using any CalDAG-GEF and/or cAMP-GEF epitope which is characteristic of that protein and which substantially distinguishes it from host proteins. Such epitopes may be identified by comparing sequences of, for example, 4-8 amino acid residues from a CalDAG-GEF and/or cAMP-GEF sequence to computer databases of protein sequences from the relevant host. Antibodies against highly conserved domains are expected to have the greatest utility for purification or identification of CalDAG-GEFs and/or cAMP-GEFs.

Amino acid residue positions which are potential antigenic sites in the CalDAG-GEF or cAMP-GEF proteins and which may be useful in generating the antibodies of the invention may be determined by using computer programs such as the IBI Pustell program. Other methods of choosing antigenic determinants are known in the art and may, of course, be employed. In addition, larger fragments (*e.g.*, 9-15 residues) including some of these epitopes may also be employed. Even larger fragments, including, for example, entire functional domains or multiple

- 28 -

functional domains may also be preferred. For an overview of antibody techniques, see Antibody Engineering: A Practical Guide, Borrebaek, ed., W.H. Freeman & Company, NY (1992), or Antibody Engineering, 2nd Ed., Borrebaek, ed., Oxford University Press, Oxford (1995).

The antibodies of the invention may be labelled or conjugated with other compounds or materials for diagnostic and/or therapeutic uses. For example, they may be coupled to radionuclides, fluorescent compounds, or enzymes for imaging or therapy, or to liposomes for the targeting of compounds contained in the liposomes to a specific tissue location.

4. Transformed Cell Lines

The present invention also provides for cells or cell lines, both prokaryotic and eukaryotic, which have been transformed or transfected with the nucleic acids of the present invention so as to cause clonal propagation of those nucleic acids and/or expression of the proteins or peptides encoded thereby. Such cells or cell lines will have utility not only in the propagation and production of the nucleic acids and proteins of the present invention but also, as further described herein, as model systems for diagnostic and therapeutic assays. As used herein, the term "transformed cell" is intended to embrace any cell, or the descendant of any cell, into which has been introduced any of the nucleic acids of the invention, whether by transformation, transfection, infection, electroporation, microinjection or other means. Methods of producing appropriate vectors, transforming cells with those vectors, and identifying transformants are well known in the art and are only briefly reviewed here (see, for example, Sambrook et al. (1989) Molecular Cloning: A Laboratory Manual, 2nd ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York).

Vectors may be introduced into the recipient or "host" cells by various methods well known in the art including, but not limited to, calcium phosphate transfection, strontium phosphate transfection, DEAE dextran transfection, electroporation, lipofection (e.g., Dosper Liposomal transfection reagent, Boehringer Mannheim, Germany), microinjection, ballistic insertion on micro-beads, protoplast fusion or, for viral or phage vectors, by infection with the recombinant virus or phage.

5. Transgenic Animal Models

The present invention also provides for the production of transgenic non-human animal models for the study of Ras-related cancers, for the screening of candidate pharmaceutical compounds, for the creation of explanted mammalian cell cultures (e.g., neuronal, glial,

organotypic or mixed cell cultures) in which mutant or wild type CalDAG-GEF and/or cAMP-GEF sequences are expressed or in which the CalDAG-GEF and/or cAMP-GEF genes have been inactivated (*e.g.*, "knock-out" deletions), and for the evaluation of potential therapeutic interventions.

5 Species suitable for use as animal models in the present invention include, but are not limited to, rats, mice, hamsters, guinea pigs, rabbits, dogs, cats, goats, sheep, pigs, and non-human primates (*e.g.*, Rhesus monkeys, chimpanzees).

Various techniques for generating transgenic animals, as well as techniques for homologous recombination or gene targeting, are now widely accepted and practiced. See, for
10 example, Hogan et al., Manipulating Mouse Embryo (1986). To create a transgene, the target sequence of interest (*e.g.*, mutant or wild-type CalDAG-GEF or cAMP-GEF sequences) is typically ligated into a cloning site located downstream of a promoter element which will regulate the expression of RNA from the CalDAG-GEF or cAMP-GEF sequence. An alternate approach to creating a transgene is to use endogenous CalDAG-GEF or cAMP-GEF regulatory
15 sequences to drive expression of the CalDAG-GEF or cAMP-GEF transgene.

6. Assays for Drugs Which Affect CalDAG-GEF and/or cAMP-GEF Expression

In another series of embodiments, the present invention provides assays for identifying small molecules or other compounds which are capable of inducing or inhibiting the expression of the CalDAG-GEF or cAMP-GEF genes and proteins. The assays may be performed *in vitro*
20 using non-transformed cells, immortalized cell lines, or recombinant cell lines, or *in vivo* using the transgenic animal models enabled herein.

In particular, the assays may detect the presence of increased or decreased expression of CalDAG-GEF, cAMP-GEF, or other CalDAG-GEF or cAMP-GEF-related genes or proteins, on the basis of increased or decreased mRNA expression (using, *e.g.*, the nucleic acid probes
25 disclosed and enabled herein), increased or decreased levels of CalDAG-GEF, cAMP-GEF or other CalDAG-GEF or cAMP-GEF-related protein products (using, *e.g.*, the anti-CalDAG-GEF or anti-cAMP-GEF antibodies disclosed and enabled herein), or increased or decreased levels of expression of a marker gene (*e.g.*, β -galactosidase or luciferase) operably joined to a CalDAG-GEF or cAMP-GEF 5' regulatory region in a recombinant construct.

30 Thus, for example, one may culture cells known to express a particular CalDAG-GEF or cAMP-GEF and add to the culture medium one or more test compounds. After allowing a sufficient period of time (*e.g.*, 0-72 hours) for the compound to induce or inhibit the expression

- 30 -

of the CalDAG-GEF or cAMP-GEF, any change in levels of expression from an established baseline may be detected using any of the techniques described above and well known in the art. In particularly preferred embodiments, the cells are from an immortalized cell line such as a human neuroblastoma, glioblastoma or a hybridoma cell line. Using the nucleic acid probes and/or antibodies disclosed and enabled herein, detection of changes in the expression of a CalDAG-GEF or cAMP-GEF and thus, identification of the compound as an inducer or repressor of CalDAG-GEF and/or cAMP-GEF expression, requires only routine experimentation.

In particularly preferred embodiments, a recombinant assay is employed in which a reporter gene such as a β -galactosidase, green fluorescent protein, alkaline phosphatase, or luciferase is operably joined to a 5' regulatory region of a CalDAG-GEF or cAMP-GEF gene. The reporter gene and regulatory regions are joined in-frame (or in each of the three possible reading frames) so that transcription and translation of the reporter gene may proceed under the control of the CalDAG-GEF or cAMP-GEF regulatory elements. The recombinant construct may then be introduced into any appropriate cell type, although mammalian cells are preferred, and human cells are most preferred. The transformed cells may be grown in culture and, after establishing the baseline level of expression of the reporter gene, test compounds may be added to the medium. The ease of detection of the expression of the reporter gene provides for a rapid, high through-put assay for the identification of inducers and repressors of the CalDAG-GEF or cAMP-GEF gene.

Compounds identified by this method will have potential utility in modifying the expression of the CalDAG-GEF, cAMP-GEF or other CalDAG-GEF or cAMP-GEF-related genes *in vivo*. These compounds may be further tested in the animal models disclosed and enabled herein to identify those compounds having the most potent *in vivo* effects. In addition, as described herein with respect to small molecules having CalDAG-GEF or cAMP-GEF-binding activity, these molecules may serve as "lead compounds" for the further development of pharmaceuticals by, for example, subjecting the compounds to sequential modifications, molecular modeling, and other routine procedures employed in rational drug design.

7. Identification of Compounds with CalDAG-GEF and/or cAMP-GEF Binding Capacity

In light of the present disclosure, one of ordinary skill in the art is enabled to practice new screening methodologies which will be useful in the identification of proteins and other compounds which bind to, or otherwise directly interact with, the CalDAG-GEFs or cAMP-GEFs. The proteins and compounds will include endogenous cellular components which interact

- 31 -

with the CalDAG-GEFs or cAMP-GEFs *in vivo* and which, therefore, provide new targets for pharmaceutical and therapeutic interventions, as well as recombinant, synthetic and otherwise exogenous compounds which may have CalDAG-GEF or cAMP-GEF binding capacity and, therefore, may be candidates for pharmaceutical agents. Thus, in one series of embodiments, cell
5 lysates or tissue homogenates (*e.g.*, human brain homogenates, leukocyte lysates) may be screened for proteins or other compounds which bind to one of the normal or mutant CalDAG-GEFs and/or cAMP-GEFs. Alternatively, any of a variety of exogenous compounds, both naturally occurring and/or synthetic (*e.g.*, libraries of small molecules or peptides), may be screened for CalDAG-GEF or cAMP-GEF binding capacity. Small molecules are particularly
10 preferred in this context because they are more readily absorbed after oral administration, have fewer potential antigenic determinants, and/or are more likely to cross the blood brain barrier than larger molecules such as nucleic acids or proteins. The methods of the present invention are particularly useful in that they may be used to identify molecules which selectively or preferentially bind to a mutant form of a CalDAG-GEF or cAMP-GEF protein (rather than a
15 normal form) and, therefore, may have particular utility in treating the heterozygous victims of a CalDAG-GEF or cAMP-GEF associated disorder.

Compounds which bind to normal, mutant or both forms of the CalDAG-GEFs or cAMP-GEFs may have utility in treatments and diagnostics. Compounds which bind only to a normal CalDAG-GEF or cAMP-GEF may, for example, act as enhancers of its normal activity and
20 thereby at least partially compensate for the lost or abnormal activity of mutant forms of the CalDAG-GEF or cAMP-GEF in victims suffering from CalDAG-GEF- or cAMP-GEF-associated disorders. Compounds which bind to both normal and mutant forms of a CalDAG-GEF or cAMP-GEF may have utility if they differentially affect the activities of the two forms so as to alleviate the overall departure from normal function. Alternatively, blocking the activity of
25 both normal and mutant forms of either CalDAG-GEF or cAMP-GEF may have less severe physiological and clinical consequences than the normal progress of the disorder and, therefore, compounds which bind to and inhibit the activity of both normal and mutant forms of a CalDAG-GEF or cAMP-GEF may be therapeutically useful. Preferably, however, compounds are identified which have a higher affinity of binding to mutant CalDAG-GEF or cAMP-GEF
30 than to normal CalDAG-GEF or cAMP-GEF, and which selectively or preferentially inhibit the activity of the mutant form. Such compounds may be identified by using any of the techniques

- 32 -

described herein, and then comparing the binding affinities of the candidate compound(s) for the normal and mutant forms of CalDAG-GEF or cAMP-GEF.

The effect of agents which bind to the CalDAG-GEFs or cAMP-GEFs (normal or mutant forms of either) can be monitored either by direct monitoring of this binding (*e.g.*, using the BIAcore assay, LKB Pharmacia, Sweden) or by indirect monitoring of binding by detecting, for example, a change in fluorescence, molecular weight, or concentration of either the binding agent or CalDAG-GEF or cAMP-GEF component, either in a soluble phase or in a substrate-bound phase.

Once identified by the methods described above, the candidate compounds may then be produced in quantities sufficient for pharmaceutical administration or testing (*e.g.*, μg or mg or greater quantities), and formulated in a pharmaceutically acceptable carrier (see, *e.g.*, REMINGTON'S PHARMACEUTICAL SCIENCES, Gennaro, A., ed., Mack Pub., (1990)). These candidate compounds may then be administered to the transformed cells of the invention, to the transgenic animal models of the invention, to cell lines derived from the animal models or from human patients, or to patients with CalDAG-GEF- or cAMP-GEF-associated disorders. The animal models described and enabled herein are of particular utility in further testing candidate compounds which bind to normal or mutant CalDAG-GEF or cAMP-GEF for their therapeutic efficacy.

In addition, once identified by the methods described above, the candidate compounds may also serve as "lead compounds" in the design and development of new pharmaceuticals. For example, as is well known in the art, sequential modification of small molecules (*e.g.*, amino acid residue replacement for peptides; functional group replacement for peptide or non-peptide compounds) is a standard approach in the pharmaceutical industry for the development of new pharmaceuticals. Such development generally proceeds from a "lead compound" which is shown to have at least some of the activity (*e.g.*, CalDAG-GEF or cAMP-GEF binding or blocking ability) of the desired pharmaceutical. In particular, when one or more compounds having at least some activity of interest (*e.g.*, modulation of CalDAG-GEF or cAMP-GEF activity) are identified, structural comparison of the molecules can greatly inform the skilled practitioner by suggesting portions of the lead compounds which should be conserved and portions which may be varied in the design of new candidate compounds. Thus, the present invention also provides a means of identifying lead compounds which may be sequentially modified to produce new candidate compounds for use in the treatment of CalDAG-GEF- or cAMP-GEF-associated

- 33 -

disorders. These new compounds then may be tested both for CalDAG-GEF or cAMP-GEF-binding or blocking (*e.g.*, in the binding assays described above) and for therapeutic efficacy (*e.g.*, in the animal models described herein). This procedure may be iterated until compounds having the desired therapeutic activity and/or efficacy are identified.

5 In each of the present series of embodiments, an assay is conducted to detect binding between a "CalDAG-GEF component" or a "cAMP-GEF component" and some other moiety. Of particular utility will be sequential assays in which compounds are tested for the ability to bind to only the normal or only the mutant forms of the CalDAG-GEF or cAMP-GEF functional domains using mutant and normal CalDAG-GEF or cAMP-GEF components in the binding
10 assays. Such compounds are expected to have the greatest therapeutic utilities, as described more fully below. The "CalDAG-GEF component" or the "cAMP-GEF component" in these assays may be a complete normal or mutant form of a CalDAG-GEF or cAMP-GEF protein (*e.g.*, an hCalDAG-GEF or hcAMP-GEF variant) but need not be. Rather, particular functional domains of the CalDAG-GEFs or cAMP-GEFs, as described above, may be employed either as
15 separate molecules or as part of a fusion protein. For example, to isolate proteins or compounds that interact with these functional domains, screening may be carried out using fusion constructs and/or synthetic peptides corresponding to these regions. Obviously, various combinations of fusion proteins and functional domains from CalDAG-GEF or cAMP-GEF are possible. In addition, the functional domains may be altered so as to aid in the assay by, for example,
20 introducing into the functional domain a reactive group or amino acid residue (*e.g.*, cysteine) which will facilitate immobilization of the domain on a substrate (*e.g.*, using sulfhydryl reactions).

Methods for screening cellular lysates, tissue homogenates, or small molecule libraries for candidate CalDAG-GEF or cAMP-GEF-binding molecules are well known in the art and, in
25 light of the present disclosure, may now be employed to identify compounds which bind to normal or mutant CalDAG-GEF or cAMP-GEF components or which modulate CalDAG-GEF or cAMP-GEF activity as defined by non-specific measures (*e.g.*, changes in intracellular Ca^{2+} , GTP/GDP ratio) or by specific measures (*e.g.*, changes in the expression of other downstream genes which can be monitored by differential display, 2D gel electrophoresis, differential
30 hybridization, or SAGE methods). The preferred methods involve variations on the following techniques: (1) direct extraction by affinity chromatography; (2) co-isolation of CalDAG-GEF or cAMP-GEF components and bound proteins or other compounds by immunoprecipitation; (3)

- 34 -

the Biomolecular Interaction Assay (BIAcore); and (4) the yeast two-hybrid systems. These and others are discussed separately below.

A. Affinity Chromatography

In light of the present disclosure, a variety of affinity binding techniques well known in the art may be employed to isolate proteins or other compounds which bind to the CalDAG-GEFs or cAMP-GEFs disclosed or otherwise enabled herein. In general, a CalDAG-GEF or cAMP-GEF component may be immobilized on a substrate (*e.g.*, a column or filter) and a solution including the test compound(s) is contacted with the CalDAG-GEF or cAMP-GEF protein, fusion or fragment under conditions which are permissive for binding. The substrate is then washed with a solution to remove unbound or weakly bound molecules. A second wash may then elute those compounds which strongly bound to the immobilized normal or mutant CalDAG-GEF or cAMP-GEF component. Alternatively, the test compounds may be immobilized and a solution containing one or more CalDAG-GEF or cAMP-GEF components may be contacted with the column, filter, or other substrate. The ability of the CalDAG-GEF or cAMP-GEF component to bind to the test compounds may be determined as above or a labeled form of the CalDAG-GEF or cAMP-GEF component (*e.g.*, a radio-labeled or chemiluminescent functional domain) may be used to more rapidly assess binding to the substrate-immobilized compound(s).

B. Co-Immunoprecipitation

Another well characterized technique for the isolation of the CalDAG-GEF or cAMP-GEF components and their associated proteins or other compounds is direct immunoprecipitation with antibodies. This procedure has been successfully used, for example, to isolate many of the synaptic vesicle associated proteins (Phizicky et al., 59 J. BIOL. CHEM. 94-123 (1994)). Thus, either normal or mutant CalDAG-GEF or cAMP-GEF components may be mixed in a solution with the candidate compound(s) under conditions which are permissive for binding, and the CalDAG-GEF or cAMP-GEF component may be immunoprecipitated. Proteins or other compounds which co-immunoprecipitate with the CalDAG-GEF or cAMP-GEF component may then be identified by standard techniques as described above. General techniques for immunoprecipitation may be found in, for example, Harlow et al., ANTIBODIES: A LABORATORY MANUAL (1988).

- 35 -

The antibodies employed in this assay, as described and enabled herein, may be polyclonal or monoclonal, and include the various antibody fragments as well as single chain antibodies, and the like.

C. The Biomolecular Interaction Assay

5 Another useful method for the detection and isolation of binding proteins is the Biomolecular Interaction Assay or "BIAcore" system developed by Pharmacia Biosensor and described in the manufacturer's protocol (LKB Pharmacia, Sweden). In light of the present disclosure, one of ordinary skill in the art is now enabled to employ this system, or a substantial equivalent, to identify proteins or other compounds having CalDAG-GEF or cAMP-GEF binding
10 capacity. The BIAcore system uses an affinity purified anti-GST antibody to immobilize GST-fusion proteins onto a sensor chip. Obviously, other fusion proteins and corresponding antibodies may be substituted. The sensor utilizes surface plasmon resonance which is an optical phenomenon that detects changes in refractive indices. A homogenate of a tissue of interest is passed over the immobilized fusion protein and protein-protein interactions are registered as
15 changes in the refractive index. This system can be used to determine the kinetics of binding and to assess whether any observed binding is of physiological relevance.

D. The Yeast Two-Hybrid System

The yeast "two-hybrid" system takes advantage of transcriptional factors that are composed of two physically separable, functional domains. One commonly used system employs
20 the yeast GAL4 transcriptional activator, consisting of a DNA binding domain and a transcriptional activation domain. Two different cloning vectors are used to generate separate fusions of the GAL4 domains to genes encoding potential binding proteins. The fusion proteins are co-expressed, targeted to the nucleus and, if interactions occur, activation of a reporter gene (*e.g.*, lacZ) produces a detectable phenotype.

25 E. Other Methods

The nucleotide sequences and protein products, including both mutant and normal forms of these nucleic acids and their corresponding proteins, can be used with the above techniques to isolate other interacting proteins, and to identify other genes whose expression is altered by the over-expression of normal CalDAG-GEF or cAMP-GEF sequences, by the under-expression of
30 normal CalDAG-GEFs or cAMP-GEFs sequences, or by the expression of mutant CalDAG-GEF and/or cAMP-GEF sequences. Identification of these interacting proteins, as well as the identification of other genes whose expression levels are altered in the face of mutant CalDAG-

- 36 -

GEF or cAMP-GEF sequences (for instance) will identify other gene targets which have direct relevance to the pathogenesis of this disease in its clinical or pathological forms. Specifically, these techniques rely on PCR-based and/or hybridization-based methods to identify genes which are differentially expressed between two conditions (a cell line expressing normal CalDAG-GEFs or cAMP-GEFs compared to the same cell type expressing a mutant CalDAG-GEF or cAMP-GEF sequence). These techniques include differential display, serial analysis of gene expression (SAGE), mass-spectrometry of protein, 2D-gels and subtractive hybridization (*See, e.g.,* Nowak, 270 Sci. 368-371 (1995); Kahn, 270 Sci. 369-370 (1995)).

8. Methods of Identifying Compounds Modulating CalDAG-GEF and/or cAMP-GEF Activity

In another series of embodiments, the present invention provides for methods of identifying compounds with the ability to modulate the activity of normal and mutant CalDAG-GEFs and/or cAMP-GEFs. As used with respect to this series of embodiments, the term "activity" broadly includes gene and protein expression, CalDAG-GEF and/or cAMP-GEF protein post-translation processing, trafficking and localization, and any functional activity (*e.g.,* enzymatic, receptor-effector, binding, channel), as well as downstream affects of any of these. Using the transformed cells and transgenic animal models of the present invention, cells obtained from subjects bearing a mutant CalDAG-GEF and/or cAMP-GEF gene, or animals or human subjects bearing naturally occurring CalDAG-GEF and/or cAMP-GEF mutations, it is now possible to screen candidate pharmaceuticals and treatments for their therapeutic effects by detecting changes in one or more of the functional characteristics or phenotypic manifestations of normal or mutant CalDAG-GEF and/or cAMP-GEF expression.

Thus, the present invention provides methods for screening or assaying for proteins, small molecules or other compounds which modulate CalDAG-GEF and/or cAMP-GEF activity by contacting a cell *in vivo* or *in vitro* with a candidate compound and assaying for a change in a marker associated with normal or mutant CalDAG-GEF and/or cAMP-GEF activity. The marker associated with CalDAG-GEF and/or cAMP-GEF activity may be any measurable biochemical, physiological, histological and/or behavioral characteristic associated with CalDAG-GEF and/or cAMP-GEF expression. In particular, useful markers will include any measurable biochemical, physiological, histological and/or behavioral characteristic which distinguishes cells, tissues, animals or individuals bearing at least one mutant CalDAG-GEF and/or cAMP-GEF gene from their normal counterparts. In addition, the marker may be any specific or non-specific measure of CalDAG-GEF and/or cAMP-GEF activity, such as the GDP/GTP bound to Rap1/Ras.

- 37 -

CalDAG-GEF and/or cAMP-GEF specific measures include measures of CalDAG-GEF and/or cAMP-GEF expression (*e.g.*, CalDAG-GEF and/or cAMP-GEF mRNA or protein levels) which may employ the nucleic acid probes or antibodies of the present invention. Non-specific measures include changes in cell physiology such as pH, intracellular calcium, cAMP levels, overall GTP/GDP ratios, phosphatidylinositol activity, protein phosphorylation, etc.; which can be monitored by known methods. The activation or inhibition of CalDAG-GEF or cAMP-GEF activity in its mutant or normal form can also be monitored by examining changes in the expression of other genes which are specific to the CalDAG-GEF and/or cAMP-GEF pathway. These can be assayed by such techniques as differential display, differential hybridization, and SAGE, as well as by 2-D gel electrophoresis of cellular lysates. In each case, the differentially-expressed genes can be ascertained by inspection of identical studies before and after application of the candidate compound. Furthermore, as noted elsewhere, the particular genes whose expression is modulated by the administration of the candidate compound can be ascertained by cloning, nucleotide sequencing, amino acid sequencing, or mass spectrometry.

In general, a cell may be contacted with a candidate compound and, after an appropriate period (*e.g.*, 0-72 hours for most biochemical measures of cultured cells), the marker of CalDAG-GEF or cAMP-GEF activity may be assayed and compared to a baseline measurement. The baseline measurement may be made prior to contacting the cell with the candidate compound or may be an external baseline established by other experiments or known in the art. The cell may be a transformed cell of the present invention or an explant from an animal or individual. In particular, the cell may be an explant from a carrier of a CalDAG-GEF or cAMP-GEF mutation or an animal model of the invention (*e.g.*, a transgenic nematode or mouse bearing a mutant CalDAG-GEF or cAMP-GEF gene). Preferred cells include those from neurological tissues such as neuronal, glial or mixed cell cultures; and cultured fibroblasts, liver, kidney, spleen, or bone marrow. The cells may be contacted with the candidate compounds in a culture *in vitro* or may be administered *in vivo* to a live animal or human subject. For live animals or human subjects, the test compound may be administered orally or by any parenteral route suitable to the compound. For clinical trials of human subjects, measurements may be conducted periodically (*e.g.*, daily, weekly or monthly) for several months or years.

In light of the identification, characterization, and disclosure herein of the CalDAG-GEF or cAMP-GEF genes and proteins, the CalDAG-GEF or cAMP-GEF nucleic acid probes and antibodies, and the CalDAG-GEF or cAMP-GEF transformed cells and transgenic animals of the

invention, one of ordinary skill in the art is now enabled by perform a great variety of assays which will detect the modulation of CalDAG-GEF or cAMP-GEF activity by candidate compounds. Particularly preferred and contemplated embodiments are discussed in some detail below.

5 A. CalDAG-GEF and/or cAMP-GEF Expression

In one series of embodiments, specific measures of CalDAG-GEF or cAMP-GEF expression are employed to screen candidate compounds for their ability to affect CalDAG-GEF or cAMP-GEF activity. Thus, using the CalDAG-GEF or cAMP-GEF nucleic acids and antibodies disclosed and otherwise enabled herein, one may use mRNA levels or protein levels
10 as a marker for the ability of a candidate compound to modulate CalDAG-GEF or cAMP-GEF activity. The use of such probes and antibodies to measure gene and protein expression is well known in the art and discussed elsewhere herein.

B. Intracellular Localization

In another series of embodiments, compounds may be screened for their ability to
15 modulate the activity of the CalDAG-GEFs or cAMP-GEFs based upon their effects on the trafficking and intracellular localization of the CalDAG-GEFs or cAMP-GEFs. Differences in localization of mutant and normal CalDAG-GEFs and/or cAMP-GEFs may contribute to the etiology of CalDAG-GEF and/or cAMP-GEF-associated diseases. Compounds which can affect the localization of the CalDAG-GEFs and/or cAMP-GEFs may, therefore, be identified as
20 potential therapeutics. Standard techniques known in the art may be employed to detect the localization of the CalDAG-GEFs and/or cAMP-GEFs. Generally, these techniques will employ the antibodies of the present invention, and in particular antibodies which selectively bind to one or more mutant CalDAG-GEFs or cAMP-GEFs but not to normal CalDAG-GEFs or cAMP-GEFs. As is well known in the art, such antibodies may be labeled by any of a variety of
25 techniques (e.g., fluorescent or radioactive tags, labeled secondary antibodies, avidin-biotin, etc.) to aid in visualizing the intracellular location of the CalDAG-GEFs or cAMP-GEFs. The CalDAG-GEFs or cAMP-GEFs may be co-localized to particular structures, as is known in the art, using antibodies to markers of those structures (e.g., TGN38 for the Golgi, transferrin receptor for post-Golgi transport vesicles, LAMP2 for lysosomes). Western blots of purified
30 fractions from cell lysates enriched for different intracellular membrane bound organelles (e.g., lysosomes, synaptosomes, Golgi) may also be employed. In addition, the relative orientation of

different domains of the CalDAG-GEFs and/or cAMP-GEFs across cellular domains may be assayed using, for example, electron microscopy and antibodies raised to those domains.

9. Screening and Diagnostics for CalDAG-GEF- or cAMP-GEF-associated disorders

A. General Diagnostic Methods

5 The CalDAG-GEF or cAMP-GEF genes and gene products, as well as the CalDAG-GEF or cAMP-GEF-derived probes, primers and antibodies, disclosed or otherwise enabled herein, are useful in the screening for carriers of alleles associated with CalDAG-GEF- or cAMP-GEF-associated disorders. Individuals at risk for such a disorder or individuals not previously known to be at risk, may be routinely screened using probes to detect the presence of a mutant CalDAG-GEF or cAMP-GEF gene or protein by a variety of techniques. Diagnosis of inherited cases of these diseases can be accomplished by methods based upon the nucleic acids (including genomic and mRNA/cDNA sequences), proteins, and/or antibodies disclosed and enabled herein, including functional assays designed to detect increases or decreases of the normal CalDAG-GEF or cAMP-GEF activity and/or the presence of specific new activities conferred by the mutant CalDAG-GEFs or cAMP-GEFs. Preferably, the methods and products are based upon the human CalDAG-GEF or cAMP-GEF nucleic acids, proteins or antibodies, as disclosed or otherwise enabled herein. For brevity of exposition, but without limiting the scope of the invention, the following description will focus upon uses of the human homologues of CalDAG-GEF and cAMP-GEF. It will be understood, however, that homologous sequences from other species, including those disclosed herein, will be equivalent for many purposes.

B. Protein Based Screens and Diagnostics

When a diagnostic assay is to be based upon CalDAG-GEF or cAMP-GEF proteins, a variety of approaches are possible. For example, diagnosis can be achieved by monitoring differences in the electrophoretic mobility of normal and mutant proteins. Such an approach will be particularly useful in identifying mutants in which insertions, deletions or substitutions have resulted in a significant change in the electrophoretic migration of the resultant protein. Alternatively, diagnosis may be based upon differences in the proteolytic cleavage patterns of normal and mutant proteins, differences in molar ratios of the various amino acid residues, or by functional assays demonstrating altered function of the gene products.

30 C. Nucleic Acid Based Screens and Diagnostics

When the diagnostic assay is to be based upon nucleic acids from a sample, the assay may be based upon mRNA, cDNA or genomic DNA. Whether mRNA, cDNA, or genomic DNA is

- 40 -

assayed, standard methods well known in the art may be used to detect the presence of a particular sequence either *in situ* or *in vitro* (See, e.g., Sambrook et al., MOLECULAR CLONING: A LABORATORY MANUAL, 2nd ed. (1989)).

(1) Appropriate Probes and Primers

5 Whether for hybridization, RNase protection, ligase-mediated detection, PCR amplification or any other standards methods described herein and well known in the art, a variety of subsequences of the CalDAG-GEF and/or cAMP-GEF sequences disclosed or otherwise enabled herein will be useful as probes and/or primers. These sequences or subsequences will include both normal CalDAG-GEF or cAMP-GEF sequences and deleterious
10 mutant sequences. In general, useful sequences will include at least 8-10, more preferably 10-15, and most preferably 15-25 consecutive nucleotides from the CalDAG-GEF or cAMP-GEF introns, exons or intron/exon boundaries. In another embodiment, useful sequences include at least 25-500 consecutive nucleotides. Depending upon the target sequence, the specificity required, and future technological developments, shorter sequences may also have utility.
15 Therefore, any CalDAG-GEF or cAMP-GEF derived sequence which is employed to isolate, clone, amplify, identify or otherwise manipulate a CalDAG-GEF or cAMP-GEF sequence may be regarded as an appropriate probe or primer.

(2) Hybridization Screening

For *in situ* detection of a normal or mutant CalDAG-GEF, cAMP-GEF or other CalDAG-
20 GEF and/or cAMP-GEF-associated nucleic acid sequence, a sample of tissue may be prepared by standard techniques and then contacted with one or more of the above-described probes, preferably one which is labeled to facilitate detection, and an assay for nucleic acid hybridization is conducted under stringent conditions which permit hybridization only between the probe and highly or perfectly complementary sequences.

25 (3) Restriction Mapping

Sequence alterations may also create or destroy fortuitous restriction enzyme recognition sites which are revealed by the use of appropriate enzyme digestion followed by electrophoresis and visualization. DNA fragments carrying the site (normal or mutant) are detected by their increase or reduction in size, or by the increase or decrease of corresponding restriction fragment
30 numbers. Such restriction fragment length polymorphism analysis (RFLP), or restriction mapping, may be employed with genomic DNA, mRNA or cDNA. The CalDAG-GEF or cAMP-GEF sequences may be amplified by PCR using the above-described primers prior to

- 41 -

restriction, in which case the lengths of the PCR products may indicate the presence or absence of particular restriction sites, and/or may be subjected to restriction after amplification. The CalDAG-GEF or cAMP-GEF fragments may be visualized by any convenient means (*e.g.*, under UV light in the presence of ethidium bromide).

5 (4) PCR Mapping

In another series of embodiments, a single base substitution mutation may be detected based on differential PCR product length or production in PCR. Thus, primers which span mutant sites or which, preferably, have 3' termini at mutation sites, may be employed to amplify a sample of genomic DNA, mRNA or cDNA from a subject. A mismatch at a mutational site may
10 be expected to alter the ability of the normal or mutant primers to promote the polymerase reaction and, thereby, result in product profiles which differ between normal subjects and heterozygous and/or homozygous CalDAG-GEF or cAMP-GEF mutants.

(5) Electrophoretic Mobility

Genetic testing based on DNA sequence differences also may be achieved by detection of
15 alterations in electrophoretic mobility of DNA, mRNA or cDNA fragments in gels. Small sequence deletions and insertions, for example, can be visualized by high resolution gel electrophoresis of single or double stranded DNA, or as changes in the migration pattern of DNA heteroduplexes in non-denaturing gel electrophoresis.

(6) Chemical Cleavage of Mismatches

20 Mutations in the CalDAG-GEFs or cAMP-GEFs may also be detected by employing the chemical cleavage of mismatch (CCM) method (*See, e.g.*, Saleeba et al., METHODS IN ENZYMOLOGY, 217: 286-295 (1993)). In this technique, probes (up to ~ 1 kb) may be mixed with a sample of genomic DNA, cDNA or mRNA obtained from a subject. The sample and probes are mixed and subjected to conditions which allow for heteroduplex formation (if any).
25 Preferably, both the probe and sample nucleic acids are double-stranded, or the probe and sample may be PCR amplified together, to ensure creation of all possible mismatch heteroduplexes. Mismatched T residues are reactive to osmium tetroxide and mismatched C residues are reactive to hydroxylamine. Because each mismatched A will be accompanied by a mismatched T, and each mismatched G will be accompanied by a mismatched C, any nucleotide differences between
30 the probe and sample (including small insertions or deletions) will lead to the formation of at least one reactive heteroduplex. After treatment with osmium tetroxide and/or hydroxylamine to modify any mismatch sites, the mixture is subjected to chemical cleavage at any modified

- 42 -

mismatch sites by, for example, reaction with piperidine. The mixture may then be analyzed by standard techniques such as gel electrophoresis to detect cleavage products which would indicate mismatches between the probe and sample.

(7) Other Methods

5 Various other methods of detecting CalDAG-GEF or cAMP-GEF mutations, based upon the CalDAG-GEF or cAMP-GEF sequences disclosed and otherwise enabled herein, will be apparent to those of ordinary skill in the art. Any of these may be employed in accordance with the present invention. These include, but are not limited to, nuclease protection assays (S1 or ligase-mediated), ligated PCR, denaturing gradient gel electrophoresis (DGGE; *see, e.g.*, Fischer
10 et al., 80 PROC. NAT'L ACAD. SCI (USA), 1578-83 (1983)), restriction endonuclease fingerprinting combined with SSCP (REF-SSCP; *see, e.g.*, Liu et al., 18 BIOTECHNIQUES 470-79 (1995)), and the like.

D. Other Screens and Diagnostics

Diagnosis also can be made by observation of alterations in CalDAG-GEF or cAMP-GEF
15 transcription, translation, and post-translational modification and processing as well as alterations in the intracellular and extracellular trafficking of CalDAG-GEF or cAMP-GEF gene products in the brain and peripheral cells. Such changes will include alterations in the amount of CalDAG-GEF or cAMP-GEF messenger RNA and/or protein, alteration in phosphorylation state, abnormal intracellular location/distribution, abnormal extracellular distribution, etc. Such assays
20 will include: Northern Blots (with CalDAG-GEF or cAMP-GEF-specific and non-specific nucleotide probes), Western blots and enzyme-linked immunosorbent assays (ELISA) (with antibodies raised specifically to a CalDAG-GEF or a cAMP-GEF functional domain, including various post-translational modification states).

E. Screening and Diagnostic Kits

25 In accordance with the present invention, diagnostic kits are also provided which will include the reagents necessary for the above-described diagnostic screens. For example, kits may be provided which include antibodies or sets of antibodies which are specific to one or more mutant epitopes. These antibodies may, in particular, be labeled by any of the standard means which facilitate visualization of binding. Alternatively, kits may be provided in which
30 oligonucleotide probes or PCR primers, as described above, are present for the detection and/or amplification of mutant CalDAG-GEF, cAMP-GEF or other CalDAG-GEF and/or cAMP-GEF-associated nucleotide sequences. Again, such probes may be labeled for easier detection of

- 43 -

specific hybridization. As appropriate to the various diagnostic embodiments described above, the oligonucleotide probes or antibodies in such kits may be immobilized to substrates and appropriate controls may be provided.

10. Methods of Treatment

5 The present invention now provides a basis for therapeutic intervention in diseases which are associated to the CalDAG-GEFs or cAMP-GEFs in that they are caused, prevented, exacerbated, or alleviated, or which may be caused, prevented, exacerbated, or alleviated, by the either normal or mutant CalDAG-GEFs or cAMP-GEFs. In considering the various therapies described below, it is understood that such therapies may be targeted at tissue other than the brain
10 where CalDAG-GEF or cAMP-GEF are also expressed.

Therapies to treat CalDAG-GEF and/or cAMP-GEF-associated diseases may be based upon (1) administration of normal CalDAG-GEF or cAMP-GEF proteins, (2) gene therapy with normal CalDAG-GEF or cAMP-GEF genes to compensate for or replace the mutant genes, (3) gene therapy based upon antisense sequences to mutant CalDAG-GEF or cAMP-GEF genes or
15 which "knock-out" the mutant genes, (4) gene therapy based upon sequences which encode a protein which blocks or corrects the deleterious effects of CalDAG-GEF or cAMP-GEF mutants, (5) immunotherapy based upon antibodies to normal and/or mutant CalDAG-GEF or cAMP-GEF proteins, or (6) small molecules (drugs) which alter CalDAG-GEF or cAMP-GEF expression, block abnormal interactions between mutant forms of CalDAG-GEF or cAMP-GEF and other
20 proteins or ligands, or which otherwise block the aberrant function of mutant CalDAG-GEF or cAMP-GEF proteins by altering the structure of the mutant proteins, by enhancing their metabolic clearance, or by inhibiting their function.

A. Protein Therapy

Treatment of CalDAG-GEF and/or cAMP-GEF-associated disorders, or disorders
25 resulting from CalDAG-GEF and/or cAMP-GEF mutations, may be performed by providing an excess of inactive mutant protein to decrease the effect of the normal function of the protein, or by providing an excess of normal protein to reduce the effect of any aberrant function of the mutant protein, by replacing a mutant protein with normal protein, or by modulating the function of the mutant protein.

30 B. Gene Therapy

In one series of embodiments, gene therapy may be employed in which normal or mutant copies of the CalDAG-GEF gene or the cAMP-GEF gene are introduced into patients to code

- 44 -

successfully for normal or mutant protein in one or more different affected cell types. The gene must be delivered to those cells in a form in which it can be taken up and code for sufficient protein to provide effective function. Thus, it is preferred that the recombinant gene be operably joined to a strong promoter so as to provide a high level of expression which will compensate for, or out-compete, the naturally-occurring proteins. As noted above, the recombinant construct may contain endogenous or exogenous regulatory elements, inducible or repressible regulatory elements, or tissue-specific regulatory elements.

In another series of embodiments, gene therapy may be employed to replace the naturally-occurring gene by homologous recombination with a recombinant construct. The recombinant construct may contain a normal or a mutant copy of the targeted CalDAG-GEF and/or cAMP-GEF gene, in which case the defect is corrected *in situ*, or may contain a "knock-out" construct which introduces a stop codon, missense mutation, or deletion which abolished function of the mutant gene. It should be noted in this respect that such a construct may knock-out both the normal and mutant copies of the targeted CalDAG-GEF and/or cAMP-GEF gene in a heterozygous individual, but the total loss of CalDAG-GEF and/or cAMP-GEF gene function may be less deleterious to the individual than continued progression of the disease state.

In another series of embodiments, antisense gene therapy may be employed. The antisense therapy is based on the fact that sequence-specific suppression of gene expression can be achieved by intracellular hybridization between mRNA or DNA and a complementary antisense species. The formation of a hybrid duplex may then interfere with the transcription of the gene and/or the processing, transport, translation and/or stability of the target CalDAG-GEF and/or cAMP-GEF mRNA. Antisense strategies may use a variety of approaches including the administration of antisense oligonucleotides or antisense oligonucleotide analogs (*e.g.*, analogs with phosphorothioate backbones) or transfection with antisense RNA expression vectors. Again, such vectors may include exogenous or endogenous regulatory regions, inducible or repressible regulatory elements, or tissue-specific regulatory elements.

In another series of embodiments, gene therapy may be used to introduce a recombinant construct encoding a protein or peptide which blocks or otherwise corrects the aberrant function caused by a naturally-occurring CalDAG-GEF and/or cAMP-GEF gene. In one embodiment, the recombinant gene may encode a peptide which corresponds to a mutant domain of a CalDAG-GEF and/or cAMP-GEF which has been found to abnormally interact with another cell protein or other cell ligand. Alternatively, the portion of a protein which interacts with a mutant, but not a

- 45 -

normal, CalDAG-GEF and/or cAMP-GEF may be encoded and expressed by a recombinant construct in order to compete with, and thereby inhibit or block, the aberrant interaction.

Retroviral vectors can be used for somatic cell gene therapy especially because of their high efficiency of infection and stable integration and expression. The targeted cells however must be able to divide and the expression of the levels of normal protein should be high. The full length CalDAG-GEF or cAMP-GEF genes, subsequences encoding functional domains of the CalDAG-GEFs or cAMP-GEFs, or any of the other therapeutic peptides described above, can be cloned into a retroviral vector and driven from its endogenous promoter, from the retroviral long terminal repeat, or from a promoter specific for the target cell type of interest. Other viral vectors which can be used include adeno-associated virus, vaccinia virus, bovine papilloma virus, or a herpes virus such as Epstein-Barr virus.

C. Immunotherapy

Antibodies may be raised to a mutant CalDAG-GEF or cAMP-GEF protein (or a portion thereof) and be administered to a patient to bind or block the mutant protein and prevent its deleterious effects. Alternatively, antibodies may be raised to specific complexes between mutant or wild-type CalDAG-GEF or cAMP-GEF and their interaction partners.

A further approach is to stimulate endogenous antibody production to the desired antigen. An immunogenic composition may be prepared as injectables, as liquid solutions or emulsions. The CalDAG-GEF or cAMP-GEF protein or other antigen may be mixed with pharmaceutically acceptable excipients compatible with the protein. Such excipients may include water, saline, dextrose, glycerol, ethanol and combinations thereof. The immunogenic composition and vaccine may further contain auxiliary substances such as emulsifying agents or adjuvants to enhance effectiveness. Immunogenic compositions and vaccines may be administered parenterally by injection subcutaneously or intramuscularly.

The immunogenic preparations and vaccines are administered in such amount as will be therapeutically effective, protective and immunogenic. Dosage depends on the route of administration and will vary according to the size of the host.

D. Small Molecule Therapeutics

As described and enabled herein, the present invention provides for a number of methods of identifying small molecules or other compounds which may be useful in the treatment of CalDAG-GEF- or cAMP-GEF-associated disorders. Thus, for example, the present invention provides for methods of identifying CalDAG-GEF or cAMP-GEF binding proteins and, in

- 46 -

particular, methods for identifying proteins or other cell components which bind to or otherwise interact with mutant CalDAG-GEFs or cAMP-GEFs but not with the normal CalDAG-GEFs or cAMP-GEFs. The invention also provides for methods of identifying small molecules which can be used to disrupt undesired interactions between CalDAG-GEFs or cAMP-GEFs and other proteins or other cell components.

By identifying these proteins and analyzing these interactions, it is possible to screen for or design compounds which counteract or prevent the interaction, thereby, providing treatment for abnormal interactions. Therapies can be designed to modulate these interactions and thereby, to modulate CalDAG-GEF- or cAMP-GEF-associated disorders. The potential efficacy of these therapies can be tested by analyzing the affinity and function of these interactions after exposure to the therapeutic agent by standard pharmacokinetic measurements of affinity (*e.g.*, K_d , V_{max}) using synthetic peptides or recombinant proteins corresponding to functional domains of the CalDAG-GEF gene, the cAMP-GEF gene or other CalDAG-GEF and/or cAMP-GEF homologues. Another method for assaying the effect of any interactions involving functional domains is to monitor changes in the intracellular trafficking and post-translational modification of the relevant genes by *in situ* hybridization, immunohistochemistry, Western blotting and metabolic pulse-chase labeling studies in the presence of, and in the absence of, the therapeutic agents. A further method is to monitor the effects of "downstream" events including changes in second messenger events, *e.g.*, cAMP, intracellular Ca^{2+} , protein kinase activities, etc.

The effect of potential therapeutic agents in cell lines and whole animals can be monitored by monitoring transcription, translation, and post-translational modification of the CalDAG-GEF and/or cAMP-GEF proteins. Methods for these studies include Western and Northern blots, immunoprecipitation after metabolic labelling (pulse-chase) with radio-labelled methionine and ATP, and immunohistochemistry. The effect of these agents can also be monitored using studies which examine the relative binding affinities and relative amounts of CalDAG-GEF or cAMP-GEF proteins involved in interactions with Rap1A, using either standard binding affinity assays or co-precipitation and Western blots using antibodies to Rap1A, CalDAG-GEF, cAMP-GEF, or other CalDAG-GEF and/or cAMP-GEF homologues.

Therapy using antisense oligonucleotides to block the expression of the mutant CalDAG-GEF gene or the mutant cAMP-GEF gene, co-ordinated with gene replacement with normal CalDAG-GEF or cAMP-GEF gene can also be applied using standard techniques of either gene therapy or protein replacement therapy.

V. Examples

Example 1: Isolation and characterization of CalDAG-GEF.

Human full-length CalDAG-GEFI cDNAs were isolated from a human frontal cortex λ ZAPII cDNA library (Stratagene) and a U937 λ ZAPII cDNA library. Mouse full-length
5 CalDAG-GEFI was identified in the mouse EST database (GenBank accession number: W71787). Rat full-length CalDAG-GEFII cDNA was isolated from a rat whole brain λ ZAPII cDNA library by using human CalDAG-GEFII as a probe. Mouse ESTs identified through BLAST searches were purchased from Genome Systems Inc. (St. Louis, MO).

CalDAG-GEFI encodes an approximately 69-kD protein (Fig. 2D) that displays in its
10 amino terminal region a GEF domain that is highly homologous to Ras-superfamily GEFs (Fig. 2A-2D). Multiple alignment analysis shows that genes of the CalDAG-GEF family form a cluster within the Ras-GEF superfamily distinct from Ras GEFs such as Sos1 and rRas-GEF (Fig. 2B). The region downstream of the GEF domain contains two tandem repeats of EF-hand Ca^{2+} binding motifs (Figs. 2A, 2E). The carboxy-terminal region displays a typical
15 diacylglycerol/phorbol ester-binding domain, which is present in most PKC family proteins (Fig. 2A, 2F). Multiple sequence alignments and phylogenetic tree analysis were carried out with the LASERGENE Software Package (DNASTAR Inc.). Abbreviations and GenBank accession numbers of the protein sequences used in Figure 2 are as follows: C3G: 474982, mCdc25: 882120, rRas-GRF: 57665, hSos1 (human son-of-sevenless 1): 476780, BUD5: 171141,
20 hCalmodulin: 115512, hCalbindin D28k: 227666, hCalcineurin B: 105504, hParvalbumin a: 131100, hTroponin C: 136043, hPKCa: 125549, hPKCb1: 125538, hPKCg: 462455.

To determine the small G protein target of CalDAG-GEFI, guanine nucleotide exchange activity *in vivo* was analyzed using intact 293T cells cotransfected with a eukaryotic expression construct of mouse CalDAG-GEFI and GST-tagged Ras family proteins. Full-length mouse
25 CalDAG-GEFI cDNA inserted into a pCMV-SPORT expression vector with a carboxy-terminal FLAG epitope was used for transfection. A PCR-amplified fragment of rat CalDAG-GEFII was subcloned into a pCAGGS expression vector with the addition of His₆-tag at its amino-terminus, resulting in pCAGGS-His-CalDAGII. pEBG-Krev1 that expresses Rap1A was used as a fusion protein to glutathione S-transferase (GST) in mammalian cells, as described in Gotoh et al., 15
30 Mol. Cell Biol. 6746-53 (1995), pEBG-R-Ras, other vectors for Ras-family proteins obtained by inserting PCR-amplified cDNAs into pEBG expression vector, pCAGGS-C3G and pCAGGS-MSos1, and pCEV-H-RasV12. CalDAG-GEFI transfection produced a dramatic increase in

- 48 -

GTP-bound Rap1A compared to the control but showed no or minimal activation of H-Ras, R-Ras, or Ral A. The increase in GTP-bound Rap1A was augmented in the presence of either the Ca^{2+} ionophore, A23187, or the phorbol ester, phorbol-12-myristate-13-acetate (TPA). Further, A23187 and TPA had additive effects when administered together.

5 To determine the effect of CalDAG-GEFI on the Erk/MAP kinase cascade, Elk1 activation was measured in 293T cells transfected with CalDAG-GEFI or constitutively active H-Ras (RasV12), or both. 293T cells were transfected by SuperFect (Qiagen) as described in Gotoh, *supra*, with expression vectors for GST-tagged Ras family proteins and with those for various GEFs. Cells were labeled 24 hours after transfection with $^{32}\text{P}_i$ for 2 hr. In some
10 experiments, cells were stimulated with either 10 μM A23187 or 1 μM phorbol-12-myristate-13-acetate (TPA) for 3 min. GST-tagged Ras family proteins were collected from cell lysates with glutathione Sepharose. Guanine nucleotides bound to Ras family proteins were separated by thin layer chromatography (TLC). Activation of Elk1 was examined by the PathDetect Elk1 transreporting system (Stratagene). 293T cells were transfected with pFR-Luc and pFA-Elk1
15 with various expression vectors, and light output was detected and analyzed by the use of LAS1000 film. CalDAG-GEFI reduced RasV12 activation of Elk1 by approximately 4-fold and did not itself activate Elk1. Thus, CalDAG-GEFI strongly inhibits Ras-dependent stimulation of the Erk/MAP kinase cascade.

Northern analysis showed that human CalDAG-GEFI is expressed strongly in the brain
20 and that CalDAG-GEFI mRNA is strikingly enriched in the striatum. Probes used included human CalDAG-GEFI: 729-bp EcoRI fragment, human CalDAG-GEFII: 584-bp SacI and HindIII fragment, rat CalDAG-GEFI: 439-bp fragment of EST clone RBC565 (GenBank accession number: C06861, and rat CalDAG-GEFII: 508-bp PCR amplified and subcloned fragment (nucleic acids 2541 to 3048 of SEQ ID NO:5). *In situ* hybridization of sections from
25 the adult rat brain confirmed these restricted distribution patterns. Intense signal was present in the striatum (caudoputamen) and the ventral striatum (nucleus accumbens, olfactory tubercle). There was weaker signal in the olfactory bulb.

A series of monoclonal antibodies against the carboxy-terminal half of mouse CalDAG-GEFI were raised. His₆-tagged mouse CalDAG-GEFI polypeptide (amino acids 349 to 608 of
30 SEQ ID NO:1) was expressed in bacteria, purified over Ni^{2+} -nitrilotriacetic acid-agarose resin, and then used to immunize BALB/c mice. The resultant polyclonal antiserum was monitored by ELISA, Western blot, immunoprecipitation, and immunofluorescence assays on CalDAG-GEFI-

- 49 -

transfected COS-7 cells. Hybridomas were generated by PEG (polyethylene glycol)-mediated fusion of donor splenocytes to the SP2/O cell line. Positive hybridoma cell lines were identified by screening in the assays described above, and purified by limiting dilution and single-cell cloning. Three hybridoma cell lines against mouse CalDAG-GEFI (mAbs 18B11, 2D9, and 18A7), in addition to the polyclonal fusion serum, were identified. Western analysis showed that mAbs 18B11 and 2D9 were specific for CalDAG-GEFI. Lightly post-fixed, cryostat-cut 10 μ m thick sections were immunostained by the ABC (Vectastain kit) method for CalDAG-GEFI with mAbs 18B11 and 2D9 and the polyclonal fusion serum, for tyrosine hydroxylase (TH) with monoclonal antibodies from INCSTAR, and for μ opioid receptor with polyclonal antiserum. Immunohistochemistry with mAb 18B11 showed a striking basal ganglia-enriched distribution pattern in sections of adult rat brain, with significant but weaker activity elsewhere. CalDAG-GEFI immunoreactivity marked the entire pathway from the striatal matrix compartment to the pallidum and substantia nigra pars reticulata, where very intense CalDAG-GEFI staining was present. Thus, CalDAG-GEFI is synthesized in striatal projection neurons and is transported to striatopallidal and striatonigral terminals.

To confirm that CalDAG-GEFI is synthesized in striatal projection neurons and transported to striatopallidal and striatonigral terminals in rats, intrastriatal injections of ibotenic acid (20 μ g/ μ l, 1.5 μ l per site, 5 day survival) were made unilaterally at 2 sites in the mid-lateral caudoputamen, with contralateral vehicle control injections were made. In other rats, unilateral subthalamic knife-cuts were made at an anteroposterior level between the entopeduncular nucleus and substantia nigra to sever the striatonigral efferents (1 and 3 days survivals), with control contralateral thalamic knife-cuts. These procedures all reduced CalDAG-GEFI staining in the substantia nigra. *In situ* hybridization was performed according to Simmons et al, 12 J. Histotechnol. 169-181 (1989). A 439bp rat EST clone RBC565 (98.4% identical to mouse CalDAG-GEFI nucleic acids 1777 to 2216 of SEQ ID NO:1) was isolated by BLAST search and used for making RNA probes with ³²P-labeled UTP (2,000 Ci/mmol, NEN, 1 Ci = 37 GBq) and T3 and T7 RNA polymerase. Brains were processed as above for CalDAG-GEFI and TH immunostaining. Thus, CalDAG-GEFI is a protein transported in striatal axons to their terminals. The terminal localization of CalDAG-GEFI was confirmed in subcellular fractionation experiments on dissected samples from the rat ventral midbrain, in which Western analysis showed the presence of CalDAG-GEFI in cytosol and in membrane fractions, including synaptosomes.

- 50 -

Because of the similarity of the GEF domains of CalDAG-GEFI and CalDAG-GEFII, the substrate specificity of CalDAG-GEFII with the same 293T cell assay system used for CalDAG-GEFI was examined. It was confirmed that CalDAG-GEFII activates Ras, and further shown that it activates H-Ras and R-Ras, but not Ral A or Rap1A. H-Ras activation was enhanced by A23187 and TPA. Moreover, CalDAG-GEFII, unlike CalDAG-GEFI, increased the transcriptional activity of Elk1 downstream to Erk/MAP kinase. Thus, in the 293T system, CalDAG-GEFI and CalDAG-GEFII target different Ras-superfamily small G proteins and have opposite effects on the MAP kinase cascade. Northern analysis further showed contrasting brain expression for CalDAG-GEFII, with highest expression being in the cerebellum, cerebral cortex, and amygdala, and low expression occurring in the striatum. Both genes are also expressed in hematopoietic organs in both human and rat.

Rap signaling is important in regulating basal ganglia output in response to Ca^{2+} and DAG. Corticostriatal inputs can activate the MAP kinase cascade in striatal projection neurons (Sgambat et al., 18 J. Neurosci. 214-26 (1993)) and phosphoinositide (PI) signaling is strongly represented in these pathways (Fotuhi et al., 13 J. Neurosci. 3300-08 (1993)). Moreover, a number of receptor systems in the striatum and its striatonigral/striatopallidal pathways are linked to Ca^{2+} and PI signaling, notably including NMDA and metabotropic glutamate receptors, D_2 -class dopamine receptors, and tachykinin receptors (Fiorillo et al., 394 Nature 78-82 (1998)). A previously unrecognized signaling target for some of these systems is likely to be Rap1, via CalDAG-GEFI. In addition, CalDAG-GEFI has a synaptic function as demonstrated by the heavy accumulation of CalDAG-GEFI in the target nuclei of striatal outputs and the localization of Rap1 in synaptosomes and synaptic vesicles. The particular basal ganglia projection systems are enriched in CalDAG-GEFI and are differentially vulnerable to neurodegeneration in Huntington's disease.

Rap and Ras functions can be regulated coordinately or disjunctively by Ca^{2+} and DAG in the brain and hematopoietic organs, depending on the relative expression of CalDAG-GEFI and CalDAG-GEFII. In neurons, Ras/MAP kinase signaling has been directly implicated in synaptic transmission and the neuroplasticity underlying learning and memory. Different CalDAG-GEFI and CalDAG-GEFII expression patterns in the brain influence region-specific neuroplasticity mediated by Ca^{2+} and DAG signaling pathways. The presence of CalDAG-GEFI and CalDAG-GEFII in the hematopoietic system demonstrates the direct input of Ca^{2+} and DAG to Ras/Rap regulation of normal growth and differentiation as well as malignant transformation.

Example 2: Isolation and characterization of cAMP-GEFs.

cAMP-GEFI and cAMP-GEFII have similar domain structures, with a cAMP binding domain at the amino terminus and a GEF domain at the carboxy terminus separated by a link region (LR) (Fig. 3A). These mammalian proteins show strong structural homology to a predicted open reading frame (T20G5.5) in *C. elegans* cAMP-GEF (cel cAMP-GEF) (Fig. 3A). The cAMP binding domains of the cAMP-GEF family proteins form a distinct group within the cyclic nucleotide-binding protein superfamily and show the closest similarity to the B domains of PKA regulatory subunits (Fig. 3B). A PR(A/T)AT motif in the cAMP binding pocket is also conserved in the cAMP-GEF proteins (Fig. 3E). The first alanine of this motif confers cAMP (alanine) as opposed to cGMP (threonine) binding specificity. All of the cAMP-GEF family members have alanine at this position, and therefore bind cAMP rather than cGMP.

The GEF domains of the cAMP-GEFs show high homology to those of Ras-GEF family proteins, but form an independent cluster distinct from Ras GEFs such as mCdc25, hSos1, and rRas-GRF (Fig. 3, C and D). The three structurally conserved regions specific to Ras-GEF family proteins (SCR1, SCR2, and SCR3) are present in all of the cAMP-GEF proteins (Fig. 3D). Multiple sequence alignments and phylogenetic tree analyses were carried out with LASERGENE (DNASTAR Inc.). Abbreviations and GenBank accession numbers of the protein sequences used in this figure: hPKARI α (human cAMP-dependent protein kinase regulatory subunit type I-alpha): 125193, hPKARI β : 1346362, hPKARII α : 125198, hPKARII β : 400115, hPKGII α (human cGMP-dependent protein kinase type I-alpha): 1255602, hPKGII β : 125379, hPKGII: 1906312, hCalDAG-GEFI: U71870, hCalDAG-GEFII: AF081195, C3G: 474982, hSos1 (human son-of-sevenless 1): 476780, mCdc25: 882120, rRas-GRF: 57665, BUD5: 171141.

In order to identify the small G protein substrate for cAMP-GEFI and II and the mode of cAMP regulation of GEF activity conferred by these proteins, the effects of cAMP-GEFI and cAMP-GEFII expression were analyzed in 293T cells on the ratio of GTP to GDP bound to different Ras family small G proteins in the presence or absence of forskolin and IBMX. Under basal conditions, in the absence of forskolin and IBMX, only Rap1 was activated significantly. In the presence of forskolin and IBMX, both cAMP-GEFI and II strongly and selectively activated Rap1A, but did not activate H-Ras, R-Ras or RalA. The effects of forskolin/IBMX treatment on cAMP-GEFI and II were dose-dependent with EC₅₀ values of 1.8 μ M and 0.3 μ M, respectively. Forskolin/IBMX treatment given alone had no effect.

- 52 -

A time-course analysis of the activation of Rap1A by forskolin/IBMX in cAMP-GEFI transfectants showed that the activation began within 10 sec, reached a maximum at 5 min, and continued for at least 60 min. Thus, cAMP-GEFI has a direct effect on Rap1A rather than secondary effects mediated by other Ras-superfamily GEFs. In addition, Sp-cAMPS, an analogue of cAMP, activated Rap1A at levels similar to those induced by forskolin/IBMX. Thus, cAMP has the capacity to activate the GEF domain of cAMP-GEFI.

Mutational analyses with cAMP-GEFI was performed to examine whether its cAMP-binding domain is required for the activation of Rap1A. In contrast to wild type cAMP-GEFI, a deletion mutant lacking a cAMP binding domain (pcDNA-rcAMP-GEFI:DcAMP(528) and (595)) did not activate Rap1A with or without forskolin/IBMX treatment. Mutants with a single amino acid substitution at the cAMP binding pocket (pcDNA-rcAMP-GEFI:R(279)K) responded minimally to forskolin/IBMX treatment. Thus, the cAMP binding domain of cAMP-GEFI is necessary for its cAMP-dependent activation of Rap1A.

To assess further the cAMP binding capacity of cAMP-GEFI, a cAMP agarose affinity bead binding assay was performed. *In vitro* translated, radiolabeled cAMP-GEFI showed selective binding to the beads that was competed by excess amounts of either cAMP or 8-Br-cAMP. cAMP-GEF protein can bind cAMP and that this binding can activate Rap1A.

cAMP-dependent activation of Rap1 has previously been ascribed to the phosphorylation of Rap1A by PKA, which raises its affinity to smgGDS, a GEF with broad substrate specificity. However, at least in the 293T cell assay system, an increase of GTP-bound Rap1A in response to increasing cAMP levels with forskolin or treatment with the cAMP analogue, Sp-cAMPS was not detected in the absence of cAMP-GEFs. In addition, even in the presence of H-89, a potent and selective inhibitor of PKA, cAMP-GEFI and II could still activate Rap1A. The activation of Rap1A induced by cAMP-GEFI and II is independent of the PKA pathway.

Intracellular cAMP has been shown to interact directly with ion channels, but the vast majority of cAMP-mediated effects in eukaryotes have been considered as sequels to cAMP binding by the regulatory subunits of the PKA tetramer. The diversity of physiological effects produced by cAMP have been attributed to the fact that, as a kinase, PKA has a large range of molecular targets. Reported herein are novel cAMP binding proteins that directly link the cAMP second messenger system to Ras superfamily signaling pathways and that appear selectively to target Rap.

cAMP can exert profound cell-type specific effects on cell growth and differentiation and that cAMP is capable of inhibiting or stimulating the Ras/mitogen-activated protein (MAP) kinase/Erk pathway. The inhibition can occur at the initial translocation step by which Ras activates Raf, whereas activation of Rap1 is thought to occur through phosphorylation by PKA.

5 Activation of Rap1 has been suggested to be part of a switch mechanism determining whether growth or differentiation occurs in response to nerve growth factor (NGF). cAMP-GEFs directly couple cAMP to Rap1, itself discovered as a negative regulator of Ras but suspected of having independent functions as well. Thus, different levels of cAMP-GEF expression confer cell-type specific regulation of Ras superfamily signaling systems.

10 The genes also exhibit developmentally regulated expression in the septum, medial thalamus and habenula, key structures in the limbic system variously linked to brain reward circuits, addiction and schizophrenia. Thus, cAMP-GEFs, in addition to PKA, underlie some of the neuronal functions of cAMP.

15 Example 3. Northern hybridization demonstrating the expression of CalDAG-GEFI and CalDAG-GEFII protein mRNAs in a variety of tissues.

Total cytoplasmic RNA was isolated from various human tissue samples including amygdala, cerebellum, corpus callosum, caudate nucleus, cortex, frontal lobe, hippocampus, kidney, liver, lung, medulla oblongata, occipital pole, putamen, spinal cord, substantia nigra, subthalamic nucleus, thalamus, and temporal lobe, obtained from surgical pathology using
20 standard procedures such as CsCl purification. The RNA was then electrophoresed on a formaldehyde gel to permit size fractionation. The nitrocellulose membrane was prepared and the RNA was then transferred onto the membrane. ³²P-labeled cDNA probes were prepared and added to the membrane in order for hybridization between the probe the RNA to occur. After washing, the membrane was wrapped in plastic film and placed into imaging cassettes containing
25 X-ray film. The autoradiographs were then allowed to develop for one to several days. Sizing was established by comparison to standard RNA markers. These northern blots demonstrated that the CalDAG-GEF genes are strongly expressed in the brain. Weaker hybridization was detectable elsewhere.

30 Example 4. Northern hybridization demonstrating the expression of cAMP-GEFI and cAMP-GEFII protein mRNAs in a variety of tissues.

- 54 -

Northern hybridization analysis was performed as in Example 3 to detect the expression of the cAMP-GEFI and cAMP-GEFII genes in a variety of human tissues. The tissues analyzed included adrenal gland, amygdala, bone marrow, cerebellum, corpus callosum, caudate nucleus, colon (mucosal lining), caudputamen, cortex, frontal lobe, hippocampus, habenula, heart, kidney, liver, lung, lymph node, medulla obongata, occipital pole, olfactory bulb, ovary, pons, pancreas, putamen, septum, small intestines, skeletal muscle, spinal cord, spleen, stomach, substantia nigra, subthalamic nucleus, testis, thalamus, temporal lobe, thymus, trachea, and thyroid.

A striking contrast in the expression patterns of human cAMP-GEFI and II was observed by Northern analysis. Human cAMP-GEFI is widely expressed, with highest levels appearing in kidney, spleen, thyroid, heart, and pancreas. Human cAMP-GEFII shows a remarkably selective enrichment in the brain and the adrenal glands. Both genes were developmentally regulated. The expression patterns of the two genes in the nervous system also differ, with cAMP-GEFI having a wider expression than cAMP-GEFII. These region-specific neuronal expression patterns were confirmed in *in situ* hybridization experiments. cAMP-GEFI mRNA is expressed broadly at low levels in the adult brain, but it is strongly and selectively expressed in parts of the neonatal brain, including the septum and the thalamus. By contrast, cAMP-GEFII is strongly expressed in the mature as well as the developing brain. Notable are the high levels of cAMP-GEFII mRNA in the cerebral cortex, the hippocampus (especially CA3 and dentate gyrus), the habenula and the cerebellum. Genes of the cAMP-GEF family have widespread influence on cAMP functions in bodily organs and also region-specific functions in the brain.

Example 5. Isolation of CalDAG-GEF or cAMP-GEF binding proteins by yeast two-hybrid system.

To identify proteins interacting with the CalDAG-GEF or cAMP-GEF proteins, a yeast expression plasmid vector (pAS2-1, Clontech) is generated by ligating an in-frame partial cDNA sequence encoding either residues of the CalDAG-GEF protein or residues of the cAMP-GEF protein into the EcoRI and BamHI sites of the vector. The resultant fusion protein contains the GAL4 DNA binding domain coupled in-frame either to residues of the CalDAG-GEF protein or to residues of the cAMP-GEF protein. These expression plasmids are co-transformed, along with purified plasmid DNA from the human brain cDNA:pACT library, into yeast using the protocols of the Clontech Matchmaker yeast-two-hybrid kit (Clontech). Yeast clones bearing human brain cDNAs which interact with the CalDAG-GEF or cAMP-GEF fragments are selected by HIS resistance and β gal⁺ activation. The clones are further selected by

- 55 -

cyclohexamide sensitivity and the inserts of the human brain cDNAs are isolated by PCR and sequenced.

Although preferred embodiments of the invention have been described herein in detail, it will be understood by those skilled in the art that variations may be made thereto without
5 departing from the spirit of the invention or the scope of the following claims.

- 56 -

CLAIMS

What is claimed is:

1. An isolated nucleic acid comprising a nucleotide sequence encoding a protein selected from the group consisting of a normal CalDAG-GEFI protein, a mutant CalDAG-GEFI protein, a normal CalDAG-GEFII protein, and a mutant CalDAG-GEFII protein.
2. An isolated nucleic acid comprising a nucleotide sequence encoding a protein selected from the group consisting of a normal cAMP-GEFI protein, a mutant cAMP-GEFI protein, a normal cAMP-GEFII protein, and a mutant cAMP-GEFII protein.
3. An isolated nucleic acid as in claim 1 wherein said nucleic acid encodes a normal CalDAG-GEF protein and wherein said nucleotide sequence is selected from the group consisting of
 - (a) a sequence encoding a protein comprising the human CalDAG-GEFI amino acid sequence of SEQ ID NO: 4;
 - (b) a sequence encoding a protein comprising the murine CalDAG-GEFI amino acid sequence of SEQ ID NO: 2;
 - (c) a sequence encoding a protein comprising the human CalDAG-GEFII amino acid sequence of SEQ ID NO: 8; and
 - (d) a sequence encoding a protein comprising the murine CalDAG-GEFII amino acid sequence of SEQ ID NO: 6; and
 - (e) a sequence encoding a normal CalDAG-GEF protein and capable of hybridizing to a sequence complementary to any sequence of (a) - (d) under stringent hybridization conditions.
4. An isolated nucleic acid as in claim 2 wherein said nucleic acid encodes a normal cAMP-GEF protein and wherein said nucleotide sequence is selected from the group consisting of
 - (a) a sequence encoding a protein comprising the human cAMP-GEFI amino acid sequence of SEQ ID NO: 12;
 - (b) a sequence encoding a protein comprising the alternatively spliced human cAMP-GEFI amino acid sequence of SEQ ID NO: 14;
 - (c) a sequence encoding a protein comprising the rat cAMP-GEFI amino acid sequence of SEQ ID NO: 10;
 - (d) a sequence encoding a protein comprising the human cAMP-GEFII amino acid sequence of SEQ ID NO: 18;

- 57 -

(e) a sequence encoding a protein comprising the rat cAMP-GEFII amino acid sequence of SEQ ID NO: 16; and

(f) a sequence encoding a normal cAMP-GEF protein and capable of hybridizing to a sequence complementary to any sequence of (a) - (e) under stringent hybridization conditions.

5. An isolated nucleic acid comprising a nucleotide sequence of at least 8 consecutive nucleotides selected from the group consisting SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, SEQ ID NO: 17, and a sequence complementary to any of these sequences.

6. An isolated nucleic acid comprising a nucleotide sequence of at least 10 consecutive nucleotides selected from the group consisting SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, SEQ ID NO: 17, and a sequence complementary to any of these sequences.

7. An isolated nucleic acid comprising a nucleotide sequence of at least 15 consecutive nucleotides selected from the group consisting SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, SEQ ID NO: 17, and a sequence complementary to any of these sequences.

8. An isolated nucleic acid comprising a nucleotide sequence encoding at least one functional domain of a CalDAG-GEF protein selected from the group consisting of a normal CalDAG-GEFI protein, a mutant CalDAG-GEFI protein, a normal CalDAG-GEFII protein, and a mutant CalDAG-GEFII protein.

9. An isolated nucleic acid comprising a nucleotide sequence encoding at least one functional domain of a cAMP-GEF protein selected from the group consisting of a normal cAMP-GEFI protein, a normal cAMP-GEFII protein, a mutant cAMP-GEFI protein, and a mutant cAMP-GEFII protein.

10. An isolated nucleic acid comprising a nucleotide sequence encoding an antigenic determinant of a CalDAG-GEF protein selected from the group consisting of a normal CalDAG-GEFI protein, a normal CalDAG-GEFII protein, a mutant CalDAG-GEFI protein, and a mutant CalDAG-GEFII protein.

11. An isolated nucleic acid comprising a nucleotide sequence encoding an antigenic determinant of a cAMP-GEF protein selected from the group consisting of a normal cAMP-GEFI protein, a normal cAMP-GEFII protein, a mutant cAMP-GEFI protein, and a mutant cAMP-GEFII protein.

- 58 -

1 12. A method for identifying an allelic variant or heterospecific homologue of a human
2 CalDAG-GEF gene comprising:

3 choosing a nucleic acid probe or primer capable of hybridizing to a human CalDAG-
4 GEF gene sequence under stringent hybridization conditions;

5 mixing said probe or primer with a sample of nucleic acids which may contain a
6 nucleic acid corresponding to said variant or homologue; and

7 detecting hybridization of said probe or primer to said nucleic acid corresponding to
8 said variant or homologue.

1 13. A method as in claim 12 wherein said human CalDAG-GEF gene sequence is
2 selected from the group consisting of SEQ ID NO: 3 and SEQ ID NO: 7.

1 14. A method as in claim 12 wherein said sample comprises a sample of nucleic acids
2 selected from the group consisting of human genomic DNA, human mRNA, and human cDNA.

1 15. A method as in claim 12 wherein said sample comprises a sample of nucleic acids
2 selected from the group consisting of mammalian genomic DNA, mammalian mRNA, and
3 mammalian cDNA.

1 16. A method as in claim 12 wherein said sample comprises a sample of nucleic acids
2 selected from the group consisting of invertebrate genomic DNA, invertebrate mRNA, and
3 invertebrate cDNA.

1 17. A method as in claim 12 further comprising the step of isolating said nucleic acid
2 corresponding to said variant or homologue.

1 18. A method as in claim 12 wherein said nucleic acid is identified by hybridization.

1 19. A method as in claim 12 wherein said nucleic acid is identified by PCR amplification.

1 20. A method for identifying allelic variants or heterospecific homologues of a human
2 cAMP-GEF gene comprising:

3 choosing a nucleic acid probe or primer capable of hybridizing to a human cAMP-
4 GEF gene sequence under stringent hybridization conditions;

5 mixing said probe or primer with a sample of nucleic acids which may contain a
6 nucleic acid corresponding to said variant or homologue; and

7 detecting hybridization of said probe or primer to said nucleic acid corresponding to
8 said variant or homologue.

1 21. A method as in claim 12 wherein said human cAMP-GEF gene sequence is selected
2 from the group consisting of SEQ ID NO: 11, SEQ ID NO: 13, and SEQ ID NO: 17.

- 59 -

- 1 22. A method as in claim 20 wherein said sample comprises a sample of nucleic acids
2 selected from the group consisting of human genomic DNA, human mRNA, and human cDNA.
- 1 23. A method as in claim 20 wherein said sample comprises a sample of nucleic acids
2 selected from the group consisting of mammalian genomic DNA, mammalian mRNA, and
3 mammalian cDNA.
- 1 24. A method as in claim 20 wherein said sample comprises a sample of nucleic acids
2 selected from the group consisting of invertebrate genomic DNA, invertebrate mRNA, and
3 invertebrate cDNA.
- 1 25. A method as in claim 20 further comprising the step of isolating said nucleic acid
2 corresponding to said variant or homologue.
- 1 26. A method as in claim 20 wherein said nucleic acid is identified by hybridization.
- 1 27. A method as in claim 20 wherein said nucleic acid is identified by PCR amplification.
- 1 28. A method for identifying an allelic variant or heterospecific homologue of a human
2 CalDAG-GEF gene comprising:
3 choosing an antibody capable of selectively binding to a human CalDAG-GEF
4 protein;
5 mixing said antibody with a sample of proteins which may contain a protein
6 corresponding to said variant or homologue; and
7 detecting binding of said antibody to said protein corresponding to said variant or
8 homologue.
- 1 29. A method as in claim 28 wherein said sample comprises a sample of proteins selected
2 from the group consisting of human proteins, human fusion proteins, and proteolytic fragments
3 thereof.
- 1 30. A method as in claim 28 wherein said sample comprises a sample of nucleic acids
2 selected from the group consisting of mammalian proteins, mammalian fusion proteins, and
3 proteolytic fragments thereof.
- 1 31. A method as in claim 28 wherein said sample comprises a sample of nucleic acids
2 selected from the group consisting of invertebrate proteins, invertebrate fusion proteins, and
3 proteolytic fragments thereof.
- 1 32. A method as in claim 28 further comprising the step of substantially purifying said
2 protein corresponding to said variant or homologue.

- 60 -

- 1 33. A method for identifying an allelic variant or heterospecific homologue of a human
2 cAMP-GEF gene comprising:
3 choosing an antibody capable of selectively binding to a human cAMP-GEF protein;
4 mixing said antibody with a sample of proteins which may contain a protein
5 corresponding to said variant or homologue; and
6 detecting binding of said antibody to said protein corresponding to said variant or
7 homologue.
- 1 34. A method as in claim 33 wherein said sample comprises a sample of proteins selected
2 from the group consisting of human proteins, human fusion proteins, and proteolytic fragments
3 thereof.
- 1 35. A method as in claim 33 wherein said sample comprises a sample of proteins selected
2 from the group consisting of mammalian proteins, mammalian fusion proteins, and proteolytic
3 fragments thereof.
- 1 36. A method as in claim 33 wherein said sample comprises a sample of proteins selected
2 from the group consisting of invertebrate proteins, invertebrate fusion proteins, and proteolytic
3 fragments thereof.
- 1 37. A method as in claim 33 further comprising the step of substantially purifying said
2 protein corresponding to said variant or homologue.
- 1 38. An isolated nucleic acid comprising an allelic variant or a heterospecific homologue
2 of a gene selected from the group consisting of a human CalDAG-GEF gene, and a human
3 cAMP-GEF gene.
- 1 39. An isolated nucleic acid encoding an allelic variant or heterospecific homologue of a
2 protein selected from the group consisting of a human CalDAG-GEF protein, and a human
3 cAMP-GEF protein.
- 1 40. An isolated nucleic acid comprising a recombinant vector including a nucleotide
2 sequence selected from the group consisting of SEQ ID NO: 1, SEQ ID NO: 3, SEQ ID NO: 5,
3 SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, SEQ ID NO:
4 17, and a sequence complementary to any of these sequences.
- 1 41. An isolated nucleic acid as in claim 40 wherein said vector is an expression vector
2 and said nucleotide sequence is operably joined to a regulatory region.
- 1 42. An isolated nucleic acid as in claim 41 wherein said expression vector may express
2 said nucleotide sequence in mammalian cells.

- 61 -

- 1 43. An isolated nucleic acid as in claim 42 wherein said cells are selected from the group
2 consisting of fibroblast, liver, kidney, spleen, bone marrow, and neurological cells.
- 1 44. An isolated nucleic acid as in claim 42 wherein said vector is selected from the group
2 consisting of vaccinia virus, adenovirus, retrovirus, neurotropic viruses, and Herpes simplex.
- 1 45. An isolated nucleic acid as in claim 41 wherein said expression vector encodes at
2 least a functional domain of a protein selected from the group consisting of normal CalDAG-
3 GEFI, a normal CalDAG-GEFII, a mutant CalDAG-GEFI, a mutant CalDAG-GEFII, a normal
4 cAMP-GEFI, a normal cAMP-GEFII, a mutant cAMP-GEFI, and a mutant cAMP-GEFII.
- 1 46. An isolated nucleic acid as in claim 41 wherein said vector further comprises
2 sequences encoding an exogenous protein operably joined to said nucleotide sequence and
3 whereby said vector encodes a fusion protein.
- 1 47. An isolated nucleic acid as in claim 46 wherein said exogenous protein is selected
2 from the group consisting of lacZ, trpE, maltose-binding protein, poly-His tags, and glutathione-
3 S-transferase.
- 1 48. An isolated nucleic acid comprising a recombinant expression vector including
2 nucleotide sequences corresponding to an endogenous regulatory region of a gene selected from
3 the group consisting of a CalDAG-GEF gene, and a cAMP-GEF gene.
- 1 49. An isolated nucleic acid as in claim 48 wherein said endogenous regulatory region is
2 operably joined to a marker gene.
- 1 50. A host cell transformed with an expression vector of any one of claims 41-49, or a
2 descendant thereof.
- 1 51. A host cell as in claim 50 wherein said host cell is selected from the group consisting
2 of bacterial cells and yeast cells.
- 1 52. A host cell as in claim 50 wherein said host cell is selected from the group consisting
2 of fetal cells, embryonic stem cells, zygotes, gametes, and germ line cells.
- 1 53. A host cell as in claim 50 wherein said cell is selected from the group consisting of
2 fibroblast, liver, kidney, spleen, bone marrow and neurological cells.
- 1 54. A host cell as in claim 50 wherein said cell is an invertebrate cell.
- 1 55. A non-human animal model for cancer, wherein a genome of said animal, or an
2 ancestor thereof, has been modified by at least one recombinant construct, and wherein said
3 recombinant construct has introduced a modification selected from the group consisting of

- 62 -

(a) insertion of nucleotide sequences encoding at least a functional domain of a heterospecific normal CalDAG-GEF gene;

(b) insertion of nucleotide sequences encoding at least a functional domain of a heterospecific mutant CalDAG-GEF gene;

(c) insertion of nucleotide sequences encoding at least a functional domain of a conspecific homologue of a heterospecific mutant CalDAG-GEF gene;

(d) inactivation of an endogenous CalDAG-GEF gene;

(e) insertion of nucleotide sequences encoding at least a functional domain of a heterospecific normal cAMP-GEF gene;

(f) insertion of nucleotide sequences encoding at least a functional domain of a heterospecific mutant cAMP-GEF gene;

(g) insertion of nucleotide sequences encoding at least a functional domain of a conspecific homologue of a heterospecific mutant cAMP-GEF gene; and

(h) inactivation of an endogenous cAMP-GEF gene.

56. A non-human animal model as in claim 55 wherein said cancer is related to the Ras-pathway.

57. A non human animal model as in claim 56 wherein said cancer is selected from the group consisting of lung cancer, pancreatic cancer, breast cancer, colorectal cancer, and myeloid leukemia.

58. An animal model as in claim 55 wherein said modification is an insertion of a nucleotide sequence encoding at least a functional domain of a protein selected from the group consisting of a normal human CalDAG-GEF, and a normal cAMP-GEF gene.

59. An animal model as in claim 55 wherein said modification is an insertion of a nucleotide sequence encoding at least a functional domain of a protein selected from the group consisting of a mutant human CalDAG-GEF, and a mutant human cAMP-GEF gene.

60. An animal as in claim 55 wherein said animal is selected from the group consisting of rats, mice, hamsters, guinea pigs, rabbits, dogs, cats, goats, sheep, pigs, and non-human primates.

61. An animal as in claim 55 wherein said animal is an invertebrate.

62. A method for producing at least a functional domain of a protein selected from the group consisting of a CalDAG-GEF protein, and a cAMP-GEF protein, said method comprising culturing a host cell of any of claims 50-54 under suitable conditions to produce said protein by expressing said nucleic acid.

- 63 -

63. A substantially pure preparation of a protein selected from the group consisting of a normal CalDAG-GEF protein, a mutant CalDAG-GEF protein, a normal cAMP-GEF protein, and a mutant cAMP-GEF protein.

64. A substantially pure preparation as in claim 63 wherein said protein comprises a normal protein selected from the group consisting of

(a) a protein comprising the amino acid sequence of SEQ ID NO: 2;

(b) a protein comprising the amino acid sequence of SEQ ID NO: 4;

(c) a protein comprising the amino acid sequence of SEQ ID NO: 6;

(d) a protein comprising the amino acid sequence of SEQ ID NO: 8;

(e) a protein comprising the amino acid sequence of SEQ ID NO: 10;

(f) a protein comprising the amino acid sequence of SEQ ID NO: 12;

(g) a protein comprising the amino acid sequence of SEQ ID NO: 14;

(h) a protein comprising the amino acid sequence of SEQ ID NO: 16; and

(i) a protein comprising the amino acid sequence of SEQ ID NO: 18.

65. A substantially pure preparation of a polypeptide comprising an amino acid sequence of at least 4 consecutive amino acid residues selected from the group consisting of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, and SEQ ID NO: 18.

66. A substantially pure preparation of a polypeptide comprising an amino acid sequence of at least 10 consecutive amino acid residues selected from the group consisting of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, and SEQ ID NO: 18.

67. A substantially pure preparation of a polypeptide comprising an amino acid sequence of at least 15 consecutive amino acid residues selected from the group consisting of SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, and SEQ ID NO: 18.

68. A substantially pure preparation of a polypeptide comprising at least one functional domain of a protein selected from the group consisting of a normal CalDAG-GEF protein, a mutant CalDAG-GEF protein, a normal cAMP-GEF protein, and a mutant cAMP-GEF protein.

69. A substantially pure preparation of a polypeptide comprising an antigenic determinant of a protein selected from the group consisting of a normal CalDAG-GEF protein, a mutant CalDAG-GEF protein, a normal cAMP-GEF protein, and a mutant cAMP-GEF protein.

- 64 -

1 70. A method of producing antibodies which selectively bind to a CalDAG-GEF protein
2 comprising the steps of

3 administering an immunogenically effective amount of a CalDAG-GEF immunogen
4 to an animal;

5 allowing said animal to produce antibodies to said immunogen; and

6 obtaining said antibodies from said animal or from a cell culture derived therefrom.

1 71. A method of producing antibodies which selectively bind to a cAMP-GEF protein
2 comprising the steps of

3 administering an immunogenically effective amount of a cAMP-GEF immunogen to
4 an animal;

5 allowing said animal to produce antibodies to said immunogen; and

6 obtaining said antibodies from said animal or from a cell culture derived therefrom.

1 72. A substantially pure preparation of an antibody which selectively binds to an
2 antigenic determinant of a protein selected from the group consisting of a normal CalDAG-GEF
3 protein, a mutant CalDAG-GEF protein, a normal cAMP-GEF protein, and a mutant cAMP-GEF
4 protein.

1 73. A substantially pure preparation of an antibody as in claim 72 wherein said antibody
2 selectively binds to an antigenic determinant of a mutant CalDAG-GEF and fails to bind to a
3 normal CalDAG-GEF protein.

1 74. A substantially pure preparation of an antibody as in claim 72 wherein said antibody
2 selectively binds to an antigenic determinant of a mutant cAMP-GEF and fails to bind to a
3 normal cAMP-GEF protein.

1 75. A cell line producing an antibody of any one of claims 72-74.

1 76. A method for identifying compounds which can modulate the expression of a
2 CalDAG-GEF gene comprising:

3 contacting a cell with a test candidate wherein said cell includes a regulatory region of
4 a CalDAG-GEF gene operably joined to a coding region; and

5 detecting a change in expression of said coding region.

1 77. A method for identifying compounds which can modulate the expression of a cAMP-
2 GEF gene comprising:

3 contacting a cell with a test candidate wherein said cell includes a regulatory region of
4 a cAMP-GEF gene operably joined to a coding region; and

- 65 -

- 5 detecting a change in expression of said coding region.
- 1 78. A method as in claim 76 or 77 wherein said change comprises a change in a level of
2 an mRNA transcript encoded by said coding region.
- 1 79. A method as in claim 78 wherein said change comprises a change in a level of a
2 protein encoded by said coding region.
- 1 80. A method as in claim 78 wherein said change is a result of an activity of a protein
2 encoded by said coding region.
- 1 81. A method as in claim 78 wherein said coding region encodes a marker protein
2 selected from the group consisting of β -galactosidase, alkaline phosphatase, green fluorescent
3 protein, and luciferase.
- 1 82. A method for identifying compounds which can selectively bind to a CalDAG-GEF
2 protein comprising the steps of
3 providing a preparation including at least one CalDAG-GEF component;
4 contacting said preparation with a sample including at least one candidate compound;
5 and
6 detecting binding of said CalDAG-GEF component to said candidate compound.
- 1 83. A method for identifying compounds which can selectively bind to a cAMP-GEF
2 protein comprising the steps of
3 providing a preparation including at least one cAMP-GEF component;
4 contacting said preparation with a sample including at least one candidate compound;
5 and
6 detecting binding of said cAMP-GEF component to said candidate compound.
- 1 84. The method in claim 82 wherein said binding to said CalDAG-GEF component is
2 detected by an assay selected from the group consisting of: affinity chromatography, co-
3 immunoprecipitation, a Biomolecular Interaction Assay, and a yeast two-hybrid system.
- 1 85. The method in claim 83 wherein said binding to said cAMP-GEF component is
2 detected by an assay selected from the group consisting of: affinity chromatography, co-
3 immunoprecipitation, a Biomolecular Interaction Assay, and a yeast two-hybrid system.
- 1 86. A method of identifying compounds which can modulate activity of a CalDAG-GEF
2 comprising the steps of
3 providing a cell expressing a normal or mutant CalDAG-GEF gene;
4 contacting said cell with at least one candidate compound; and

- 66 -

5 detecting a change in a marker of said activity.

1 87. A method of identifying compounds which can modulate activity of a cAMP-GEF
2 comprising the steps of

3 providing a cell expressing a normal or mutant cAMP-GEF gene;

4 contacting said cell with at least one candidate compound; and

5 detecting a change in a marker of said activity.

1 88. A method as in claim 86 wherein measurement of said marker indicates a difference
2 between cells bearing an expressed mutant CalDAG-GEF gene and otherwise identical cells free
3 of an expressed mutant CalDAG-GEF gene.

1 89. A method as in claim 86 wherein said change comprises a change in a non-specific
2 marker of cell physiology selected from the group consisting of pH, intracellular calcium, cyclic
3 AMP levels, GTP/GDP ratios, phosphatidylinositol activity, and protein phosphorylation.

1 90. A method as in claim 86 wherein said change comprises a change in expression of
2 said CalDAG-GEF.

1 91. A method as in claim 86 wherein said change comprises a change in occurrence or
2 rate of apoptosis or cell death.

1 92. A method as in claim 86 wherein said cell is a cell cultured *in vitro*.

1 93. A method as in claim 92 wherein said cell is a transformed host cell of any one of
2 claims 50-54.

1 94. A method as in claim 92 wherein said cell is explanted from a host bearing at least
2 one mutant CalDAG-GEF gene.

1 95. A method as in claim 92 wherein said cell is explanted from a transgenic animal of
2 any one of claims 55-61.

1 96. A method as in claim 86 wherein said cell is a cell in a live animal.

1 97. A method as in claim 96 wherein said cell is a cell of a transgenic animal of any one
2 of claims 55-61.

1 98. A method as in claim 86 wherein said cell is in a human subject in a clinical trial.

1 99. A method as in claim 87 wherein measurement of said marker indicates a difference
2 between cells bearing an expressed mutant cAMP-GEF gene and otherwise identical cells free of
3 an expressed mutant cAMP-GEF gene.

- 67 -

1 100. A method as in claim 87 wherein said change comprises a change in a non-specific
2 marker of cell physiology selected from the group consisting of pH, intracellular calcium, cyclic
3 AMP levels, GTP/GDP ratios, phosphatidylinositol activity, and protein phosphorylation.

1 101. A method as in claim 87 wherein said change comprises a change in expression of
2 said cAMP-GEF.

1 102. A method as in claim 87 wherein said change comprises a change in occurrence or
2 rate of apoptosis or cell death.

1 103. A method as in claim 87 wherein said cell is a cell cultured *in vitro*.

1 104. A method as in claim 103 wherein said cell is a transformed host cell of any one of
2 claims 50-54.

1 105. A method as in claim 103 wherein said cell is explanted from a host bearing at least
2 one mutant cAMP-GEF gene.

1 106. A method as in claim 103 wherein said cell is explanted from a transgenic animal of
2 any one of claims 55-61.

1 107. A method as in claim 87 wherein said cell is a cell in a live animal.

1 108. A method as in claim 107 wherein said cell is a cell of a transgenic animal of any one
2 of claims 55-61.

1 109. A method as in claim 87 wherein said cell is in a human subject in a clinical trial.

1 110. A diagnostic method for determining if a subject bears a mutant CalDAG-GEF gene
2 comprising the steps of

3 providing a biological sample of said subject; and

4 detecting in said sample a mutant CalDAG-GEF nucleic acid, a mutant CalDAG-GEF
5 protein, or a mutant CalDAG-GEF activity.

1 111. A method as in claim 111, wherein a mutant CalDAG-GEF nucleic acid is detected
2 by an assay selected from the group consisting of direct nucleotide sequencing, probe specific
3 hybridization, restriction enzyme digest and mapping, PCR mapping, ligase-mediated PCR
4 detection, RNase protection, electrophoretic mobility shift detection, and chemical mismatch
5 cleavage.

1 112. A method as in claim 110, wherein a mutant CalDAG-GEF protein is detected by an
2 assay selected from the group consisting of an immunoassay, a protease assay, and an
3 electrophoretic mobility assay.

- 68 -

1 113. A diagnostic method for determining if a subject bears a mutant cAMP-GEF gene
2 comprising the steps of

3 providing a biological sample of said subject; and

4 detecting in said sample a mutant cAMP-GEF nucleic acid, a mutant cAMP-GEF
5 protein, or a mutant cAMP-GEF activity.

1 114. A method as in claim 113, wherein a mutant cAMP-GEF nucleic acid is detected by
2 an assay selected from the group consisting of direct nucleotide sequencing, probe specific
3 hybridization, restriction enzyme digest and mapping, PCR mapping, ligase-mediated PCR
4 detection, RNase protection, electrophoretic mobility shift detection, and chemical mismatch
5 cleavage.

1 115. A method as in claim 113, wherein a mutant cAMP-GEF protein is detected by an
2 assay selected from the group consisting of an immunoassay, a protease assay, and an
3 electrophoretic mobility assay.

1 116. A pharmaceutical preparation comprising a substantially pure CalDAG-GEF protein
2 and a pharmaceutically acceptable carrier.

1 117. A pharmaceutical preparation comprising a substantially pure cAMP-GEF protein and
2 a pharmaceutically acceptable carrier.

1 118. A pharmaceutical preparation comprising an expression vector operably encoding a
2 protein selected from the group consisting of a CalDAG-GEF protein, and a cAMP-GEF protein,
3 wherein said expression vector may express said CalDAG-GEF protein or said cAMP-GEF
4 protein in a human subject, and a pharmaceutically acceptable carrier.

1 119. A pharmaceutical preparation comprising an expression vector operably encoding a
2 CalDAG-GEF antisense sequence, wherein said expression vector may express said CalDAG-
3 GEF antisense sequence in a human subject, and a pharmaceutically acceptable carrier.

1 120. A pharmaceutical preparation comprising an expression vector operably encoding a
2 cAMP-GEF antisense sequence, wherein said expression vector may express said cAMP-GEF
3 antisense sequence in a human subject, and a pharmaceutically acceptable carrier.

1 121. A pharmaceutical preparation comprising a substantially pure antibody, and a
2 pharmaceutically acceptable carrier,

3 wherein said antibody selectively binds to a mutant protein selected from the group
4 consisting of a mutant CalDAG-GEF protein, and a mutant cAMP-GEF protein.

- 69 -

1 122. A pharmaceutical preparation as in claim 121 wherein said preparation is essentially
2 free of an antibody which selectively binds a normal CalDAG-GEF protein.

1 123. A pharmaceutical preparation as in claim 121 wherein said preparation is essentially
2 free of an antibody which selectively binds a normal cAMP-GEF protein.

1 124. A pharmaceutical preparation comprising a substantially pure preparation of an
2 antigenic determinant of a mutant CalDAG-GEF protein or a mutant cAMP-GEF protein.

1 125. A pharmaceutical preparation as in claim 124 wherein said preparation is essentially
2 free of an antigenic determinant of a normal CalDAG-GEF protein.

1 126. A pharmaceutical preparation as in claim 124 wherein said preparation is essentially
2 free of an antigenic determinant of a normal cAMP-GEF protein.

1 127. A method of treatment for a patient bearing a mutant CalDAG-GEF gene comprising
2 the step of administering to said patient a therapeutically effective amount of the pharmaceutical
3 preparation of claim 116.

1 128. A method for identifying compounds according to claim 82, wherein the CalDAG-
2 GEF component is a CalDAG-GEF domain selected from the group consisting of SCR1, SCR2,
3 SCR3, DAG-binding and an EF hand domain.

1 129. A substantially pure preparation of a polypeptide comprising a domain selected from
2 the group consisting of a CalDAG-GEF SCR1 domain, a CalDAG-GEF SCR2 domain,
3 CalDAG-GEF SCR3 domain, CalDAG-GEF DAG-binding domain, CalDAG-GEF EF hand
4 domain.

1 130. A substantially pure preparation of a polypeptide comprising a domain selected from
2 the group consisting of a cAMP-GEF SCR1 domain, a cAMP-GEF SCR2 domain, cAMP-GEF
3 SCR3 domain, cAMP-GEF cAMP-binding domain.




FIG. 1




FIG. 2A




FIG. 2B




FIG. 2D-1	FIG. 2D-2
--------------	--------------

FIG. 2D

hCaIDAG-GEFI	MAGTLDLDKGC	TVEEELLRGC	IEAFDDSGK	VRDPQLVR	MFMMH	PWYIPSS
mCaIDAG-GEFI	MA[S]TLDLDKGC	TVEEELLRGC	IEAFDDSGK	VRDPQLVR	MFMMH	PWYIPSS
hCaIDAG-GEFI	KE L K A L L D Q E G N R R H S S L I D I D S V P T Y K W K R Q V T Q R N P V G Q K K R K M S L L F					
mCaIDAG-GEFI	KE L K A L L D Q E G N R R H S S L I D I [E] S V P T Y K W K R Q V T Q R N P V [E] Q K K R K M S L L F					
hCaIDAG-GEFI	N S V S Q W V Q L M I L S K P T A P Q R A L V I T H F V H V A E K L L Q L Q N F N T L M A V V G G L					
mCaIDAG-GEFI	N S V S Q W V Q L M I L S K P T A [T] Q R A L V I T H F V H V A E K L L Q L Q N F N T L M A V V G G L					
hCaIDAG-GEFI	F P I L G V H L K D L V A L Q L A L P D W L D P A R T R L N G A K M K Q L F S I L E E L A M V T S L					
mCaIDAG-GEFI	F P I L G V H L K D L V A L Q L A L P D W L D P [G] R T R L N G A K M [R] Q L F S I L E E L A M V T S L					
hCaIDAG-GEFI	P P R P P V L E E W Y S A A K P K L D Q A L V V E H I E K M V E S V F R N F D V D G D G H I S Q E E					
mCaIDAG-GEFI	P P R P P V L E E W Y S [Y] A K P K L D Q A L V [A] E H I E K M V E S V F R N F D V D G D G H I S Q E E					
hCaIDAG-GEFI	F Q E S N S L R P V A C R H C K A L I L G I Y K Q G L K C R A C G V N C H K Q C K D R R L S L E C R R					
mCaIDAG-GEFI	F Q E S N S L R P V A C R H C K A L I L G I Y K Q G L K C R A C G V N C H K Q C K [E] R L S L E C R R					
hCaIDAG-GEFI	EDGVFDIHL					
mCaIDAG-GEFI	EDGVFDIHL					

FIG. 2D-1

Q L A A K L L H I Y Q Q S R K D N S N S L Q Y K T C H L V R Y W I S A F P A E F D L N P E L A E Q I	100
Q L A S K L L H F Y Q Q S R K D N S N S L Q Y K T C H L V R Y W V S A F P A E F D L N P E L A E P I	100
D H L E P M E L A E H L T Y L E Y R S F C K I L F Q D D Y H S F V T H G C T V D N P V L E R F I S L F	200
D H L E P M E L A E H L T Y L E Y R S F C K I L F Q D D Y H S F V T H G C T V D N P V L E R F I S L F	200
S H S S I S R L K E T H S H V S P E T I K L W E G L T E L V T A T G N Y G N Y R R R L A A C V G F R	300
S H S S I S R L K E T H S H V S P D T I K L W E G L T E L V T A T G N Y S N Y R R R L A A C V G F R	300
R P P V Q A N P D L L S L L T V S L D Q Y Q T E D E L Y Q L S L Q R E P R S K S S P T S P T S C T P	400
R P P V Q A K P S L L S L L T V S L D Q Y Q T E D E L Y Q L S L Q K E P R S K S S P T S P T S C T P	400
F Q I I R G N F P Y L S A F G D L O Q N Q D G C I S R E E M V S Y F L R S S S V L G G R M G F V H N	500
F Q I I R G N F P Y L S A F G D L O Q N Q D G C I S R E E M I S Y F L R S S S V L G G R M G F V H N	500
R A Q S V S L E G S A P S P S P M H S H H H R A F S F S L P R P G R R G S R P P E I R E E E V Q T V	600
R A Q S V S L E G S A P S P S P T H T H H H R A F S F S L P R P G R R G S R P P E I R E E E V Q S V	599
	609
	608

FIG. 2D-2

6/12

F2583.3	A	V	E	K	H	Y	D	H	D	R	D	G	F	I	S	Q	E	E	F	Q	473
hCaIDAG-GEFI	S	V	F	R	N	F	D	V	D	G	D	G	H	I	S	Q	E	E	F	Q	452
hCaIDAG-GEFII	S	V	F	K	N	Y	D	H	D	Q	D	G	Y	I	S	Q	E	E	F	E	447
hCalmodulin	E	A	F	S	L	F	D	K	D	G	D	G	T	I	T	K	F	L	G		34
hCalbindin D28K	L	M	L	K	L	F	D	S	N	N	D	G	K	L	E	L	T	E	M	A	168
hCalcieneurin B	F	A	F	R	I	Y	D	M	D	K	D	G	Y	I	S	N	G	E	L	F	113
hParvalbumin α	K	V	F	H	M	L	D	K	D	K	S	G	F	I	E	E	D	E	L	G	65
hTroponin C	E	C	F	R	I	F	D	R	K	A	D	G	Y	I	D	P	E	E	L	A	117

FIG. 2E

F2583.3	H	N	F	H	E	T	T	F	L	T	P	T	T	C	N	H	C	N	K	L	L	W	G	I	L	R	Q	G	F	K	C	K	D	C	G	L	A	V	H	S	C	C	K	S	N	A	V	A	E	C	570
hCaIDAG-GEFI	H	N	F	Q	E	S	N	S	L	R	P	V	A	C	R	H	C	K	A	L	I	L	G	I	Y	K	Q	G	L	K	C	R	A	C	G	V	N	C	H	K	Q	C	K	D	R	L	S	V	E	C	548
hCaIDAG-GEFII	H	N	F	Q	E	T	T	Y	L	K	P	T	F	C	D	N	C	A	G	F	L	W	G	V	I	K	Q	G	Y	R	C	C	D	C	C	M	N	C	H	K	Q	C	K	D	L	V	F	E	C	542	
hPKC α	H	K	F	I	A	R	F	F	K	Q	P	T	F	C	S	H	C	T	D	F	I	W	G	F	G	K	Q	G	F	Q	C	Q	V	C	C	F	V	V	H	K	R	C	H	E	F	V	T	F	S	C	86
hPKC β 1	H	K	F	T	A	R	F	F	K	Q	P	T	F	C	S	H	C	T	D	F	I	W	G	F	G	K	Q	G	F	Q	C	Q	V	C	C	F	V	V	H	K	R	C	H	E	F	V	T	F	S	C	86
hPKC γ	H	K	F	T	A	R	F	F	K	Q	P	T	F	C	S	H	C	T	D	F	I	W	G	I	G	K	Q	G	L	Q	C	Q	V	C	S	F	V	V	H	R	C	H	E	F	V	T	F	E	C	85	

FIG. 2F

7/12




FIG. 3A




FIG. 3B

	10	20	30	40	
hcAMP-GEFI	R A Q L L K K F I K L A A H L K E Q K N V N S F F A V M F G L S N S P I S R L A N T W	731			
rcAMP-GEFI	R A Q L L R K F I K L A A H L K E Q K N L N S F F A V M F G L S N S A I S R L A N T W	734			
hcAMP-GEFII	R V Q L L K K F I K L I A A H C K E Y K N L N S F F A I V M G L S N I A V S R L A L T W	882			
rcAMP-GEFII	R V Q L L K K C I K L I A A H C K E Y K N L N S F F G I V M G L S N V A E S R L A L T W	307			
celcAMP-GEF	R M E I L K K F I S I A T I A R E Y R D L L T V F A I T L G L S M T S I S R L L T W	1101			
hCaIDEG-GEFI	R A L V I T H F V H V A E K L L Q L Q N F N T L M A V V G G L S H S S I S R L K E T H	262			
hCaIDAG-GEFII	R A E V F I K F I Q V A Q K L N Q L Q N F N T L M A V I G G L S C H S S I S R L K K T S	313			
C3G	R E R L L K F I K I M K H L K K L M N F N S Y L A I L S A L D S A P I R R L E - W	946			
CDC25	R S K L T Q Y F V T V A Q H C K E L M N F S S M T A I V S A L Y S S P I Y R L K K T W	1415			
rRas-GRF	R A S T K W V A V A D I C R C L M N Y N A V L E I T S S I N R S A I F R L K K T W	1119			
hSos1	R V A V S R I E I L Q V F Q E L N N F N Q V L E V S A M N S P V Y R L D K T F	890			
BUD5	Q T N T I S Y W L Q V A L A C L Y L R N L N S L A S I T S L Q N H S I R R L S L P I	406			

FIG. 3D

9/12





FIG. 3F-1	FIG. 3F-2	FIG. 3F-3
-----------	-----------	-----------

FIG. 3F

FIG. 3E

10/12

hcAMP-GEFI	M V L - - - - -
hcAMP-GEFII	M V A A H A A H S S S S A E W I A C L D K
hcAMP-GEFI	- - - - - H Q H P
hcAMP-GEFII	Y A V L A G S L D V K V S E T S S H Q D A
hcAMP-GEFI	- - - - - T P
hcAMP-GEFII	P Y G V M E T G S M M D R I P D K K N T P
hcAMP-GEFI	L V D G I L A L G L G V H S R S Q V V G I
hcAMP-GEFII	L V D W M M Q Q T P C V H S R T Q A V G M
hcAMP-GEFI	S Q R G P D A L L T V A L R K P P G Q R T
hcAMP-GEFII	S Q M G P D A M M R W I L R K P P G Q R T
hcAMP-GEFI	G S N K V V T N G K G L V T T L H K G D D
hcAMP-GEFII	G S N S V V I Y G K G V V C T L H K G D D
hcAMP-GEFI	A S Q G A - G P S R P P T P G R N R Y T V
hcAMP-GEFII	V P A G N R R S N Q G N S Q P Q Q K Y T V
hcAMP-GEFI	G G S E Q E R S T Y V C M E R Q Q I L R L
hcAMP-GEFII	Q G T E Q E K K D Y A L M N K K K V I R L
hcAMP-GEFI	S P Q K K A R M L P V W L P N Q D K P L P
hcAMP-GEFII	A P Q R K H K V L L Q Q F N T G D K R - A
hcAMP-GEFI	G D A I G L Q P D A R G V A T S L G L N E
hcAMP-GEFII	G K K V V L K P N D V S V F T F L T I N G
hcAMP-GEFI	H Y V L G P Q H L R D V T T A K L E R F N
hcAMP-GEFII	Y M T F Q R M N F K K - T T A K L D L F L
hcAMP-GEFI	R L A M T W E R L P H K V R K L Y S A L K
hcAMP-GEFII	R L A L T W E K L P S K F K K F Y A E F K
hcAMP-GEFI	A A R M L H H C R S R N P V P L S F L R S
hcAMP-GEFII	T A R T V R Y Y R S Q - - - P F N F - - -

FIG. 3F-1

11/12

R	R	M	H	R	P	R	S	-	-	-	-	-	-	-	-	-	-	-	-													
R	F	L	E	K	S	S	E	D	V	D	I	I	F	T	R	L	K	E	V	K	A	F	E	K	F	H	P	N	L	L	M	Q
S	-	-	C	I	Q	G	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
V	T	I	C	T	L	G	I	G	T	A	F	G	E	S	I	L	D	N	T	F	R	N	A	T	I	V	T	R	E	S	S	E
L	T	N	S	E	E	S	L	D	F	S	E	S	L	E	Q	A	S	T	E	R	V	L	R	A	G	R	Q	L	K	Q	N	I
L	I	E	P	H	V	P	L	R	P	A	N	T	I	T	K	V	P	S	E	K	T	L	R	A	G	K	I	L	R	N	A	I
C	Q	V	L	L	D	K	G	A	L	C	H	V	K	K	D	W	A	F	Q	D	K	D	A	Q	F	Y	R	F	P	G	P	E
W	Q	V	L	L	K	D	G	V	L	N	H	V	D	Q	E	H	N	F	Q	D	K	-	Y	L	F	Y	R	F	L	D	D	E
D	E	E	L	D	L	I	F	E	E	L	L	H	I	K	A	V	A	N	L	S	N	S	V	K	R	E	L	A	A	V	L	L
V	D	D	L	E	I	I	Y	E	E	L	L	H	I	K	A	L	S	N	L	S	T	T	V	R	R	E	L	A	G	V	L	I
F	G	Q	L	A	L	V	N	D	A	P	R	A	A	T	I	I	L	R	E	Y	N	C	H	F	L	R	V	D	K	Q	D	F
F	G	K	L	A	L	V	N	D	A	P	R	A	A	S	I	V	L	R	E	D	N	C	H	F	L	R	V	D	K	E	D	F
M	S	G	T	P	D	K	I	L	E	L	L	L	E	A	N	G	L	D	S	E	A	H	D	P	K	E	T	T	L	S	D	F
M	S	G	T	P	E	K	I	L	E	H	F	L	E	T	I	R	L	E	A	T	L	N	E	A	T	D	S	V	L	N	D	F
V	S	Q	W	V	A	L	Y	G	S	M	L	H	T	D	P	V	A	T	S	F	L	M	K	L	S	D	L	V	G	R	D	T
V	L	Q	W	A	A	M	Y	G	D	L	L	Q	E	D	D	V	S	K	A	F	L	E	E	F	Y	V	S	V	S	D	D	A
G	S	S	C	A	I	Q	V	G	D	K	V	P	Y	D	I	C	R	P	D	H	S	V	L	T	L	Q	L	P	V	T	A	S
Q	K	R	Q	P	I	R	S	S	D	E	V	L	F	K	V	Y	C	K	D	H	T	T	T	T	T	R	V	P	V	A	T	S
R	L	F	V	V	N	P	Q	E	V	H	E	L	I	P	H	P	D	Q	L	G	P	T	V	G	S	A	E	G	L	D	L	V
R	L	F	A	C	P	R	E	Q	F	D	S	L	T	P	L	P	E	Q	K	G	P	T	V	G	T	V	G	T	F	E	L	M
R	R	F	N	E	L	Q	Y	W	V	A	T	E	L	C	L	C	P	V	P	G	P	R	A	Q	L	L	K	K	F	I	K	L
R	R	F	N	E	I	Q	F	W	V	V	T	E	I	C	L	C	S	Q	L	B	K	R	V	Q	L	L	K	K	F	I	K	I
R	L	L	D	P	S	W	N	H	R	V	Y	R	L	A	L	A	K	L	S	P	P	Y	I	P	F	M	P	L	L	L	K	D
S	L	M	D	P	S	R	N	H	R	A	Y	R	L	T	V	A	K	L	E	P	P	L	I	P	F	M	P	L	L	I	K	D
R	V	S	H	L	K	G	D	S	Q	V																						

FIG. 3F-2

SUBSTITUTE SHEET (RULE 26)

12/12

- - - C S Y Q L L L K - - - - -	20
I C L C G Y Y E N L K K G I T L F R Q G D I G T N W	80
- L R - - - - - W - - - - -	32
L L R I E Q K D F K A L W E K Y R Q Y M A G L L A P	160
L A T C P N L I R D R K Y H L R L Y R Q C C S G R E	92
L S R A P H M I R D R K Y H L K T Y R Q C C V G T E	240
F E - - P V G T H E M - - - - E E E L A E A V A L L	166
H E D A P L P T E E E K K E C D E E L Q D T M L L L	319
F E P H S K A G T V L F S Q G D K G T S W Y X X W K	246
F E S H A K G G T V L F N Q G E E G T S W Y X X L K	399
N R I I K D V E A K T M K L E G K G K V V L V L K R	326
N R I L R D V E A N T V K L K G K D Q D V L V L K K	479
L L T H R V F M P S A Q L C A A L L H H F H V E P A	405
I M M H C V F M P N T Q L C P A L V A H Y H A Q P S	559
R L S N L L R E Q W P E R R R C H R L E N G C G M A	485
R M I A A L K E Q L P E L E K I V K Q I S E - D A K	638
V R E V M A A L A Q E D G W T K G Q V L V K V N S A	565
V K E V I S A V A D K L G S G E G L I I V K M S S G	717
S A K D L A G Q L T D H D W S L F N S I H Q V E L I	645
S S K D L A Y Q M T I Y D W E L F N C V H R L E L I	797
A A H L R R Q K N V N S F F A V M F G L G K S P I S	725
A A H C R R Y K N L N S F F A I V X G L G K I A V S	876
M T F I H E G N H T L W E N L I N F G R M R N M A R	805
M T F T H E G N K T F I D N L V N F G R M R N I A N	956
V Q Q L K V I D N Q R E L S R L S R E L E P	881
V R Q L N V I D N Q R T L S Q N S H R L E P R R F	1011

FIG. 3F-3

SUBSTITUTE SHEET (RULE 26)

SEQUENCE LISTING

<110> Kawasaki, Hiroaki
Graybiel, Ann
Housman, David

<120> Genes Integrating Signal Transduction Pathways

<130> MIT-103

<140>

<141>

<150> US 60/105,507

<151> 1998-10-23

<150> US 60/108,685

<151> 1998-11-16

<160> 18

<170> PatentIn Ver. 2.0

<210> 1

<211> 2250

<212> DNA

<213> Mus musculus

<220>

<221> CDS

<222> (166)..(1989)

<223> CalDAG-GEFI

<400> 1

```

cgaaggatca gaggctgagc tggttcaagt gaacagaagg tctgggaggt gaactgcatt 60
cggggtttgca ttctgaagta aaggacttgg gacaggggta cgaatcgagc actgtgggag 120
gctctgagag tgtaacttgg gtctagccca ctggcaccgg cagcc atg gcg agc act 177
                                     Met Ala Ser Thr
                                     1

ctg gac ctg gac aag ggt tgc acc gtg gag gag ctg ctc cgt ggc tgt 225
Leu Asp Leu Asp Lys Gly Cys Thr Val Glu Glu Leu Leu Arg Gly Cys
5 10 15 20

atc gaa gcc ttt gat gac tct gga aag gtg cga gat cca cag cta gtg 273
Ile Glu Ala Phe Asp Asp Ser Gly Lys Val Arg Asp Pro Gln Leu Val
25 30 35

cgc atg ttt ctc atg atg cac ccc tgg tac ata cct tcc tct cag ctg 321
Arg Met Phe Leu Met Met His Pro Trp Tyr Ile Pro Ser Ser Gln Leu
40 45 50

gct tcg aaa ctg ctc cac ttc tat cag caa tcc cgg aag gac aac tcc 369
Ala Ser Lys Leu Leu His Phe Tyr Gln Gln Ser Arg Lys Asp Asn Ser
55 60 65

aat tcc cta cag gtg aaa acc tgt cac ctg gtc agg tac tgg gtc tca 417
Asn Ser Leu Gln Val Lys Thr Cys His Leu Val Arg Tyr Trp Val Ser
70 75 80

gcc ttc cca gca gag ttc gac ttg aac cca gag ctg gct gaa ccg atc 465
Ala Phe Pro Ala Glu Phe Asp Leu Asn Pro Glu Leu Ala Glu Pro Ile
85 90 95 100

aag gag ctg aag gct ctg tta gac caa gaa gga aac cgc agg cac agc 513
Lys Glu Leu Lys Ala Leu Leu Asp Gln Glu Gly Asn Arg Arg His Ser
105 110 115

agc ctc atc gac atc gag agt gtc ccc acc tac aag tgg aag cgg cag 561
Ser Leu Ile Asp Ile Glu Ser Val Pro Thr Tyr Lys Trp Lys Arg Gln
120 125 130

gtg acc cag cgg aac cct gtg gaa cag aaa aag cgc aag atg tcc ctg 609
Val Thr Gln Arg Asn Pro Val Glu Gln Lys Lys Arg Lys Met Ser Leu
135 140 145

```

- 2 -

ttg	ttt	gat	cac	ttg	gag	cct	atg	gaa	ctg	gca	gaa	cat	ctc	acc	tac	657
Leu	Phe	Asp	His	Leu	Glu	Pro	Met	Glu	Leu	Ala	Glu	His	Leu	Thr	Tyr	
	150					155					160					
ttg	gag	tat	cgg	tcc	ttc	tgc	aag	atc	ctg	ttc	cag	gac	tat	cac	agc	705
Leu	Glu	Tyr	Arg	Ser	Phe	Cys	Lys	Ile	Leu	Phe	Gln	Asp	Tyr	His	Ser	
	165				170					175					180	
ttt	gtg	act	cat	ggc	tgc	act	gta	gac	aat	ccg	gtc	ctg	gag	cga	ttc	753
Phe	Val	Thr	His	Gly	Cys	Thr	Val	Asp	Asn	Pro	Val	Leu	Glu	Arg	Phe	
				185					190					195		
atc	tcc	ctc	ttc	aac	agt	gtc	tct	cag	tgg	gtc	caa	ctc	atg	atc	ctc	801
Ile	Ser	Leu	Phe	Asn	Ser	Val	Ser	Gln	Trp	Val	Gln	Leu	Met	Ile	Leu	
			200					205					210			
agc	aag	ccc	aca	gcc	acg	cag	cgg	gcg	ctg	gtc	atc	aca	cat	ttc	gtg	849
Ser	Lys	Pro	Thr	Ala	Thr	Gln	Arg	Ala	Leu	Val	Ile	Thr	His	Phe	Val	
		215					220					225				
cat	gtg	gca	gag	aag	ctg	ctg	cag	ctg	cag	aac	ttc	aac	acg	ttg	atg	897
His	Val	Ala	Glu	Lys	Leu	Gln	Gln	Leu	Gln	Asn	Phe	Asn	Thr	Leu	Met	
	230					235					240					
gcc	gtc	gtg	gga	ggc	ctg	agc	cac	agc	tcc	atc	tca	cgc	ctc	aag	gag	945
Ala	Val	Val	Gly	Gly	Leu	Ser	His	Ser	Ser	Ile	Ser	Arg	Leu	Lys	Glu	
	245				250					255					260	
acc	cac	agc	cat	gtc	agc	cct	gac	acc	atc	aag	ctc	tgg	gaa	ggc	ctg	993
Thr	His	Ser	His	Val	Ser	Pro	Asp	Thr	Ile	Lys	Leu	Trp	Glu	Gly	Leu	
				265					270					275		
aca	gaa	cta	gtg	aca	gct	act	ggc	aac	tac	agc	aac	tac	cgg	cga	agg	1041
Thr	Glu	Leu	Val	Thr	Ala	Thr	Gly	Asn	Tyr	Ser	Asn	Tyr	Arg	Arg	Arg	
			280					285					290			
ctg	gcg	gcc	tgc	gtg	ggc	ttc	cgc	ttt	cct	atc	ctg	ggc	gtg	cac	ctc	1089
Leu	Ala	Ala	Cys	Val	Gly	Phe	Arg	Phe	Pro	Ile	Leu	Gly	Val	His	Leu	
		295					300					305				
aag	gat	cta	gtg	gct	ctg	cag	ctg	gct	ctg	cct	gac	tgg	ctg	gac	cca	1137
Lys	Asp	Leu	Val	Ala	Leu	Gln	Leu	Ala	Leu	Pro	Asp	Trp	Leu	Asp	Pro	
	310					315					320					
ggc	cgg	acc	cgg	ctc	aat	gga	gcc	aag	atg	agg	cag	ctt	ttc	agc	att	1185
Gly	Arg	Thr	Arg	Leu	Asn	Gly	Ala	Lys	Met	Arg	Gln	Leu	Phe	Ser	Ile	
	325				330					335					340	
ctg	gag	gag	ttg	gct	atg	gtg	acc	agt	ctt	cga	cca	cca	gtg	caa	gcc	1233
Leu	Glu	Glu	Leu	Ala	Met	Val	Thr	Ser	Leu	Arg	Pro	Pro	Val	Gln	Ala	
				345					350					355		
aac	ccc	gac	ctg	ctg	agt	ctg	ctc	acg	gtg	tcc	ctg	gac	cag	tat	cag	1281
Asn	Pro	Asp	Leu	Leu	Ser	Leu	Leu	Thr	Val	Ser	Leu	Asp	Gln	Tyr	Gln	
			360					365					370			
acg	gag	gat	gag	ctg	tat	cag	ctc	tct	ctg	cag	cga	gag	cca	cgt	tcc	1329
Thr	Glu	Asp	Glu	Leu	Tyr	Gln	Leu	Ser	Leu	Gln	Arg	Glu	Pro	Arg	Ser	
		375					380					385				
aag	tca	tcg	ccc	acc	agc	ccc	acc	agc	tgc	acc	ccg	cct	ccc	cgg	ccg	1377
Lys	Ser	Ser	Pro	Thr	Ser	Pro	Thr	Ser	Cys	Thr	Pro	Pro	Pro	Arg	Pro	
	390					395					400					
cct	gtg	ctg	gaa	gag	tgg	acc	tca	gtt	gcc	aag	cct	aag	ctg	gac	caa	1425
Pro	Val	Leu	Glu	Glu	Trp	Thr	Ser	Val	Ala	Lys	Pro	Lys	Leu	Asp	Gln	
					410					415					420	
gcc	ttg	gtg	gcc	gag	cac	att	gag	aag	atg	gtg	gag	tct	gtg	ttc	cgg	1473
Ala	Leu	Val	Ala	Glu	His	Ile	Glu	Lys	Met	Val	Glu	Ser	Val	Phe	Arg	
				425					430					435		
aac	ttt	gac	gtt	gat	ggg	gac	ggt	cac	atc	tcc	cag	gag	gag	ttc	cag	1521
Asn	Phe	Asp	Val	Asp	Gly	Asp	Gly	His	Ile	Ser	Gln	Glu	Glu	Phe	Gln	

- 3 -

440										445					450					
atc	atc	cgg	ggc	aac	ttc	cct	tat	ctc	agc	gcc	ttt	ggg	gac	ttg	gac	1569				
Ile	Ile	Arg	Gly	Asn	Phe	Pro	Tyr	Leu	Ser	Ala	Phe	Gly	Asp	Leu	Asp					
		455					460					465								
cag	aac	cag	gat	ggc	tgc	atc	agc	cgg	gag	gag	atg	att	tcc	tac	ttc	1617				
Gln	Asn	Gln	Asp	Gly	Cys	Ile	Ser	Arg	Glu	Glu	Met	Ile	Ser	Tyr	Phe					
	470					475					480									
ctg	cgc	tcc	agc	tcc	gtg	ctg	gga	ggc	cgc	atg	ggc	ttc	gta	cac	aac	1665				
Leu	Arg	Ser	Ser	Ser	Val	Leu	Gly	Gly	Arg	Met	Gly	Phe	Val	His	Asn					
485					490					495					500					
ttc	cag	gag	agt	aac	tgc	cta	agg	ccg	gtc	gcc	tgc	cga	cac	tgc	aaa	1713				
Phe	Gln	Glu	Ser	Asn	Ser	Leu	Arg	Pro	Val	Ala	Cys	Arg	His	Cys	Lys					
				505					510					515						
gct	ctg	atc	ctg	ggc	atc	tac	aag	cag	ggc	ctc	aaa	tgt	aga	gct	tgt	1761				
Ala	Leu	Ile	Leu	Gly	Ile	Tyr	Lys	Gln	Gly	Leu	Lys	Cys	Arg	Ala	Cys					
			520					525					530							
ggt	gtg	aac	tgc	cac	aag	cag	tgc	aaa	gac	cga	ctg	tca	gtg	gaa	tgt	1809				
Gly	Val	Asn	Cys	His	Lys	Gln	Cys	Lys	Asp	Arg	Leu	Ser	Val	Glu	Cys					
		535				540						545								
cgc	cgc	cgc	gcc	cag	agt	gtg	agc	ctg	gag	ggc	tct	gca	ccc	tct	ccc	1857				
Arg	Arg	Arg	Ala	Gln	Ser	Val	Ser	Leu	Glu	Gly	Ser	Ala	Pro	Ser	Pro					
		550				555					560									
tca	ccc	aca	cat	acc	cac	cat	cgg	gcc	ttc	agc	ttc	tcc	ctg	cct	cgc	1905				
Ser	Pro	Thr	His	Thr	His	His	Arg	Ala	Phe	Ser	Phe	Ser	Leu	Pro	Arg					
565					570					575					580					
cca	ggc	agg	cgc	agc	tct	cgg	cct	cca	gag	atc	cgt	gaa	gag	gag	gtg	1953				
Pro	Gly	Arg	Arg	Ser	Ser	Arg	Pro	Pro	Glu	Ile	Arg	Glu	Glu	Glu	Val					
				585					590					595						
cag	act	gtg	gaa	gat	ggt	gtg	ttc	gac	atc	cac	tta	taagacgctg				1999				
Gln	Thr	Val	Glu	Asp	Gly	Val	Phe	Asp	Ile	His	Leu									
			600					605												
tgactatcaa	ggactcattc	ctgccttggg	gaaaagactt	ggagcagagc	agggagccag	2059														
ggattctggg	gcaggaggtt	ggggctgaag	gtgggggaag	ttgaagggtg	catgcactga	2119														
aaaaaaggcc	agggctggtg	tccttaaggt	tgtacagact	tctgtgaata	tttgtatttt	2179														
ccagatggaa	taaaaaggcc	cgaataatta	acctcgaaaa	aaaaaaaaaa	aaaaaaaaaa	2239														
aaaaaaaaaa	a					2250														

<210> 2
 <211> 608
 <212> PRT
 <213> Mus musculus

<400> 2
 Met Ala Ser Thr Leu Asp Leu Asp Lys Gly Cys Thr Val Glu Glu Leu
 1 5 10 15
 Leu Arg Gly Cys Ile Glu Ala Phe Asp Asp Ser Gly Lys Val Arg Asp
 20 25 30
 Pro Gln Leu Val Arg Met Phe Leu Met Met His Pro Trp Tyr Ile Pro
 35 40 45
 Ser Ser Gln Leu Ala Ser Lys Leu Leu His Phe Tyr Gln Gln Ser Arg
 50 55 60
 Lys Asp Asn Ser Asn Ser Leu Gln Val Lys Thr Cys His Leu Val Arg
 65 70 75 80
 Tyr Trp Val Ser Ala Phe Pro Ala Glu Phe Asp Leu Asn Pro Glu Leu

85

90

95

SUBSTITUTE SHEET (RULE 26)

- 5 -

485 490 495

Phe Val His Asn Phe Gln Glu Ser Asn Ser Leu Arg Pro Val Ala Cys
500 505 510

Arg His Cys Lys Ala Leu Ile Leu Gly Ile Tyr Lys Gln Gly Leu Lys
515 520 525

Cys Arg Ala Cys Gly Val Asn Cys His Lys Gln Cys Lys Asp Arg Leu
530 535 540

Ser Val Glu Cys Arg Arg Arg Ala Gln Ser Val Ser Leu Glu Gly Ser
545 550 555 560

Ala Pro Ser Pro Ser Pro Thr His Thr His His Arg Ala Phe Ser Phe
565 570 575

Ser Leu Pro Arg Pro Gly Arg Arg Ser Ser Arg Pro Pro Glu Ile Arg
580 585 590

Glu Glu Glu Val Gln Thr Val Glu Asp Gly Val Phe Asp Ile His Leu
595 600 605

<210> 3
<211> 2236
<212> DNA
<213> Homo sapiens

<220>
<221> CDS
<222> (161)..(1987)
<223> CalDAG-GEFI

<400> 3
ggggactcaa ggctggcctg gctcaagtga acagcacgtc caggaggcga cctcggtccgc 60
gggtttgcat tctgggggtgg acgagctggg gggtcggtcc gagcccggtg ggaggctccc 120
ggagcgcagc ctggggcccag cccaccccgc gccggcgggc atg gca ggc acc ctg 175
Met Ala Gly Thr Leu
1 5

gac ctg gac aag ggc tgc acg gtg gag gag ctg ctc cgc ggc tgc atc 223
Asp Leu Asp Lys Gly Cys Thr Val Glu Glu Leu Leu Arg Gly Cys Ile
10 15 20

gaa gcc ttc gat gac tcc ggg aag gtg cgg gac ccg cag ctg gtg cgc 271
Glu Ala Phe Asp Asp Ser Gly Lys Val Arg Asp Pro Gln Leu Val Arg
25 30 35

atg ttc ctc atg atg cac ccc tgg tac atc ccc tcc tct cag ctg gcg 319
Met Phe Leu Met Met His Pro Trp Tyr Ile Pro Ser Ser Gln Leu Ala
40 45 50

gcc aag ctg ctc cac atc tac caa caa tcc cgg aag gac aac tcc aat 367
Ala Lys Leu Leu His Ile Tyr Gln Gln Ser Arg Lys Asp Asn Ser Asn
55 60 65

tcc ctg cag gtg aaa acg tgc cac ctg gtc agg tac tgg atc tcc gcc 415
Ser Leu Gln Val Lys Thr Cys His Leu Val Arg Tyr Trp Ile Ser Ala
70 75 80 85

ttc cca gcg gag ttt gac ttg aac ccg gag ttg gct gag cag atc aag 463
Phe Pro Ala Glu Phe Asp Leu Asn Pro Glu Leu Ala Glu Gln Ile Lys
90 95 100

gag ctg aag gct ctg cta gac caa gaa ggg aac cga cgg cac agc agc 511
Glu Leu Lys Ala Leu Leu Asp Gln Glu Gly Asn Arg Arg His Ser Ser
105 110 115

cta atc gac ata gac agc gtc cct acc tac aag tgg aag cgg cag gtg 559
Leu Ile Asp Ile Asp Ser Val Pro Thr Tyr Lys Trp Lys Arg Gln Val
120 125 130

- 6 -

act Thr 135	cag Gln 135	cgg Arg 135	aac Asn 135	cct Pro 135	gtg Val 140	gga Gly 140	cag Gln 140	aaa Lys 140	aag Lys 145	cgc Arg 145	aag Lys 145	atg Met 145	tcc Ser 145	ctg Leu 145	ttg Leu 145	607
ttt Phe 150	gac Asp 150	cac His 150	ctg Leu 155	gag Glu 155	ccc Pro 155	atg Met 155	gag Glu 160	ctg Leu 160	gag Ala 160	cat His 160	ctc Leu 160	acc Met 160	tac Thr 165	ttg Tyr 165	Leu 165	655
gag Glu 170	tat Tyr 170	cgc Arg 170	tcc Ser 170	ttc Phe 170	tgc Cys 170	aag Lys 175	atc Ile 175	ctg Leu 175	ttt Phe 175	cag Gln 180	gac Asp 180	tat Tyr 180	cac His 180	agt Ser 180	ttc Phe 180	703
gtg Val 185	act Thr 185	cat His 185	ggc Gly 185	tgc Cys 185	act Thr 190	gtg Val 190	gac Asp 190	aac Asn 190	ccc Pro 195	gtc Val 195	ctg Leu 195	gag Glu 195	cgg Arg 195	ttc Phe 195	atc Ile 195	751
tcc Ser 200	ctc Leu 200	ttc Phe 200	aac Asn 200	agc Ser 205	gtc Val 205	tca Ser 205	cag Gln 205	tgg Trp 205	gtg Val 210	cag Gln 210	ctc Leu 210	atg Met 210	atc Ile 210	ctc Leu 210	agc Ser 210	799
aaa Lys 215	ccc Pro 215	aca Thr 215	gcc Ala 220	ccg Pro 220	cag Gln 220	cgg Arg 220	gcc Ala 220	ctg Leu 220	gtc Val 225	atc Ile 225	aca Thr 225	cac His 225	ttt Phe 225	gtc Val 225	cac His 225	847
gtg Val 230	gag Ala 230	gag Glu 230	aag Lys 235	ctg Leu 235	cta Leu 235	cag Gln 235	ctg Leu 235	cag Gln 240	aac Asn 240	ttc Phe 240	aac Asn 240	acg Thr 240	ctg Leu 245	atg Met 245	gca Ala 245	895
gtg Val 250	gtc Val 250	ggg Gly 250	ggc Gly 250	ctg Leu 250	agc Ser 255	cac His 255	agc Ser 255	tcc Ser 255	atc Ile 255	tcc Ser 255	cgc Arg 255	ctc Leu 255	aag Lys 260	gag Glu 260	acc Thr 260	943
cac His 265	agc Ser 265	cac His 265	gtt Val 265	agc Ser 270	cct Pro 270	gag Glu 270	acc Thr 270	atc Ile 270	aag Lys 270	ctc Leu 275	tgg Trp 275	gag Glu 275	ggt Gly 275	ctc Leu 275	acg Thr 275	991
gaa Glu 280	cta Leu 280	gtg Val 280	acg Thr 285	gag Ala 285	aca Thr 285	ggc Gly 285	aac Asn 285	tat Tyr 285	ggc Gly 290	aac Asn 290	tac Tyr 290	cgg Arg 290	cgt Arg 290	cgg Arg 290	ctg Leu 290	1039
gca Ala 295	gcc Ala 295	tgt Cys 295	gtg Val 300	ggc Gly 300	ttc Phe 300	cgc Arg 300	ttc Phe 300	ccg Pro 305	atc Ile 305	ctg Leu 305	ggt Gly 305	gtg Val 305	cac His 305	ctc Leu 305	aag Lys 305	1087
gac Asp 310	ctg Leu 310	gtg Val 315	gcc Ala 315	ctg Leu 315	cag Gln 315	ctg Leu 315	gca Ala 320	ctg Leu 320	cct Pro 320	gac Asp 320	tgg Trp 320	ctg Leu 320	gac Asp 320	cca Pro 325	gcc Ala 325	1135
cgg Arg 330	acc Thr 330	cgg Arg 330	ctc Leu 330	aac Asn 330	ggg Gly 330	gcc Ala 335	aag Lys 335	atg Met 335	aag Lys 335	cag Gln 335	ctc Leu 335	ttt Phe 340	agc Ser 340	atc Ile 340	ctg Leu 340	1183
gag Glu 345	gag Glu 345	ctg Leu 345	gcc Ala 345	atg Met 345	gtg Val 350	acc Thr 350	agc Ser 350	ctg Leu 350	cgg Arg 350	cca Pro 350	cca Pro 355	gta Val 355	cag Gln 355	gcc Ala 355	aac Asn 355	1231
ccc Pro 360	gac Asp 360	ctg Leu 360	ctg Leu 360	agc Ser 365	ctg Leu 365	ctc Leu 365	acg Thr 365	gtg Val 365	tct Ser 370	ctg Leu 370	gat Asp 370	cag Gln 370	tat Tyr 370	cag Gln 370	acg Thr 370	1279
gag Glu 375	gat Asp 375	gag Glu 375	ctg Leu 380	tac Tyr 380	cag Gln 380	ctg Leu 380	tcc Ser 380	ctg Leu 385	cag Gln 385	cgg Arg 385	gag Glu 385	ccg Pro 385	cgc Arg 385	tcc Ser 385	aag Lys 385	1327
tct Ser 390	tcg Ser 390	cca Pro 395	acc Thr 395	agc Ser 395	ccc Pro 395	acg Thr 395	agt Ser 395	tgc Cys 400	acc Thr 400	cca Pro 400	cca Pro 400	ccc Pro 400	cgg Arg 405	ccc Pro 405	ccg Pro 405	1375
gta Val 410	ttg Leu 410	gag Glu 410	gag Glu 410	tgg Trp 410	acc Thr 415	tcg Ser 415	gct Ala 415	gcc Ala 415	aaa Lys 415	ccc Pro 415	aag Lys 420	ctg Leu 420	gat Asp 420	cag Gln 420	gcc Ala 420	1423
ctc Leu 425	gtg Val 425	gtg Val 425	gag Glu 425	cac His 430	atc Ile 430	gag Glu 430	aag Lys 430	atg Met 430	gtg Val 435	gag Glu 435	tct Ser 435	gtg Val 435	ttc Phe 435	cgg Arg 435	aac Asn 435	1471

- 7 -

ttt gac gtc gat ggg gat ggc cac atc tca cag gaa gaa ttc cag atc 1519
 Phe Asp Val Asp Gly Asp Gly His Ile Ser Gln Glu Glu Phe Gln Ile
 440 445 450
 atc cgt ggg aac ttc cct tac ctc agc gcc ttt ggg gac ctc gac cag 1567
 Ile Arg Gly Asn Phe Pro Tyr Leu Ser Ala Phe Gly Asp Leu Asp Gln
 455 460 465
 aac cag gat ggc tgc atc agc agg gag gag atg gtt tcc tat ttc ctg 1615
 Asn Gln Asp Gly Cys Ile Ser Arg Glu Glu Met Val Ser Tyr Phe Leu
 470 475 480 485
 cgc tcc agc tct gtg ttg ggg ggg cgc atg ggc ttc gta cac aac ttc 1663
 Arg Ser Ser Ser Val Leu Gly Gly Arg Met Gly Phe Val His Asn Phe
 490 495 500
 cag gag agc aac tcc ttg cgc ccc gtc gcc tgc cgc cac tgc aaa gcc 1711
 Gln Glu Ser Asn Ser Leu Arg Pro Val Ala Cys Arg His Cys Lys Ala
 505 510 515
 ctg atc ctg ggc atc tac aag cag ggc ctc aaa tgc cga gcc tgt gga 1759
 Leu Ile Leu Gly Ile Tyr Lys Gln Gly Leu Lys Cys Arg Ala Cys Gly
 520 525 530
 gtg aac tgc cac aag cag tgc aag gat cgc ctg tca gtt gag tgt cgg 1807
 Val Asn Cys His Lys Gln Cys Lys Asp Arg Leu Ser Val Glu Cys Arg
 535 540 545
 cgc agg gcc cag agt gtg agc ctg gag ggg tct gca ccc tca ccc tca 1855
 Arg Arg Ala Gln Ser Val Ser Leu Glu Gly Ser Ala Pro Ser Pro Ser
 550 555 560 565
 ccc atg cac agc cac cat cac cgc gcc ttc agc ttc tct ctg ccc cgc 1903
 Pro Met His Ser His His His Arg Ala Phe Ser Phe Ser Leu Pro Arg
 570 575 580
 cct ggc agg cga ggc tcc agg cct cca gag atc cgt gag gag gag gta 1951
 Pro Gly Arg Arg Gly Ser Arg Pro Pro Glu Ile Arg Glu Glu Glu Val
 585 590 595
 cag acg gtg gag gat ggg gtg ttt gac atc cac ttg taatagatgc 1997
 Gln Thr Val Glu Asp Gly Val Phe Asp Ile His Leu
 600 605

tgtggttgga tcaaggactc attcctgcct tggagaaaat acttcaacca gagcagggag 2057
 cctggggggtg tcgggggcagg aggcctgggga tggggggtggg atatgagggt ggcatgcagc 2117
 tgaggggcagg gccagggctg gtgtccctaa gggtgtacag actcttgtga atatttgtat 2177
 tttccagatg gaataaaaag gcccggtgtaa ttaaccttca aaaaaaaaaa aaaaaaaaaa 2236

<210> 4
 <211> 609
 <212> PRT
 <213> Homo sapiens

<400> 4
 Met Ala Gly Thr Leu Asp Leu Asp Lys Gly Cys Thr Val Glu Glu Leu
 1 5 10 15
 Leu Arg Gly Cys Ile Glu Ala Phe Asp Asp Ser Gly Lys Val Arg Asp
 20 25 30
 Pro Gln Leu Val Arg Met Phe Leu Met Met His Pro Trp Tyr Ile Pro
 35 40 45
 Ser Ser Gln Leu Ala Ala Lys Leu Leu His Ile Tyr Gln Gln Ser Arg
 50 55 60
 Lys Asp Asn Ser Asn Ser Leu Gln Val Lys Thr Cys His Leu Val Arg
 65 70 75 80

Tyr	Trp	Ile	Ser	Ala 85	Phe	Pro	Ala	Glu	Phe 90	Asp	Leu	Asn	Pro	Glu 95	Leu
Ala	Glu	Gln	Ile 100	Lys	Glu	Leu	Lys	Ala 105	Leu	Leu	Asp	Gln	Glu 110	Gly	Asn
Arg	Arg	His 115	Ser	Ser	Leu	Ile	Asp 120	Ile	Asp	Ser	Val	Pro 125	Thr	Tyr	Lys
Trp	Lys 130	Arg	Gln	Val	Thr	Gln 135	Arg	Asn	Pro	Val	Gly 140	Gln	Lys	Lys	Arg
Lys 145	Met	Ser	Leu	Leu	Phe 150	Asp	His	Leu	Glu	Pro 155	Met	Glu	Leu	Ala	Glu 160
His	Leu	Thr	Tyr	Leu 165	Glu	Tyr	Arg	Ser	Phe 170	Cys	Lys	Ile	Leu	Phe 175	Gln
Asp	Tyr	His	Ser 180	Phe	Val	Thr	His	Gly 185	Cys	Thr	Val	Asp	Asn 190	Pro	Val
Leu	Glu	Arg 195	Phe	Ile	Ser	Leu	Phe 200	Asn	Ser	Val	Ser	Gln 205	Trp	Val	Gln
Leu	Met 210	Ile	Leu	Ser	Lys	Pro 215	Thr	Ala	Pro	Gln	Arg 220	Ala	Leu	Val	Ile
Thr 225	His	Phe	Val	His 230	Val	Ala	Glu	Lys	Leu	Leu 235	Gln	Leu	Gln	Asn	Phe 240
Asn	Thr	Leu	Met	Ala 245	Val	Val	Gly	Gly	Leu 250	Ser	His	Ser	Ser	Ile 255	Ser
Arg	Leu	Lys	Glu 260	Thr	His	Ser	His	Val 265	Ser	Pro	Glu	Thr	Ile 270	Lys	Leu
Trp	Glu	Gly 275	Leu	Thr	Glu	Leu	Val 280	Thr	Ala	Thr	Gly	Asn 285	Tyr	Gly	Asn
Tyr	Arg 290	Arg	Arg	Leu	Ala	Ala 295	Cys	Val	Gly	Phe	Arg 300	Phe	Pro	Ile	Leu
Gly 305	Val	His	Leu	Lys	Asp 310	Leu	Val	Ala	Leu	Gln 315	Leu	Ala	Leu	Pro	Asp 320
Trp	Leu	Asp	Pro	Ala 325	Arg	Thr	Arg	Leu	Asn 330	Gly	Ala	Lys	Met	Lys 335	Gln
Leu	Phe	Ser	Ile 340	Leu	Glu	Glu	Leu	Ala 345	Met	Val	Thr	Ser	Leu 350	Arg	Pro
Pro	Val	Gln 355	Ala	Asn	Pro	Asp	Leu 360	Leu	Ser	Leu	Leu 365	Thr	Val	Ser	Leu
Asp	Gln 370	Tyr	Gln	Thr	Glu	Asp 375	Glu	Leu	Tyr	Gln	Leu 380	Ser	Leu	Gln	Arg
Glu 385	Pro	Arg	Ser	Lys	Ser 390	Ser	Pro	Thr	Ser	Pro 395	Thr	Ser	Cys	Thr	Pro 400
Pro	Pro	Arg	Pro	Pro 405	Val	Leu	Glu	Glu	Trp 410	Thr	Ser	Ala	Ala	Lys 415	Pro
Lys	Leu	Asp	Gln 420	Ala	Leu	Val	Val	Glu 425	His	Ile	Glu	Lys	Met 430	Val	Glu
Ser	Val	Phe 435	Arg	Asn	Phe	Asp	Val 440	Asp	Gly	Asp	Gly	His 445	Ile	Ser	Gln
Glu	Glu 450	Phe	Gln	Ile	Ile	Arg 455	Gly	Asn	Phe	Pro	Tyr 460	Leu	Ser	Ala	Phe
Gly 465	Asp	Leu	Asp	Gln 470	Asn	Gln	Asp	Gly	Cys	Ile 475	Ser	Arg	Glu	Glu	Met 480

- 9 -

Val Ser Tyr Phe Leu Arg Ser Ser Ser Val Leu Gly Gly Arg Met Gly
 485 490 495
 Phe Val His Asn Phe Gln Glu Ser Asn Ser Leu Arg Pro Val Ala Cys
 500 505 510
 Arg His Cys Lys Ala Leu Ile Leu Gly Ile Tyr Lys Gln Gly Leu Lys
 515 520 525
 Cys Arg Ala Cys Gly Val Asn Cys His Lys Gln Cys Lys Asp Arg Leu
 530 535 540
 Ser Val Glu Cys Arg Arg Arg Ala Gln Ser Val Ser Leu Glu Gly Ser
 545 550 555 560
 Ala Pro Ser Pro Ser Pro Met His Ser His His His Arg Ala Phe Ser
 565 570 575
 Phe Ser Leu Pro Arg Pro Gly Arg Arg Gly Ser Arg Pro Pro Glu Ile
 580 585 590
 Arg Glu Glu Glu Val Gln Thr Val Glu Asp Gly Val Phe Asp Ile His
 595 600 605
 Leu

<210> 5
 <211> 3624
 <212> DNA
 <213> Rattus norvegicus

<220>
 <221> CDS
 <222> (145)..(2529)
 <223> CalDAG-GEFII

<400> 5
 gcgcctgggt cggctcgccg ggctccgaga gtggccgggt ccgggctgcg agggccggcg 60
 gcggccgagg gatgcgccgc tcccggccga gccgatccca ccgctccagg tgagacggct 120
 ccagggggcg agagagccgc ggcc atg gga acc ctg ggc aag gcg aga gag 171
 Met Gly Thr Leu Gly Lys Ala Arg Glu
 1 5
 gct ccg cgg aaa cct tgc cat ggc tcc aga gct ggc ccc aaa gga aga 219
 Ala Pro Arg Lys Pro Cys His Gly Ser Arg Ala Gly Pro Lys Gly Arg 25
 10 15 20
 cta gag gcc aaa tca acc aac agt cct ctc cct gcc cag ccc agc ttg 267
 Leu Glu Ala Lys Ser Thr Asn Ser Pro Leu Pro Ala Gln Pro Ser Leu 40
 30 35 40
 gcc cag atc acc cag ttc cga atg atg gtg tcc ctg gga cat ctg gcc 315
 Ala Gln Ile Thr Gln Phe Arg Met Met Val Ser Leu Gly His Leu Ala 55
 45 50 55
 aaa gga gcc agc ctg gat gat ctt att gac agc tgc att caa tct ttc 363
 Lys Gly Ala Ser Leu Asp Asp Leu Ile Asp Ser Cys Ile Gln Ser Phe 70
 60 65 70
 gat gca gat gga aac ctg tgt cga agt aac cag ctg tta caa gtc atg 411
 Asp Ala Asp Gly Asn Leu Cys Arg Ser Asn Gln Leu Leu Gln Val Met 85
 75 80 85
 cta acc atg cac cga atc atc atc tcc tcg gcc gag ctg ctg caa aaa 459
 Leu Thr Met His Arg Ile Ile Ile Ser Ser Ala Glu Leu Leu Gln Lys 105
 90 95 100 105
 ctc atg aat cta tat aag gac gcc ctg gaa aag aat tct cca gga att 507
 Leu Met Asn Leu Tyr Lys Asp Ala Leu Glu Lys Asn Ser Pro Gly Ile 120
 110 115 120

- 10 -

tgc	ttg	aag	atc	tgc	tat	ttt	gtc	agg	tat	tgg	ata	aca	gaa	ttc	tgg	555
Cys	Leu	Lys	Ile	Cys	Tyr	Phe	Val	Arg	Tyr	Trp	Ile	Thr	Glu	Phe	Trp	
			125					130					135			
atc	atg	ttc	aaa	atg	gat	gcc	agc	ttg	acc	agc	acc	atg	gaa	gag	ttt	603
Ile	Met	Phe	Lys	Met	Asp	Ala	Ser	Leu	Thr	Ser	Thr	Met	Glu	Glu	Phe	
		140					145					150				
cag	gac	ctg	gtg	aaa	gcc	aat	ggc	gag	gag	tcc	cac	tgc	cac	ctc	atc	651
Gln	Asp	Leu	Val	Lys	Ala	Asn	Gly	Glu	Glu	Ser	His	Cys	His	Leu	Ile	
	155					160					165					
gac	acg	aca	caa	att	aat	tct	cga	gac	tgg	tcc	agg	aaa	ctg	act	cag	699
Asp	Thr	Thr	Gln	Ile	Asn	Ser	Arg	Asp	Trp	Ser	Arg	Lys	Leu	Thr	Gln	
170					175				180						185	
agg	ata	aaa	tca	aat	acc	agc	aag	aag	cgg	aaa	gtg	tcc	ctg	ctg	ttt	747
Arg	Ile	Lys	Ser	Asn	Thr	Ser	Lys	Lys	Arg	Lys	Val	Ser	Leu	Leu	Phe	
				190					195					200		
gac	cat	ctt	gaa	cct	gaa	gaa	ctg	tct	gaa	cac	ctc	acc	tac	ctt	gag	795
Asp	His	Leu	Glu	Pro	Glu	Glu	Leu	Ser	Glu	His	Leu	Thr	Tyr	Leu	Glu	
		205						210				215				
ttc	aag	tcc	ttc	cga	cgg	ata	tct	ttc	tct	gat	tat	cag	aat	tac	ctt	843
Phe	Lys	Ser	Phe	Arg	Arg	Ile	Ser	Phe	Ser	Asp	Tyr	Gln	Asn	Tyr	Leu	
		220					225					230				
gta	aac	agc	tgc	gta	aag	gag	aac	ccc	acc	atg	gag	cgg	tcc	att	gcc	891
Val	Asn	Ser	Cys	Val	Lys	Glu	Asn	Pro	Thr	Met	Glu	Arg	Ser	Ile	Ala	
	235					240					245					
ctg	tgc	aat	ggc	atc	tcc	cag	tgg	gta	caa	ctg	atg	gtt	ctc	agc	cgt	939
Leu	Cys	Asn	Gly	Ile	Ser	Gln	Trp	Val	Gln	Leu	Met	Val	Leu	Ser	Arg	
250					255				260						265	
ccc	acc	cca	cag	ctc	cgg	gca	gag	gtc	ttc	atc	aag	ttc	atc	cat	gtg	987
Pro	Thr	Pro	Gln	Leu	Arg	Ala	Glu	Val	Phe	Ile	Lys	Phe	Ile	His	Val	
				270					275					280		
gct	cag	aag	ctc	cac	cag	cta	cag	aac	ttc	aac	acg	cta	atg	gct	gtg	1035
Ala	Gln	Lys	Leu	His	Gln	Leu	Gln	Asn	Phe	Asn	Thr	Leu	Met	Ala	Val	
			285					290					295			
atc	ggg	gga	ctg	tgt	cac	agc	tcc	atc	tcc	agg	ctc	aag	gag	aca	agt	1083
Ile	Gly	Gly	Leu	Cys	His	Ser	Ser	Ile	Ser	Arg	Leu	Lys	Glu	Thr	Ser	
		300					305					310				
tca	cat	gtc	cca	cat	gag	atc	aat	aag	gtt	ctg	ggt	gag	atg	act	gaa	1131
Ser	His	Val	Pro	His	Glu	Ile	Asn	Lys	Val	Leu	Gly	Glu	Met	Thr	Glu	
		315				320					325					
ctg	ctg	tcc	tcc	tgc	aga	aac	tat	gac	aac	tac	agg	cga	gcc	tat	ggt	1179
Leu	Leu	Ser	Ser	Cys	Arg	Asn	Tyr	Asp	Asn	Tyr	Arg	Arg	Ala	Tyr	Gly	
330					335					340					345	
gag	tgc	acc	cac	ttc	aaa	atc	ccc	ata	ctg	ggt	gtg	cac	ctc	aag	gac	1227
Glu	Cys	Thr	His	Phe	Lys	Ile	Pro	Ile	Leu	Gly	Val	His	Leu	Lys	Asp	
				350					355					360		
ctc	ata	tcc	cta	tat	gaa	gcc	atg	cct	gac	tac	ctg	gaa	gac	ggg	aag	1275
Leu	Ile	Ser	Leu	Tyr	Glu	Ala	Met	Pro	Asp	Tyr	Leu	Glu	Asp	Gly	Lys	
			365					370					375			
gtg	aat	gtc	caa	aag	ctc	ctg	gcc	ctt	tac	aat	cac	atc	aat	gag	ttg	1323
Val	Asn	Val	Gln	Lys	Leu	Leu	Ala	Leu	Tyr	Asn	His	Ile	Asn	Glu	Leu	
		380					385					390				
gtc	cag	ctg	cag	gac	gtg	gcc	cca	cca	ttg	gat	gcc	aac	aag	gac	ctg	1371
Val	Gln	Leu	Gln	Asp	Val	Ala	Pro	Pro	Leu	Asp	Ala	Asn	Lys	Asp	Leu	
		395				400					405					
gtg	cac	ctg	ctg	acg	tta	tcc	ctg	gat	cta	tac	tac	acc	gaa	gat	gaa	1419
Val	His	Leu	Leu	Thr	Leu	Ser	Leu	Asp	Leu	Tyr	Tyr	Thr	Glu	Asp	Glu	
410					415					420					425	

- 11 -

atc	tat	gag	ctt	tcc	tac	gcc	cgt	gaa	cca	agg	aac	cac	agg	gcc	ccg	1467
Ile	Tyr	Glu	Leu	Ser	Tyr	Ala	Arg	Glu	Pro	Arg	Asn	His	Arg	Ala	Pro	
				430					435					440		
cca	ctg	aca	cct	tcg	aag	cca	cca	gtt	gta	gtg	gac	tgg	gcc	tct	gga	1515
Pro	Leu	Thr	Pro	Ser	Lys	Pro	Pro	Val	Val	Val	Asp	Trp	Ala	Ser	Gly	
			445					450					455			
gtg	tct	ccc	aaa	cct	gac	ccg	aag	acc	atc	agc	aaa	cac	gtc	caa	agg	1563
Val	Ser	Pro	Lys	Pro	Asp	Pro	Lys	Thr	Ile	Ser	Lys	His	Val	Gln	Arg	
			460				465					470				
atg	gtg	gat	tct	gtc	ttt	aag	aac	tat	gat	ctc	gac	cag	gat	gga	tat	1611
Met	Val	Asp	Ser	Val	Phe	Lys	Asn	Tyr	Asp	Leu	Asp	Gln	Asp	Gly	Tyr	
			475			480					485					
atc	tct	cag	gag	gag	ttt	gaa	aag	att	gct	gcg	agc	ttt	cca	ttt	tcc	1659
Ile	Ser	Gln	Glu	Glu	Phe	Glu	Lys	Ile	Ala	Ala	Ser	Phe	Pro	Phe	Ser	
					495					500					505	
ttc	tgt	gtg	atg	gac	aaa	gat	agg	gag	ggc	ctc	atc	agc	aga	gac	gag	1707
Phe	Cys	Val	Met	Asp	Lys	Asp	Arg	Glu	Gly	Leu	Ile	Ser	Arg	Asp	Glu	
				510					515					520		
atc	aca	gcc	tac	ttc	atg	agg	gcc	agc	tcc	atc	tat	tcc	aag	ctg	ggc	1755
Ile	Thr	Ala	Tyr	Phe	Met	Arg	Ala	Ser	Ser	Ile	Tyr	Ser	Lys	Leu	Gly	
			525				530						535			
cta	ggc	ttt	cca	cac	aac	ttt	caa	gag	acc	act	tac	ctg	aag	ccc	acc	1803
Leu	Gly	Phe	Pro	His	Asn	Phe	Gln	Glu	Thr	Thr	Tyr	Leu	Lys	Pro	Thr	
			540				545					550				
ttc	tgt	gac	aac	tgt	gct	ggc	ttt	ctc	tgg	ggt	gtg	atc	aag	caa	ggc	1851
Phe	Cys	Asp	Asn	Cys	Ala	Gly	Phe	Leu	Trp	Gly	Val	Ile	Lys	Gln	Gly	
						560					565					
tat	cgc	tgt	aaa	gac	tgt	ggg	atg	aac	tgc	cac	aaa	cag	tgc	aaa	gac	1899
Tyr	Arg	Cys	Lys	Asp	Cys	Gly	Met	Asn	Cys	His	Lys	Gln	Cys	Lys	Asp	
						575				580					585	
ctg	gta	gtg	ttt	gag	tgc	aag	aaa	cga	tcc	aag	agc	ccg	gcg	gta	tcc	1947
Leu	Val	Val	Phe	Glu	Cys	Lys	Lys	Arg	Ser	Lys	Ser	Pro	Ala	Val	Ser	
				590					595					600		
aca	gaa	aac	atc	agc	tct	gtg	gtg	cca	atg	tcc	act	ctt	tgt	cca	ctg	1995
Thr	Glu	Asn	Ile	Ser	Ser	Val	Val	Pro	Met	Ser	Thr	Leu	Cys	Pro	Leu	
			605					610					615			
gga	acc	aaa	gat	ctg	ctc	cat	gca	ccc	gaa	gaa	gga	tct	ttc	att	ttc	2043
Gly	Thr	Lys	Asp	Leu	Leu	His	Ala	Pro	Glu	Glu	Gly	Ser	Phe	Ile	Phe	
			620				625					630				
cag	aat	gga	gag	gtt	gtg	gac	cac	agt	gag	gag	agc	aag	gat	agg	acc	2091
Gln	Asn	Gly	Glu	Val	Val	Asp	His	Ser	Glu	Glu	Ser	Lys	Asp	Arg	Thr	
						640					645					
atc	atg	ctc	ttg	ggc	gta	tcc	tca	cag	aaa	att	tca	gtt	cgg	ctg	aag	2139
Ile	Met	Leu	Leu	Gly	Val	Ser	Ser	Gln	Lys	Ile	Ser	Val	Arg	Leu	Lys	
					655					660					665	
agg	act	gtt	gcc	cac	aag	acc	acc	cag	aca	gaa	tca	ttt	cct	tgg	gtt	2187
Arg	Thr	Val	Ala	His	Lys	Thr	Thr	Gln	Thr	Glu	Ser	Phe	Pro	Trp	Val	
				670				675						680		
ggc	ggc	gag	atg	ccc	cct	ggt	cac	ttt	gtg	ctg	act	tct	cca	aga	aag	2235
Gly	Gly	Glu	Met	Pro	Pro	Gly	His	Phe	Val	Leu	Thr	Ser	Pro	Arg	Lys	
			685					690					695			
tca	gca	caa	ggg	gct	ctt	tat	gtg	cac	agt	cca	gca	tct	ccg	tgc	ccc	2283
Ser	Ala	Gln	Gly	Ala	Leu	Tyr	Val	His	Ser	Pro	Ala	Ser	Pro	Cys	Pro	
			700				705					710				
agc	cca	gca	ctg	gtc	cgg	aag	cgg	gca	ttc	gtc	aag	tgg	gag	aac	aaa	2331
Ser	Pro	Ala	Leu	Val	Arg	Lys	Arg	Ala	Phe	Val	Lys	Trp	Glu	Asn	Lys	

- 12 -

715	720	725	
gag tcc ctt atc aaa cca aaa cca gag ctt cac ctc agg ctc cgg acc			2379
Glu Ser Leu Ile Lys Pro Lys Pro Glu Leu His Leu Arg Leu Arg Thr			
730	735	740	745
tac caa gaa ctg gaa cag gag gta aat acc ctg agg gca gat aac gat			2427
Tyr Gln Glu Leu Glu Gln Glu Val Asn Thr Leu Arg Ala Asp Asn Asp			
	750	755	760
gct ctg aag atc cag ctg aag tat gca cag aaa caa ata gaa tcc ctg			2475
Ala Leu Lys Ile Gln Leu Lys Tyr Ala Gln Lys Gln Ile Glu Ser Leu			
	765	770	775
cag ctt ggc aaa agc aat cac gtc tta gca cag atg gac cac ggt gat			2523
Gln Leu Gly Lys Ser Asn His Val Leu Ala Gln Met Asp His Gly Asp			
	780	785	790
ggt act taatccagaa attcaaggaa cagaatctgc agacggggttt actgggatct			2579
Gly Thr			
795			
cacttcaaaa ctgattgcag aggttcagca actttagact gattgaacttt taaagggcag			2639
agatagccac tgttattggt gtccttggtt tttccctaa ctgtcttttaa tgtgagtcgt			2699
gggttttcag tcagttgagt aaaaggaag aaaagttcca gcatgtaaaa cactgagcag			2759
tcataattcta accttttctt tcttttactg aaaccaatac caaacatgtg ctaaaataag			2819
agtatagttt cctgacagtg tctcaggcag cctgctttac tgattcgcac ttagataaag			2879
agccctggt gaatcaattt gctgccttcc ccagggttcc tgcaaactgg gagttacttt			2939
gttctaccgg aaaacctgct cataatggaa acaataccct aaaagtgggtg actttgacat			2999
gcctgcattg tttgtgaagc tgatccctac cttattcact acctcagatc tcaagagcct			3059
ctttccctgt cctttacttg cccctattt cctcccttgc ttgtaggcac atcagcatat			3119
ctacctagaa gtgacctcct agagatgtag cctgtgttta aatccagagc ttcttaattg			3179
aacttgacat tgtctgattg caggccaact tatagctcta ggtcttagtc attatgaaat			3239
acaaaataaa actaccata atcatcaatt accatgtcat acacagaact cattatctaa			3299
gtcaaacgac gtaaacacgc ctgtccagat agtctttctt tagtggatag aattaacaat			3359
ctgtgacctt ttaaaagaca tgggtctgtga acagtgttta gttcttcacg ttctagtctc			3419
tctctctctc tctctctctc tctctctctc tctctctctc tctctcaagt aaactatcgt			3479
aaggctctct tttgaactga atctgtgctt aaaattgtct tcacttttat tatctacaaa			3539
taagctatgg gaggcgatgg ctttaacagc tgacagcatt tacctatgtg tagaatatgt			3599
gtatatagta tcaggcatgc atgga			3624
 <210> 6			
<211> 795			
<212> PRT			
<213> Rattus norvegicus			
 <400> 6			
Met Gly Thr Leu Gly Lys Ala Arg Glu Ala Pro Arg Lys Pro Cys His			
1	5	10	15
Gly Ser Arg Ala Gly Pro Lys Gly Arg Leu Glu Ala Lys Ser Thr Asn			
	20	25	30
Ser Pro Leu Pro Ala Gln Pro Ser Leu Ala Gln Ile Thr Gln Phe Arg			
	35	40	45
Met Met Val Ser Leu Gly His Leu Ala Lys Gly Ala Ser Leu Asp Asp			

- 13 -

50	55	60
Leu Ile Asp Ser Cys 65	Ile Gln Ser Phe Asp 70	Ala Asp Gly Asn Leu Cys 75 80
Arg Ser Asn Gln 85	Leu Gln Val Met 90	Leu Thr Met His Arg Ile Ile 95
Ile Ser Ser 100	Ala Glu Leu Leu Gln 105	Lys Leu Met Asn Leu Tyr Lys Asp 110
Ala Leu Glu 115	Lys Asn Ser Pro Gly 120	Ile Cys Leu Lys Ile Cys Tyr Phe 125
Val Arg Tyr Trp 130	Ile Thr Glu Phe Trp 135	Ile Met Phe Lys Met Asp Ala 140
Ser Leu Thr Ser Thr 145	Met Glu Glu Phe Gln 150	Asp Leu Val Lys Ala Asn 155 160
Gly Glu Glu Ser 165	His Cys His Leu Ile 170	Asp Thr Thr Gln Ile Asn Ser 175
Arg Asp Trp 180	Ser Arg Lys Leu Thr 185	Gln Arg Ile Lys Ser Asn Thr Ser 190
Lys Lys Arg 195	Lys Val Ser Leu Leu 200	Phe Asp His Leu Glu Pro Glu Glu 205
Leu Ser Glu His 210	Leu Thr Tyr Leu Glu 215	Phe Lys Ser Phe Arg Arg Ile 220
Ser Phe Ser Asp Tyr 225	Gln Asn Tyr Leu Val 230	Asn Ser Cys Val Lys Glu 235 240
Asn Pro Thr Met 245	Glu Arg Ser Ile Ala 250	Leu Cys Asn Gly Ile Ser Gln 255
Trp Val Gln 260	Leu Met Val Leu Ser 265	Arg Pro Thr Pro Gln Leu Arg Ala 270
Glu Val Phe 275	Ile Lys Phe Ile His 280	Val Ala Gln Lys Leu His Gln Leu 285
Gln Asn Phe Asn Thr 290	Leu Met Ala Val Ile 295	Gly Gly Leu Cys His Ser 300
Ser Ile Ser Arg 305	Leu Lys Glu Thr Ser 310	Ser His Val Pro His Glu Ile 315 320
Asn Lys Val Leu 325	Gly Glu Met Thr Glu 330	Leu Leu Ser Ser Cys Arg Asn 335
Tyr Asp Asn Tyr 340	Arg Arg Ala Tyr 345	Gly Glu Cys Thr His Phe Lys Ile 350
Pro Ile Leu Gly 355	Val His Leu Lys 360	Asp Leu Ile Ser Leu Tyr Glu Ala 365
Met Pro Asp Tyr 370	Leu Glu Asp Gly 375	Lys Val Asn Val Gln Lys Leu Leu 380
Ala Leu Tyr Asn His 385	Ile Asn Glu Leu Val 390	Gln Leu Gln Asp Val Ala 395 400
Pro Pro Leu Asp 405	Ala Asn Lys Asp Leu 410	Val His Leu Leu Thr Leu Ser 415
Leu Asp Leu Tyr 420	Tyr Thr Glu Asp 425	Glu Ile Tyr Glu Leu Ser Tyr Ala 430
Arg Glu Pro Arg 435	Asn His Arg Ala 440	Pro Pro Leu Thr Pro Ser Lys Pro 445
Pro Val Val Val 450	Asp Trp Ala Ser 455	Gly Val Ser Pro Lys Pro Asp Pro 460

- 14 -

450					455					460					
Lys	Thr	Ile	Ser	Lys	His	Val	Gln	Arg	Met	Val	Asp	Ser	Val	Phe	Lys
465					470					475					480
Asn	Tyr	Asp	Leu	Asp	Gln	Asp	Gly	Tyr	Ile	Ser	Gln	Glu	Glu	Phe	Glu
				485					490					495	
Lys	Ile	Ala	Ala	Ser	Phe	Pro	Phe	Ser	Phe	Cys	Val	Met	Asp	Lys	Asp
			500					505					510		
Arg	Glu	Gly	Leu	Ile	Ser	Arg	Asp	Glu	Ile	Thr	Ala	Tyr	Phe	Met	Arg
		515					520					525			
Ala	Ser	Ser	Ile	Tyr	Ser	Lys	Leu	Gly	Leu	Gly	Phe	Pro	His	Asn	Phe
	530					535					540				
Gln	Glu	Thr	Thr	Tyr	Leu	Lys	Pro	Thr	Phe	Cys	Asp	Asn	Cys	Ala	Gly
545					550					555					560
Phe	Leu	Trp	Gly	Val	Ile	Lys	Gln	Gly	Tyr	Arg	Cys	Lys	Asp	Cys	Gly
				565					570					575	
Met	Asn	Cys	His	Lys	Gln	Cys	Lys	Asp	Leu	Val	Val	Phe	Glu	Cys	Lys
			580					585					590		
Lys	Arg	Ser	Lys	Ser	Pro	Ala	Val	Ser	Thr	Glu	Asn	Ile	Ser	Ser	Val
		595					600					605			
Val	Pro	Met	Ser	Thr	Leu	Cys	Pro	Leu	Gly	Thr	Lys	Asp	Leu	Leu	His
	610					615					620				
Ala	Pro	Glu	Glu	Gly	Ser	Phe	Ile	Phe	Gln	Asn	Gly	Glu	Val	Val	Asp
625					630					635					640
His	Ser	Glu	Glu	Ser	Lys	Asp	Arg	Thr	Ile	Met	Leu	Leu	Gly	Val	Ser
				645					650					655	
Ser	Gln	Lys	Ile	Ser	Val	Arg	Leu	Lys	Arg	Thr	Val	Ala	His	Lys	Thr
			660					665					670		
Thr	Gln	Thr	Glu	Ser	Phe	Pro	Trp	Val	Gly	Gly	Glu	Met	Pro	Pro	Gly
		675					680					685			
His	Phe	Val	Leu	Thr	Ser	Pro	Arg	Lys	Ser	Ala	Gln	Gly	Ala	Leu	Tyr
	690					695					700				
Val	His	Ser	Pro	Ala	Ser	Pro	Cys	Pro	Ser	Pro	Ala	Leu	Val	Arg	Lys
705					710					715					720
Arg	Ala	Phe	Val	Lys	Trp	Glu	Asn	Lys	Glu	Ser	Leu	Ile	Lys	Pro	Lys
				725					730					735	
Pro	Glu	Leu	His	Leu	Arg	Leu	Arg	Thr	Tyr	Gln	Glu	Leu	Glu	Gln	Glu
			740					745					750		
Val	Asn	Thr	Leu	Arg	Ala	Asp	Asn	Asp	Ala	Leu	Lys	Ile	Gln	Leu	Lys
		755					760					765			
Tyr	Ala	Gln	Lys	Gln	Ile	Glu	Ser	Leu	Gln	Leu	Gly	Lys	Ser	Asn	His
	770					775					780				
Val	Leu	Ala	Gln	Met	Asp	His	Gly	Asp	Gly	Thr					
785						790					795				

<210> 7
 <211> 5075
 <212> DNA
 <213> Homo sapiens

 <220>
 <221> CDS
 <222> (104)..(2494)
 <223> CalDAG-GEFII

<400> 7																		60
ccccggggggg	agagagccgg	caggcggcgg	cggtggtggc	gggggcgatg	cgccgcgccc													
ggccgcgcta	ggtgagccgg	caccggggagc	gcggggccgcg	gcc	atg	ggc	acc	ctg										115
					Met	Gly	Thr	Leu										
					1													
ggc	aag	gcg	aga	gag	gct	ccg	cgg	aaa	cct	tcc	cat	ggc	tgc	aga	gct			163
Gly	Lys	Ala	Arg	Glu	Ala	Pro	Arg	Lys	Pro	Ser	His	Gly	Cys	Arg	Ala			
5					10					15					20			
gcc	tct	aaa	gca	aga	cta	gag	gca	aag	cca	gcc	aac	agc	ccc	ttc	ccc			211
Ala	Ser	Lys	Ala	Arg	Leu	Glu	Ala	Lys	Pro	Ala	Asn	Ser	Pro	Phe	Pro			
				25					30					35				
tcc	cat	ccc	agc	ttg	gcc	cac	atc	acc	cag	ttc	cga	atg	atg	gtg	tct			259
Ser	His	Pro	Ser	Leu	Ala	His	Ile	Thr	Gln	Phe	Arg	Met	Met	Val	Ser			
				40					45					50				
ctg	gga	cat	tta	gcc	aaa	gga	gcc	agc	ctg	gac	gat	ctc	att	gac	agc			307
Leu	Gly	His	Leu	Ala	Lys	Gly	Ala	Ser	Leu	Asp	Asp	Leu	Ile	Asp	Ser			
				55					60									
tgc	att	caa	tct	ttt	gat	gca	gat	gga	aac	ctg	tgt	cga	agt	aac	caa			355
Cys	Ile	Gln	Ser	Phe	Asp	Ala	Asp	Gly	Asn	Leu	Cys	Arg	Ser	Asn	Gln			
				70					75									
ctg	ttg	caa	gtc	atg	ctg	acc	atg	cac	cga	att	gtc	atc	tcc	tct	gca			403
Leu	Leu	Gln	Val	Met	Leu	Thr	Met	His	Arg	Ile	Val	Ile	Ser	Ser	Ala			
				85					90					95			100	
gaa	ctg	ctc	caa	aaa	gtt	atc	acc	ctc	tat	aag	gat	gct	ttg	gca	aag			451
Glu	Leu	Leu	Gln	Lys	Val	Ile	Thr	Leu	Tyr	Lys	Asp	Ala	Leu	Ala	Lys			
				105					110					115				
aat	tca	cca	gga	ctt	tgc	ctg	aag	atc	tgt	tat	ttt	gta	agg	tat	tgg			499
Asn	Ser	Pro	Gly	Leu	Cys	Leu	Lys	Ile	Cys	Tyr	Phe	Val	Arg	Tyr	Trp			
				120					125					130				
ata	aca	gaa	ttc	tgg	gtc	atg	ttt	aaa	atg	gac	gcc	agc	ttg	aca	gac			547
Ile	Thr	Glu	Phe	Trp	Val	Met	Phe	Lys	Met	Asp	Ala	Ser	Leu	Thr	Asp			
				135					140					145				
act	atg	gag	gag	ttt	cag	gaa	ctg	gtg	aaa	gct	aag	ggg	gag	gag	tta			595
Thr	Met	Glu	Glu	Phe	Gln	Glu	Leu	Val	Lys	Ala	Lys	Gly	Glu	Glu	Leu			
				150					155					160				
cat	tgc	cgc	ctg	att	gac	aca	act	caa	atc	aat	gcc	cgt	gac	tgg	tcc			643
His	Cys	Arg	Leu	Ile	Asp	Thr	Thr	Gln	Ile	Asn	Ala	Arg	Asp	Trp	Ser			
				165					170					175			180	
agg	aaa</																	

- 16 -

aag ttc atc cag gtg gct cag aag ctc cac caa cta cag aac ttc aat	979
Lys Phe Ile Gln Val Ala Gln Lys Leu His Gln Leu Gln Asn Phe Asn	
280 285 290	
aca ctg atg gct gtg ata ggt ggg ctg tgt cac agc tca atc tcg agg	1027
Thr Leu Met Ala Val Ile Gly Gly Leu Cys His Ser Ser Ile Ser Arg	
295 300 305	
ctc aag gag aca agt tcg cat gtc cca cat gaa atc aat aag gtt ctc	1075
Leu Lys Glu Thr Ser Ser His Val Pro His Glu Ile Asn Lys Val Leu	
310 315 320	
ggt gag atg act gag ctg ctg tcc tcc tcc aga aac tac gac aat tac	1123
Gly Glu Met Thr Glu Leu Leu Ser Ser Ser Arg Asn Tyr Asp Asn Tyr	
325 330 335 340	
cgg cga gcc tat gga gag tgc acc gac ttc aag atc ccc att ctg ggt	1171
Arg Arg Ala Tyr Gly Glu Cys Thr Asp Phe Lys Ile Pro Ile Leu Gly	
345 350 355	
gtg cat ctc aag gac ctc atc tcc ctg tat gaa gcc atg cct gac tat	1219
Val His Leu Lys Asp Leu Ile Ser Leu Tyr Glu Ala Met Pro Asp Tyr	
360 365 370	
ctg ggg gac ggg aaa gtg aac gtc cat aag cta ctg gcc cta tac aat	1267
Leu Gly Asp Gly Lys Val Asn Val His Lys Leu Leu Ala Leu Tyr Asn	
375 380 385	
cat atc agt gaa ttg gtc cag ctg caa gag gtg gcc cca ccc ttg gag	1315
His Ile Ser Glu Leu Val Gln Leu Gln Glu Val Ala Pro Pro Leu Glu	
390 395 400	
gct aac aag gac ttg gta cac ttg ctg acg tta tcc ctg gat ctt tac	1363
Ala Asn Lys Asp Leu Val His Leu Leu Thr Leu Ser Leu Asp Leu Tyr	
405 410 415 420	
tac act gag gat gaa atc tat gag ctt tcc tat gcc cgg gaa cca agg	1411
Tyr Thr Glu Asp Glu Ile Tyr Glu Leu Ser Tyr Ala Arg Glu Pro Arg	
425 430 435	
aac cac aga gct cca cca cta aca cct tca aag cca cca gta gta gtg	1459
Asn His Arg Ala Pro Pro Leu Thr Pro Ser Lys Pro Pro Val Val Val	
440 445 450	
gac tgg gct tct gga gtg tct ccc aaa cct gat cca aaa acc att agc	1507
Asp Trp Ala Ser Gly Val Ser Pro Lys Pro Asp Pro Lys Thr Ile Ser	
455 460 465	
aaa cac gtc cag agg atg gtg gat tct gtc ttc aag aac tat gat cac	1555
Lys His Val Gln Arg Met Val Asp Ser Val Phe Lys Asn Tyr Asp His	
470 475 480	
gac cag gat gga tac att tct cag gaa gaa ttt gaa aag att gct gcg	1603
Asp Gln Asp Gly Tyr Ile Ser Gln Glu Glu Phe Glu Lys Ile Ala Ala	
485 490 495 500	
agt ttt cca ttt tcc ttc tgt gtg atg gac aaa gac agg gaa ggc ctc	1651
Ser Phe Pro Phe Ser Phe Cys Val Met Asp Lys Asp Arg Glu Gly Leu	
505 510 515	
atc agc agg gat gag atc aca gcc tac ttc atg aga gcc agc tca atc	1699
Ile Ser Arg Asp Glu Ile Thr Ala Tyr Phe Met Arg Ala Ser Ser Ile	
520 525 530	
tat tcc aag ctg ggc ctg ggc ttt cct cac aac ttc caa gag acc acc	1747
Tyr Ser Lys Leu Gly Leu Gly Phe Pro His Asn Phe Gln Glu Thr Thr	
535 540 545	
tac ctg aag ccc act ttt tgt gac aac tgt gct gga ttt ctc tgg gga	1795
Tyr Leu Lys Pro Thr Phe Cys Asp Asn Cys Ala Gly Phe Leu Trp Gly	
550 555 560	
gtg atc aaa caa gga tat cga tgt aaa gac tgc ggg atg aac tgt cac	1843
Val Ile Lys Gln Gly Tyr Arg Cys Lys Asp Cys Gly Met Asn Cys His	

- 17 -

565	570	575	580	
aaa caa tgc aaa gat ctg gtt gtg ttt gag tgt aag aag cga gcc aag	1891			
Lys Gln Cys Lys Asp 585 Leu Val Val Phe Glu Cys Lys Lys Arg Ala Lys 595				
aac cca gta gct ccc aca gag aac aac act tct gtg ggg cca gtg tcc	1939			
Asn Pro Val Ala 600 Thr Glu Asn Asn Thr Ser Val Gly Pro Val Ser 610				
aac ctt tgc tca ttg gga gcc aaa gat ctg ctc cat gca cct gag gaa	1987			
Asn Leu Cys 615 Ser Leu Gly Ala Lys Asp Leu Leu His Ala Pro Glu Glu 625				
gga cct ttt aca ttc cct aat ggg gag gct gtg gaa cat ggt gag gag	2035			
Gly Pro Phe Thr Phe 630 Pro Asn 635 Gly Glu Ala Val Glu His Gly Glu Glu 640				
agt aag gat cgg acc atc atg ctg atg gga gtg tcc tca cag aag att	2083			
Ser Lys Asp Arg Thr 645 Ile Met Leu Met Gly Val Ser Ser Gln Lys Ile 660				
tct ctt cgg ctg aag agg gct gtt gcc cac aag gcc acc cag act gaa	2131			
Ser Leu Arg Leu Lys 665 Arg Ala Val Ala His Lys Ala Thr Gln Thr Glu 675				
tca cag cct tgg att ggc agt gag ggc cct tca ggt ccc ttt gtg ctg	2179			
Ser Gln Pro Trp 680 Ile Gly Ser Glu Gly Pro Ser Gly Pro Phe Val Leu 690				
tct tcc cca agg aag aca gcc cag gat act cta tat gtg ctt ccc agt	2227			
Ser Ser Pro Arg Lys Thr Ala Gln Asp Thr Leu Tyr Val Leu Pro Ser 705				
ccc acc tct cca tgt cct agc cca gtc ttg gtc aga aag cgg gct ttt	2275			
Pro Thr Ser Pro Cys 710 Pro Ser 715 Pro Val Leu Val Arg Lys Arg Ala Phe 720				
gtc aag tgg gag aat aaa gac tcc ctc ata aaa tca aag gag gag ctc	2323			
Val Lys Trp Glu Asn Lys Asp Ser Leu Ile Lys 735 Ser Lys Glu Glu Leu 740				
cgt cac ctc aga ctg cct acc tac caa gaa ctg gaa cag gaa ata aat	2371			
Arg His Leu Arg Leu 745 Pro Thr Tyr Gln Glu Leu Glu Gln Glu Ile Asn 755				
act ctg aaa gca gat aat gat gcc cta aag atc caa ctg aaa tat gca	2419			
Thr Leu Lys Ala Asp Asn Asp Ala Leu Lys Ile Gln Leu Lys Tyr Ala 770				
cag aag aaa ata gaa tcc ctc cag ctt gaa aaa agc aat cat gtc tta	2467			
Gln Lys Lys Ile Glu Ser Leu 780 Gln Leu Glu Lys Ser Asn His Val Leu 785				
gct caa atg gag cag ggt gac tgt tct tagccccagaa actaagtagc	2514			
Ala Gln Met Glu Gln Gly 790 Asp Cys Ser 795				
acaatctgta gatgagtata gtgatctcat ttccctaaact gtaatgcaca gacctgagga	2574			
actttacact gaccagcttt aaaacagtac tttaaaagga aaagcctgtt actgtttatt	2634			
tacctaaaag attcctaattg tgcagcactg ttttctcttt cagttagttg actcaaaggg	2694			
ggaaaaactaa agaatgcaaa acttttgcta ttcataccac tgatgttcat caaaagtatg	2754			
cgatctaaaa catgactatc atttcttgac aatggggcat ctcggtggcc tgcttggtg	2814			
attcttccctt aaaactaaaa tctctggaaa atggatttgc ttcttaccct gtgttttctg	2874			
caaactgact tactttgttc cagccaaagc ttgctaataa tagaaaaacta cccatattcc	2934			
aaaagtagat ttccctctgta tcccagcata ctttgtgaac ctggtctcctt cttcactacc	2994			
tcagatctaa tcaattagtc catgtaccct ccttcctcac tacagtcata acacatgagc	3054			

- 18 -

atatctacct agaagccaat ttctactgat gtagccacaa ctttttagag gcctattaaa 3114
catgacgtta tccaattgca ggtcaactta tagttgtage cttacaactt acaggcatca 3174
gaaaataagt aatcaaatta ggtacctgga aacatagcta ttaccatctc atattactgt 3234
ctaattaaaa taaacataac gcaaacatgt ttgtccttat atattctaca gtggatagaa 3294
ttaggaattg atggcttaaa aaaaaaagtc tatgaagagt ctgtttaact cttcatgttc 3354
catctttctc ttctgaagta aactattttg aaagtctctt ttttgaaatg aatttgtgct 3414
taactgtctt cactattaat actattttaga aataagctaa ttggatcagt ggcttaaata 3474
atagctgact gtgtgtacat atgtatataa tatgtatata caatatcagg catgcatgtg 3534
gcttggaatt ttgtttcctc cataaaatgt ggaagtgaat taaacaagtt ttagtcattt 3594
atacaaagtc acaaatataa agttcagttt gtcacaagat taaattgctc acaaggtaaa 3654
attgtattgt ttggcaaaat cacaagtaac aatcctgtga gttttctatt atgaaggtta 3714
ataataaatg ggctcattta gttgcctggg cacctattca caaattcatt tgtcagcctc 3774
tttttagttc tcttaaaaaa aaaaaaatca tatgacattt ttcttttttg ggggtactta 3834
gcttccatgc ctataaagtc tggtagcaga ctgacttgaa attcataaac aagttgtcca 3894
attgccaaga atatgttaac aattaaaagt tccaaactaa agccaatagc accaagtctt 3954
cataagaata caaagtatac atacagtatt gcttacctgg aggattcaga tcatttagga 4014
attctctttg atgaaagatc agttcccatt tgagttcctc cttgcactga gtttttagtga 4074
tatagaacta gcttgtagtt agtgtttcat tacattataa agaatagttt tacacacgta 4134
tttaccgttt tccaaattta aactcagaaa tacccaaagc aggctgctt aagcccaacta 4194
cctggcatat aaacttatag taacactttg ttactttctt tttaatagga caagcatgag 4254
ttaggacaaa ctctaaaaat tcatattctt cactattctt gttttccttt gattgatata 4314
gaccaaagat ggtgtactct aattttttta aacagtaatg gaacacaatt tttttcattc 4374
ttcctcctct ccattogaag taaagatccc cagttagttt ttatataaat aatctatagg 4434
gattcaaaaag gtgtcacagt ccacttaatt agtcaaatta gcaatggcta aacagtatca 4494
agtactgcag aattttatcac tgaaatggat aagaggaaat agtttagtca caggttttta 4554
cagtccagca agggccaaag aggtatagta tacaagttaa tagtatttgt gttgagcaac 4614
atggggctag tgggatcaca gaaatctgga aaaaaaaaaa aaaaggcttt ggcttatcaa 4674
gcctagtgtg aattttctgca tctcacagca ctttagtttg gccagggtatt tatctgccaa 4734
aacaaggaca aatcttggtg tattaacagc agggtcactt ctcattttct ttgctgactt 4794
acctttttac tgaccgttgt gaattttctgt ctcaaatgt ataatataga aatgcaagaa 4854
aaaaacaaat gtacagattg taaagttttt tgatacctaa tgtaagtttt ctttgtgtaa 4914
tatttatatg ataaaagaca ttaggatccc tacaaaaaaaa aaaaaaaaaa aaaaaaaaaa 4974
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaggg atcgacaatt 5034
acgactcccc gggggggccc ggtccaattc ccccaaaagg g 5075

<210> 8
<211> 797
<212> PRT
<213> Homo sapiens

- 19 -

<400> 8
Met Gly Thr Leu Gly Lys Ala Arg Glu Ala Pro Arg Lys Pro Ser His
1 5 10 15
Gly Cys Arg Ala Ala Ser Lys Ala Arg Leu Glu Ala Lys Pro Ala Asn
20 25 30
Ser Pro Phe Pro Ser His Pro Ser Leu Ala His Ile Thr Gln Phe Arg
35 40 45
Met Met Val Ser Leu Gly His Leu Ala Lys Gly Ala Ser Leu Asp Asp
50 55 60
Leu Ile Asp Ser Cys Ile Gln Ser Phe Asp Ala Asp Gly Asn Leu Cys
65 70 75 80
Arg Ser Asn Gln Leu Leu Gln Val Met Leu Thr Met His Arg Ile Val
85 90 95
Ile Ser Ser Ala Glu Leu Leu Gln Lys Val Ile Thr Leu Tyr Lys Asp
100 105 110
Ala Leu Ala Lys Asn Ser Pro Gly Leu Cys Leu Lys Ile Cys Tyr Phe
115 120 125
Val Arg Tyr Trp Ile Thr Glu Phe Trp Val Met Phe Lys Met Asp Ala
130 135 140
Ser Leu Thr Asp Thr Met Glu Glu Phe Gln Glu Leu Val Lys Ala Lys
145 150 155 160
Gly Glu Glu Leu His Cys Arg Leu Ile Asp Thr Thr Gln Ile Asn Ala
165 170 175
Arg Asp Trp Ser Arg Lys Leu Thr Gln Arg Ile Lys Ser Asn Thr Ser
180 185 190
Lys Lys Arg Lys Val Ser Leu Leu Phe Asp His Leu Glu Pro Glu Glu
195 200 205
Leu Ser Glu His Leu Thr Tyr Leu Glu Phe Lys Ser Phe Arg Arg Ile
210 215 220
Ser Phe Ser Asp Tyr Gln Asn Tyr Leu Val Asn Ser Cys Val Lys Glu
225 230 235 240
Asn Pro Thr Met Glu Arg Ser Ile Ala Leu Cys Asn Gly Ile Ser Gln
245 250 255
Trp Val Gln Leu Met Val Leu Ser Arg Pro Thr Pro Gln Leu Arg Ala
260 265 270
Glu Val Phe Ile Lys Phe Ile Gln Val Ala Gln Lys Leu His Gln Leu
275 280 285
Gln Asn Phe Asn Thr Leu Met Ala Val Ile Gly Gly Leu Cys His Ser
290 295 300
Ser Ile Ser Arg Leu Lys Glu Thr Ser Ser His Val Pro His Glu Ile
305 310 315 320
Asn Lys Val Leu Gly Glu Met Thr Glu Leu Leu Ser Ser Ser Arg Asn
325 330 335
Tyr Asp Asn Tyr Arg Arg Ala Tyr Gly Glu Cys Thr Asp Phe Lys Ile
340 345 350
Pro Ile Leu Gly Val His Leu Lys Asp Leu Ile Ser Leu Tyr Glu Ala
355 360 365
Met Pro Asp Tyr Leu Gly Asp Gly Lys Val Asn Val His Lys Leu Leu
370 375 380
Ala Leu Tyr Asn His Ile Ser Glu Leu Val Gln Leu Gln Glu Val Ala
385 390 395 400

- 20 -

Pro	Pro	Leu	Glu	Ala	Asn	Lys	Asp	Leu	Val	His	Leu	Leu	Thr	Leu	Ser
				405					410						415
Leu	Asp	Leu	Tyr	Tyr	Thr	Glu	Asp	Glu	Ile	Tyr	Glu	Leu	Ser	Tyr	Ala
			420					425					430		
Arg	Glu	Pro	Arg	Asn	His	Arg	Ala	Pro	Pro	Leu	Thr	Pro	Ser	Lys	Pro
		435					440					445			
Pro	Val	Val	Val	Asp	Trp	Ala	Ser	Gly	Val	Ser	Pro	Lys	Pro	Asp	Pro
	450					455					460				
Lys	Thr	Ile	Ser	Lys	His	Val	Gln	Arg	Met	Val	Asp	Ser	Val	Phe	Lys
465					470					475					480
Asn	Tyr	Asp	His	Asp	Gln	Asp	Gly	Tyr	Ile	Ser	Gln	Glu	Glu	Phe	Glu
				485					490					495	
Lys	Ile	Ala	Ala	Ser	Phe	Pro	Phe	Ser	Phe	Cys	Val	Met	Asp	Lys	Asp
			500					505					510		
Arg	Glu	Gly	Leu	Ile	Ser	Arg	Asp	Glu	Ile	Thr	Ala	Tyr	Phe	Met	Arg
		515					520					525			
Ala	Ser	Ser	Ile	Tyr	Ser	Lys	Leu	Gly	Leu	Gly	Phe	Pro	His	Asn	Phe
	530					535					540				
Gln	Glu	Thr	Thr	Tyr	Leu	Lys	Pro	Thr	Phe	Cys	Asp	Asn	Cys	Ala	Gly
545					550					555					560
Phe	Leu	Trp	Gly	Val	Ile	Lys	Gln	Gly	Tyr	Arg	Cys	Lys	Asp	Cys	Gly
				565					570					575	
Met	Asn	Cys	His	Lys	Gln	Cys	Lys	Asp	Leu	Val	Val	Phe	Glu	Cys	Lys
			580					585					590		
Lys	Arg	Ala	Lys	Asn	Pro	Val	Ala	Pro	Thr	Glu	Asn	Asn	Thr	Ser	Val
		595					600					605			
Gly	Pro	Val	Ser	Asn	Leu	Cys	Ser	Leu	Gly	Ala	Lys	Asp	Leu	Leu	His
		610				615					620				
Ala	Pro	Glu	Glu	Gly	Pro	Phe	Thr	Phe	Pro	Asn	Gly	Glu	Ala	Val	Glu
625					630					635					640
His	Gly	Glu	Glu	Ser	Lys	Asp	Arg	Thr	Ile	Met	Leu	Met	Gly	Val	Ser
				645					650					655	
Ser	Gln	Lys	Ile	Ser	Leu	Arg	Leu	Lys	Arg	Ala	Val	Ala	His	Lys	Ala
			660					665					670		
Thr	Gln	Thr	Glu	Ser	Gln	Pro	Trp	Ile	Gly	Ser	Glu	Gly	Pro	Ser	Gly
		675					680					685			
Pro	Phe	Val	Leu	Ser	Ser	Pro	Arg	Lys	Thr	Ala	Gln	Asp	Thr	Leu	Tyr
	690					695					700				
Val	Leu	Pro	Ser	Pro	Thr	Ser	Pro	Cys	Pro	Ser	Pro	Val	Leu	Val	Arg
705					710					715					720
Lys	Arg	Ala	Phe	Val	Lys	Trp	Glu	Asn	Lys	Asp	Ser	Leu	Ile	Lys	Ser
				725					730					735	
Lys	Glu	Glu	Leu	Arg	His	Leu	Arg	Leu	Pro	Thr	Tyr	Gln	Glu	Leu	Glu
			740					745					750		
Gln	Glu	Ile	Asn	Thr											

```
<210> 9
<211> 3373
<212> DNA
<213> Rattus norvegicus
```

```
<220>
<221> CDS
<222> (197) .. (2848)
<223> cAMP-GEFI
```

SUBSTITUTE SHEET (RULE 26)

- 22 -

ttt Phe	gaa Glu	cca Pro	cac His	agc Ser 225	aag Lys	gca Ala	gga Gly	act Thr 230	gtg Val	ttg Leu	ttc Phe	agc Ser	cag Gln	ggg Gly 235	gac Asp	904
aag Lys	ggt Gly	acc Thr	tca Ser 240	tgg Trp	tac Tyr	att Ile	atc Ile	tgg Trp 245	aag Lys	gga Gly	tct Ser	gtc Val	aat Asn 250	gtg Val	gtg Val	952
acc Thr	cgt Arg	ggc Gly 255	aag Lys	ggg Gly	ctg Leu	gtg Val	acc Thr 260	acg Thr	ttg Leu	cac His	gag Glu	gga Gly 265	gat Asp	gac Asp	ttt Phe	1000
gga Gly 270	cag Gln	ctg Leu	gct Ala	ctg Leu	gtg Val	aac Asn 275	gac Asp	gca Ala	cct Pro	cga Arg	gca Ala 280	gcc Ala	acc Thr	atc Ile	atc Ile	1048
ctt Leu 285	cga Arg	gaa Glu	aat Asn	aac Asn	tgt Cys 290	cac His	ttt Phe	ctg Leu	cgt Arg	gtg Val 295	gac Asp	aag Lys	cag Gln	gac Asp	ttc Phe 300	1096
aac Asn	cgc Arg	atc Ile	atc Ile	aag Lys 305	gat Asp	gtg Val	gaa Glu	gca Ala	aaa Lys 310	acc Thr	atg Met	aga Arg	ctg Leu	gaa Glu 315	gaa Glu	1144
cac His	ggc Gly	aaa Lys	gtg Val 320	gtg Val	tta Leu	gtt Val	ttg Leu 325	gag Glu	aga Arg	acc Thr	tct Ser	cag Gln	ggt Gly 330	gct Ala	ggc Gly	1192
cct Pro	tcc Ser	cgc Arg 335	cct Pro	ccg Pro	acc Thr	cca Pro	ggc Gly 340	agg Arg	aac Asn	cga Arg	tat Tyr	acg Thr 345	gta Val	atg Met	tct Ser	1240
ggc Gly 350	acc Thr	cca Pro	gag Glu	aaa Lys	atc Ile	cta Leu 355	gaa Glu	ctc Leu	ctg Leu	ttg Leu	gag Glu 360	gct Ala	atg Met	aga Arg	ccg Pro	1288
gat Asp 365	tcc Ser	agt Ser	gct Ala	cat His	gac Asp 370	cca Pro	aca Thr	gag Glu	aca Thr	ttc Phe 375	ctc Leu	agt Ser	gac Asp	ttc Phe	ctg Leu 380	1336
ctg Leu	acg Thr	cac His	agt Ser	gtc Val 385	ttc Phe	atg Met	ccc Pro	tgc Cys	aca Thr 390	cag Gln	ctc Leu	ttt Phe	gcc Ala	gcc Ala 395	ctc Leu	1384
ctg Leu	cac His	cac His	ttc Phe 400	cac His	gtg Val	gag Glu	cca Pro	tca Ser 405	gag Glu	cct Pro	gcc Ala	ggg Gly	ggc Gly 410	agc Ser	gag Glu	1432
cag Gln	gaa Glu	cgc Arg 415	agc Ser	acc Thr	tac Tyr	atc Ile	tgc Cys 420	aac Asn	aag Lys	agg Arg	cag Gln	cag Gln	att Ile	ctg Leu	cgt Arg	1480
ctg Leu	gtc Val 430	agc Ser	cgg Arg	tgg Trp	gtg Val	gcc Ala 435	ctc Leu	tac Tyr	agc Ser	ccc Pro	atg Met 440	ctc Leu	cgc Arg	tca Ser	gat Asp	1528
ccc Pro 445	gtg Val	gcc Ala	acc Thr	agc Ser	ttc Phe 450	ctc Leu	cag Gln	aaa Lys	ctc Leu	tca Ser 455	gac Asp	ctg Leu	gtg Val	agc Ser	aga Arg 460	1576
gat Asp	acc Thr	cga Arg	ctt Leu	agc Ser 465	aac Asn	ttg Leu	ctg Leu	agg Arg	gaa Glu 470	cag Gln	tat Tyr	ccg Pro	gag Glu	aga Arg 475	cgg Arg	1624
cga Arg	cac His	cac His	agg Arg 480	ttg Leu	gag Glu	aat Asn	ggc Gly 485	tgt Cys	ggg Gly	aat Asn	gta Val	tct Ser	cct Pro 490	cag Gln	acc Thr	1672
aag Lys	gcc Ala	cgg Arg 495	aat Asn	gca Ala	cct Pro	gtt Val	tgg Trp 500	ttt Phe	cct Pro	aac Asn	cat His	gag Glu 505	gaa Glu	ccc Pro	ctc Leu	1720
cca Pro	agc Ser 510	agc Ser	gct Ala	ggg Gly	gcc Ala	atc Ile 515	cga Arg	gtc Val	ggg Gly	gac Asp	aaa Lys 520	gtc Val	ccc Pro	tat Tyr	gat Asp	1768

- 23 -

atc Ile 525	tgc Cys	aga Arg	ccc Pro	gac Asp	cac His 530	tcg Ser	gtg Val	ctg Leu	acc Thr 535	ctg Leu	cac His	ctg Leu	ccg Pro	gtg Val	aca Thr 540	1816
gcc Ala	tcc Ser	gtg Val	agg Arg	gaa Glu 545	gtg Val	atg Met	gca Ala	gct Ala	ttg Leu 550	gcc Ala	cat His	gag Glu	gac Asp	cac His 555	tgg Trp	1864
acc Thr	aag Lys	ggg Gly	cag Gln 560	gtg Val	ctg Leu	gta Val	aag Lys	gtc Val 565	aat Asn	tct Ser	gcc Ala	ggt Gly	gat Asp 570	gtc Val	gtt Val	1912
ggc Gly	ttg Leu	cag Gln 575	cca Pro	gat Asp	gcc Ala	cgc Arg	ggt Gly 580	gtg Val	gcc Ala	aca Thr	tcc Ser	ctg Leu 585	ggg Gly	ctc Leu	aat Asn	1960
gag Glu 590	cgg Arg	atc Ile	ttt Phe	gtt Val	gtc Val	gac Asp 595	cca Pro	cag Gln	gaa Glu	gtg Val	cac His 600	gag Glu	ctg Leu	acc Thr	cca Pro	2008
cac His 605	cct Pro	gag Glu	cag Gln	ctg Leu	ggg Gly 610	ccc Pro	act Thr	ctg Leu	ggt Gly	tct Ser 615	tct Ser	gag Glu	atg Met	ctg Leu	gac Asp 620	2056
cta Leu	gtg Val	agt Ser	gcc Ala	aag Lys 625	gac Asp	ctg Leu	gca Ala	ggc Gly	cag Gln 630	ctc Leu	aca Thr	gag Glu	cat His	gac Asp 635	tgg Trp	2104
aac Asn	ctc Leu	ttc Phe	aac Asn 640	agg Arg	atc Ile	cac His	cag Gln	gtg Val 645	gag Glu	ctg Leu	atc Ile	cac His	tat Tyr 650	gta Val	ctg Leu	2152
ggc Gly	ccc Pro	cag Gln 655	cac His	ctg Leu	cgg Arg	gac Asp	gtc Val 660	acc Thr	act Thr	gca Ala	aac Asn	ctg Leu 665	gag Glu	cgc Arg	ttc Phe	2200
atg Met 670	cga Arg	cgc Arg	ttc Phe	aac Asn	gag Glu	ctg Leu 675	cag Gln	tac Tyr	tgg Trp	gtg Val	gcc Ala 680	acc Thr	gaa Glu	ctc Leu	tgt Cys	2248
ctc Leu 685	tgc Cys	ccg Pro	gtt Val	cct Pro	ggc Gly 690	ccc Pro	cgg Arg	gct Ala	cag Gln	cta Leu 695	ctc Leu	cgg Arg	aag Lys	ttc Phe	atc Ile 700	2296
aag Lys	ctg Leu	gca Ala	gcc Ala	cac His 705	ctc Leu	aag Lys	gag Glu	cag Gln	aag Lys 710	aat Asn	ctc Leu	aac Asn	tct Ser	ttc Phe 715	ttt Phe	2344
gcg Ala	gtc Val	atg Met	ttt Phe 720	ggc Gly	ctc Leu	agc Ser	aac Asn	tcg Ser 725	gcc Ala	atc Ile	agc Ser	cgc Arg	ctg Leu 730	gcc Ala	cac His	2392
acc Thr	tgg Trp	gag Glu	cgt Arg	ctg Leu	ccc Pro	cat His	aaa Lys 740	gta Val	cgg Arg	aag Lys	ctg Leu	tac Tyr 745	tcg Ser	gcc Ala	ctg Leu	2440
gaa Glu 750	agg Arg	ttg Leu	ctg Leu	gac Asp	cct Pro	tcc Ser 755	tgg Trp	aac Asn	cac His	cga Arg	gtg Val 760	tac Tyr	cga Arg	ttg Leu	gct Ala	2488
ctc Leu 765	acc Thr	aag Lys	ctc Leu	tct Ser	cct Pro 770	cct Pro	gtc Val	atc Ile	cct Pro 775	ttc Phe	atg Met	ccc Pro	ctg Leu	cta Leu	ctc Leu 780	2536
aaa Lys	gac Asp	atg Met	acc Thr	ttc Phe 785	att Ile	cat His	gaa Glu	ggg Gly	aac Asn 790	cac His	aca Thr	ctg Leu	gta Val	gaa Glu 795	aac Asn	2584
ctc Leu	atc Ile	aac Asn	ttt Phe 800	gag Glu	aag Lys	atg Met	cga Arg	atg Met 805	gcc Ala	aga Arg	gcc Ala	gtg Val 810	cgg Arg	atg Met		2632
ctc Leu	cac His	cac His	tgc Cys	cga Arg	agc Ser	cac His	agc Ser	acc Thr	gcg Ala	cct Pro	cta Leu	tca Ser	cca Pro	ctc Leu	aga Arg	2680

- 24 -

815	820	825	
agc cgg gtt tcc cac atc	cac gag gac agc cag	gca tca aga atc tcc	2728
Ser Arg Val Ser His Ile	His Glu Asp Ser Gln	Ala Ser Arg Ile Ser	
830	835	840	
aca tgt tcc gag cag tcc	ctg agc acc cgg agt	cca gcc agc acc tgg	2776
Thr Cys Ser Glu Gln Ser	Leu Ser Thr Arg Ser	Pro Ala Ser Thr Trp	
845	850	855	860
gct tat gtc cag cag	ctg aag gtc att gac aac	cag cgg gaa ctg tcc	2824
Ala Tyr Val Gln Gln	Leu Lys Val Ile Asp Asn	Gln Arg Glu Leu Ser	
	865	870	875
cgc ctc tcc cgg gaa	ctg gaa cca	tgaggaagga ctggctggag	caggcacttc 2878
Arg Leu Ser Arg Glu	Leu Glu Pro		
880			
tctcagagaa agccagagcc	tgtgcaacca agaggtccag	aggccagcca cagctgggca	2938
gggctctcca cagagcggac	tcaaggccct ggagtgggca	gtgtcgaggc agctgtcctc	2998
tgtgatgact gtcagctgtg	aagatctttg atgttcacgg	ccaaggaaaa ggggccattg	3058
agggcccaga ggggtaggag	agctgggagg tgcaggactc	tggttcagta gagagccctc	3118
ccaccgggct ttctgcatgt	ctgtatgtct gtacatgcag	ctgtgtgtcc tggatgccag	3178
gccctgtgct tgtattcaca	ggagccagcg agctcacatc	tgcattgtgt gtgtgtgtgt	3238
gtgtgtgtgt gtgtgtgtgt	gtgtgtgtga gcaactgggtg	gtgccttgcc tgggagaggg	3298
tggggaggttc tgctattctc	accacacatc tgagataaac	agctgggtgt gggcaaaaaa	3358
aaaaaaaaa aaaaa			3373

<210> 10
 <211> 884
 <212> PRT
 <213> Rattus norvegicus

<400> 10
 Met Val Leu Lys Arg Met His Arg Pro Arg Cys Cys Ser Tyr Gln Leu
 1 5 10 15
 Val Phe Glu His Arg Arg Pro Ser Cys Ile Gln Gly Leu Arg Trp Thr
 20 25 30
 Pro Leu Thr Asn Ser Glu Gly Ser Leu Asp Phe Arg Val Ser Leu Glu
 35 40 45
 Gln Ala Thr Thr Glu His Val His Lys Ala Gly Lys Leu Leu Tyr Arg
 50 55 60
 His Leu Leu Ala Thr Tyr Pro Thr Leu Ile Arg Asp Arg Lys Tyr His
 65 70 75 80
 Leu Arg Leu His Arg Gln Cys Cys Ser Gly Arg Glu Leu Val Asp Gly
 85 90 95
 Ile Leu Ala Leu Gly Leu Gly Val His Ser Arg Ser Gln Ala Val Gly
 100 105 110
 Ile Cys Gln Val Leu Leu Asp Glu Gly Ala Leu Cys His Val Lys His
 115 120 125
 Asp Trp Thr Phe Gln Asp Arg Asp Ala Gln Phe Tyr Arg Phe Pro Gly
 130 135 140
 Pro Glu Pro Gln Pro Ala Gly Thr His Asp Val Glu Glu Glu Leu Val
 145 150 155 160
 Glu Ala Met Ala Leu Leu Ser Gln Arg Gly Pro Asp Ala Leu Leu Thr
 165 170 175

- 25 -

Val Ala Leu Arg Lys Ser Pro Gly Gln Arg Thr Asp Glu Glu Leu Asp
 180 185 190
 Leu Ile Phe Glu Glu Leu Val His Ile Lys Ala Val Ala His Leu Ser
 195 200 205
 Asn Ser Val Lys Arg Glu Leu Ala Ala Val Leu Leu Phe Glu Pro His
 210 215 220
 Ser Lys Ala Gly Thr Val Leu Phe Ser Gln Gly Asp Lys Gly Thr Ser
 225 230 235 240
 Trp Tyr Ile Ile Trp Lys Gly Ser Val Asn Val Val Thr Arg Gly Lys
 245 250 255
 Gly Leu Val Thr Thr Leu His Glu Gly Asp Asp Phe Gly Gln Leu Ala
 260 265 270
 Leu Val Asn Asp Ala Pro Arg Ala Ala Thr Ile Ile Leu Arg Glu Asn
 275 280 285
 Asn Cys His Phe Leu Arg Val Asp Lys Gln Asp Phe Asn Arg Ile Ile
 290 295 300
 Lys Asp Val Glu Ala Lys Thr Met Arg Leu Glu Glu His Gly Lys Val
 305 310 315 320
 Val Leu Val Leu Glu Arg Thr Ser Gln Gly Ala Gly Pro Ser Arg Pro
 325 330 335
 Pro Thr Pro Gly Arg Asn Arg Tyr Thr Val Met Ser Gly Thr Pro Glu
 340 345 350
 Lys Ile Leu Glu Leu Leu Leu Glu Ala Met Arg Pro Asp Ser Ser Ala
 355 360 365
 His Asp Pro Thr Glu Thr Phe Leu Ser Asp Phe Leu Leu Thr His Ser
 370 375 380
 Val Phe Met Pro Cys Thr Gln Leu Phe Ala Ala Leu Leu His His Phe
 385 390 395 400
 His Val Glu Pro Ser Glu Pro Ala Gly Gly Ser Glu Gln Glu Arg Ser
 405 410 415
 Thr Tyr Ile Cys Asn Lys Arg Gln Gln Ile Leu Arg Leu Val Ser Arg
 420 425 430
 Trp Val Ala Leu Tyr Ser Pro Met Leu Arg Ser Asp Pro Val Ala Thr
 435 440 445
 Ser Phe Leu Gln Lys Leu Ser Asp Leu Val Ser Arg Asp Thr Arg Leu
 450 455 460
 Ser Asn Leu Leu Arg Glu Gln Tyr Pro Glu Arg Arg Arg His His Arg
 465 470 475 480
 Leu Glu Asn Gly Cys Gly Asn Val Ser Pro Gln Thr Lys Ala Arg Asn
 485 490 495
 Ala Pro Val Trp Phe Pro Asn His Glu Glu Pro Leu Pro Ser Ser Ala
 500 505 510
 Gly Ala Ile Arg Val Gly Asp Lys Val Pro Tyr Asp Ile Cys Arg Pro
 515 520 525
 Asp His Ser Val Leu Thr Leu His Leu Pro Val Thr Ala Ser Val Arg
 530 535 540
 Glu Val Met Ala Ala Leu Ala His Glu Asp His Trp Thr Lys Gly Gln
 545 550 555 560
 Val Leu Val Lys Val Asn Ser Ala Gly Asp Val Val Gly Leu Gln Pro
 565 570 575

- 26 -

Asp Ala Arg Gly Val Ala Thr Ser Leu Gly Leu Asn Glu Arg Ile Phe
 580 585 590
 Val Val Asp Pro Gln Glu Val His Glu Leu Thr Pro His Pro Glu Gln
 595 600 605
 Leu Gly Pro Thr Leu Gly Ser Ser Glu Met Leu Asp Leu Val Ser Ala
 610 615 620
 Lys Asp Leu Ala Gly Gln Leu Thr Glu His Asp Trp Asn Leu Phe Asn
 625 630 635 640
 Arg Ile His Gln Val Glu Leu Ile His Tyr Val Leu Gly Pro Gln His
 645 650 655
 Leu Arg Asp Val Thr Thr Ala Asn Leu Glu Arg Phe Met Arg Arg Phe
 660 665 670
 Asn Glu Leu Gln Tyr Trp Val Ala Thr Glu Leu Cys Leu Cys Pro Val
 675 680 685
 Pro Gly Pro Arg Ala Gln Leu Leu Arg Lys Phe Ile Lys Leu Ala Ala
 690 695 700
 His Leu Lys Glu Gln Lys Asn Leu Asn Ser Phe Phe Ala Val Met Phe
 705 710 715 720
 Gly Leu Ser Asn Ser Ala Ile Ser Arg Leu Ala His Thr Trp Glu Arg
 725 730 735
 Leu Pro His Lys Val Arg Lys Leu Tyr Ser Ala Leu Glu Arg Leu Leu
 740 745 750
 Asp Pro Ser Trp Asn His Arg Val Tyr Arg Leu Ala Leu Thr Lys Leu
 755 760 765
 Ser Pro Pro Val Ile Pro Phe Met Pro Leu Leu Leu Lys Asp Met Thr
 770 775 780
 Phe Ile His Glu Gly Asn His Thr Leu Val Glu Asn Leu Ile Asn Phe
 785 790 795 800
 Glu Lys Met Arg Met Met Ala Arg Ala Val Arg Met Leu His His Cys
 805 810 815
 Arg Ser His Ser Thr Ala Pro Leu Ser Pro Leu Arg Ser Arg Val Ser
 820 825 830
 His Ile His Glu Asp Ser Gln Ala Ser Arg Ile Ser Thr Cys Ser Glu
 835 840 845
 Gln Ser Leu Ser Thr Arg Ser Pro Ala Ser Thr Trp Ala Tyr Val Gln
 850 855 860
 Gln Leu Lys Val Ile Asp Asn Gln Arg Glu Leu Ser Arg Leu Ser Arg
 865 870 875 880
 Glu Leu Glu Pro

<210> 11
 <211> 3394
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (216)..(2858)
 <223> CAMP-GEFI

<400> 11
 ggatccccctt atcaaagctg atgggggtcg ctgggggaccc cccagccttt tgctagagcc 60

tg	cac	gg	ctg	tggg	ag	cttg	aaa	aga	aa	aca	tga	aggt	ggg	ctgg	cc	caggt	gag	ag	ctg	ct	120
gg	caggt	ggg	cct	gg	ctg	tg	gag	ga	tag	cc	cag	ct	ctg	ggg	ag	cacc	cg	gtg	ggg	ag	180
tc	ctg	ac	gt	gg	gc	gg	gag	ggg	ac	act	ac	atg	gtg	ttg	aga	agg	atg				233
												Met	Val	Leu	Arg	Arg	Met				
												1				5					
cac	cg	ccc	cga	agc	tgc	tcc	tac	cag	ctg	ctg	ctg	gag	cac	cag	cat						281
His	Arg	Pro	Arg	Ser	Cys	Ser	Tyr	Gln	Leu	Leu	Leu	Glu	His	Gln	His						
			10					15					20								
ccg	agc	tgc	atc	cag	ggg	ctg	cgc	tgg	aca	cca	ctc	acc	aac	agc	gag						329
Pro	Ser	Cys	Ile	Gln	Gly	Leu	Arg	Trp	Thr	Pro	Leu	Thr	Asn	Ser	Glu						
		25					30					35									
gag	tcc	ctg	gat	ttc	agc	gag	agc	ctg	gag	cag	gcc	tcc	aca	gag	cgg						377
Glu	Ser	Leu	Asp	Phe	Ser	Glu	Ser	Leu	Glu	Gln	Ala	Ser	Thr	Glu	Arg						
	40					45					50										
gtg	ctc	agg	gct	ggg	agg	cag	ctg	cat	cag	cat	cta	ctg	gcc	acc	tgc						425
Val	Leu	Arg	Ala	Gly	Arg	Gln	Leu	His	Gln	His	Leu	Leu	Ala	Thr	Cys						
	55				60					65					70						
cca	aac	ctc	atc	cga	gac	cgg	aag	tac	cac	ctt	agg	ctc	tat	cgg	cag						473
Pro	Asn	Leu	Ile	Arg	Asp	Arg	Lys	Tyr	His	Leu	Arg	Leu	Tyr	Arg	Gln						
				75					80					85							
tgc	tgc	tct	ggc	cgg	gag	ctg	gtg	gat	ggg	atc	ttg	gcc	ctg	gga	ctt						521
Cys	Cys	Ser	Gly	Arg	Glu	Leu	Val	Asp	Gly	Ile	Leu	Ala	Leu	Gly	Leu						
			90					95				100									
ggg	gtc	cat	tcc	cgg	agc	caa	gtt	gtg	gga	atc	tgc	cag	gtg	ctg	ctg						569
Gly	Val	His	Ser	Arg	Ser	Gln	Val	Val	Gly	Ile	Cys	Gln	Val	Leu	Leu						
		105				110						115									
gat	gaa	gg	gcc	ctc	tgc	cat	gtg	aaa	cac	gac	tgg	gcc	ttc	cag	gac						617
Asp	Glu	Gly	Ala	Leu	Cys	His	Val	Lys	His	Asp	Trp	Ala	Phe	Gln	Asp						
	120					125					130										
cga	gat	gcc	caa	ttc	tac	cgg	ttc	ccc	ggg	ccc	gag	ccc	gag	ccc	gtg						665
Arg	Asp	Ala	Gln	Phe	Tyr	Arg	Phe	Pro	Gly	Pro	Glu	Pro	Glu	Pro	Val						
					140				145						150						
gga	act	cat	gag	atg	gag	gag	gag	ttg	gcc	gaa	gct	gtg	gcc	ctg	ctc						713
Gly	Thr	His	Glu																		

- 28 -

cgg	gca	gcc	acc	atc	atc	ctg	cga	gaa	tac	aac	tgt	cat	ttc	ctg	cgt	1097
Arg	Ala	Ala	Thr	Ile	Ile	Leu	Arg	Glu	Tyr	Asn	Cys	His	Phe	Leu	Arg	
	280					285					290					
gtg	gac	aag	cag	gac	ttc	aac	cgt	atc	atc	aag	gat	gtg	gag	gca	aag	1145
Val	Asp	Lys	Gln	Asp	Phe	Asn	Arg	Ile	Ile	Lys	Asp	Val	Glu	Ala	Lys	
295					300					305					310	
acc	atg	cgg	ctg	gaa	gaa	cat	ggc	aaa	gtg	gtg	ctg	gtg	ctg	gag	aga	1193
Thr	Met	Arg	Leu	Glu	Glu	His	Gly	Lys	Val	Val	Leu	Val	Leu	Glu	Arg	
				315					320					325		
gcc	tct	cag	ggc	gcc	ggc	cct	tcc	cga	ccc	cca	acc	cca	ggc	agg	aac	1241
Ala	Ser	Gln	Gly	Ala	Gly	Pro	Ser	Arg	Pro	Pro	Thr	Pro	Gly	Arg	Asn	
			330					335					340			
cgg	tat	aca	gtg	atg	tct	ggc	act	cca	gat	aag	atc	cta	gag	ctt	ctg	1289
Arg	Tyr	Thr	Val	Met	Ser	Gly	Thr	Pro	Asp	Lys	Ile	Leu	Glu	Leu	Leu	
	345					350						355				
ttg	gag	gcc	atg	gga	cta	gat	tcc	agt	gct	cat	gac	cca	aaa	gaa	aca	1337
Leu	Glu	Ala	Met	Gly	Leu	Asp	Ser	Ser	Ala	His	Asp	Pro	Lys	Glu	Thr	
360						365					370					
ttc	ctc	agc	gac	ttc	ctc	ctg	acc	cac	agg	gtc	ttc	atg	ccc	agc	gcc	1385
Phe	Leu	Ser	Asp	Phe	Leu	Leu	Thr	His	Arg	Val	Phe	Met	Pro	Ser	Ala	
375					380					385					390	
caa	ctc	tgc	gct	gcc	ctt	ctg	cac	cac	ttc	cat	gtg	gag	cct	gcg	ggt	1433
Gln	Leu	Cys	Ala	Ala	Leu	Leu	His	His	Phe	His	Val	Glu	Pro	Ala	Gly	
				395					400					405		
ggc	agc	gag	cag	gag	cgc	agc	acc	tac	gtc	tgc	aac	aag	agg	cag	cag	1481
Gly	Ser	Glu	Gln	Glu	Arg	Ser	Thr	Tyr	Val	Cys	Asn	Lys	Arg	Gln	Gln	
			410					415					420			
atc	ttg	cgg	ctg	gtc	agc	cag	tgg	gtg	gcc	ctg	tat	ggc	tcc	atg	ctc	1529
Ile	Leu	Arg	Leu	Val	Ser	Gln	Trp	Val	Ala	Leu	Tyr	Gly	Ser	Met	Leu	
	425						430					435				
cac	act	gac	cct	gtg	gcc	acc	agc	ttc	ctc	cac	aaa	ctc	tca	gac	ctg	1577
His	Thr	Asp	Pro	Val	Ala	Thr	Ser	Phe	Leu	His	Lys	Leu	Ser	Asp	Leu	
	440					445					450					
gtg	ggc	agg	gac	acc	cga	ctc	agc	aac	ctg	ctg	agg	gag	cag	tgg	cca	1625
Val	Gly	Arg	Asp	Thr	Arg	Leu	Ser	Asn	Leu	Leu	Arg	Glu	Gln	Trp	Pro	
455					460					465					470	
gag	agg	cgg	cga	tgc	cac	agg	ttg	gag	aat	ggc	tgt	ggg	aat	gca	tct	1673
Glu	Arg	Arg	Arg	Cys	His	Arg	Leu	Glu	Asn	Gly	Cys	Gly	Asn	Ala	Ser	
				475					480					485		
cct	cag	atg	aag	gcc	cgg	aac	ttg	cct	gtt	tgg	ctc	ccc	aac	cag	gac	1721
Pro	Gln	Met	Lys	Ala	Arg	Asn	Leu	Pro	Val	Trp	Leu	Pro	Asn	Gln	Asp	
			490					495					500			
gag	ccc	ctt	cct	ggc	agc	agc	tgt	gcc	atc	caa	gtt	ggg	gat	aaa	gtc	1769
Glu	Pro	Leu	Pro	Gly	Ser	Ser	Cys	Ala	Ile	Gln	Val	Gly	Asp	Lys	Val	
	505						510					515				
ccc	tat	gac	atc	tgc	cgg	cca	gac	cac	tca	gtg	ttg	acc	ctg	cag	ctg	1817
Pro	Tyr	Asp	Ile	Cys	Arg	Pro	Asp	His	Ser	Val	Leu	Thr	Leu	Gln	Leu	
	520					525					530					
cct	gtg	aca	gcc	tcc	gtg	aga	gag	gtg	atg	gca	gcg	ttg	gcc	cag	gag	1865
Pro	Val	Thr	Ala	Ser	Val	Arg	Glu	Val	Met	Ala	Ala	Leu	Ala	Gln	Glu	
535					540					545					550	
gat	ggc	tgg	acc	aag	ggg	cag	gtg	ctg	gtg	aag	gtc	aat	tct	gca	ggt	1913
Asp	Gly	Trp	Thr	Lys	Gly	Gln	Val	Leu	Val	Lys	Val	Asn	Ser	Ala	Gly	
				555					560					565		
gat	gcc	att	ggc	ctg	cag	cca	gat	gcc	cgt	ggt	gtg	gcc	aca	tct	ctg	1961
Asp	Ala	Ile	Gly	Leu	Gln	Pro	Asp	Ala	Arg	Gly	Val	Ala	Thr	Ser	Leu	

- 29 -

570					575					580					
ggg ctc aat gag cgt ctc ttt gtt gtc aac cca cag gaa gtg cat gag	2009														
Gly Leu Asn Glu Arg Leu Phe Val Val Asn Pro Gln Glu Val His Glu															
585 590 595															
ctg atc cca cac cct gac cag ctg ggg ccc act gtg ggc tct gct gag	2057														
Leu Ile Pro His Pro Asp Gln Leu Gly Pro Thr Val Gly Ser Ala Glu															
600 605 610															
ggg ctg gac ctg gtg agt gcc aag gac ctg gca ggc cag ctg acg gac	2105														
Gly Leu Asp Leu Val Ser Ala Lys Asp Leu Ala Gly Gln Leu Thr Asp															
615 620 625 630															
cac gac tgg agc ctc ttc aac agt atc cac cag gtg gag ctg atc cac	2153														
His Asp Trp Ser Leu Phe Asn Ser Ile His Gln Val Glu Leu Ile His															
635 640 645															
tat gtg ctg ggc ccc cag cat ctg cgg gat gtc acc acc gcc aac ctg	2201														
Tyr Val Leu Gly Pro Gln His Leu Arg Asp Val Thr Thr Ala Asn Leu															
650 655 660															
gag cgc ttc atg cgc cgc ttc aat gag ctg cag tac tgg gtg gcc acc	2249														
Glu Arg Phe Met Arg Arg Phe Asn Glu Leu Gln Tyr Trp Val Ala Thr															
665 670 675															
gag ctg tgt ctc tgc ccc gtg ccc ggc ccc cgg gcc cag ctg ctc aaa	2297														
Glu Leu Cys Leu Cys Pro Val Pro Gly Pro Arg Ala Gln Leu Leu Lys															
680 685 690															
aag ttc att aag ctg gcg gcc cac ctc aag gag cag aag aat gtc aat	2345														
Lys Phe Ile Lys Leu Ala Ala His Leu Lys Glu Gln Lys Asn Val Asn															
695 700 705 710															
tcc ttc ttt gcc gtc atg ttt ggc ctc agc aac tcg ccc atc agc cgc	2393														
Ser Phe Phe Ala Val Met Phe Gly Leu Ser Asn Ser Pro Ile Ser Arg															
715 720 725															
cta gcc cac acc tgg gag cgg ctg cct cac aaa gtc cgg aag ctg tac	2441														
Leu Ala His Thr Trp Glu Arg Leu Pro His Lys Val Arg Lys Leu Tyr															
730 735 740															
tcc gcc ctc gag agg ctg ctg gat ccc tca tgg aac cac cgg gta tac	2489														
Ser Ala Leu Glu Arg Leu Leu Asp Pro Ser Trp Asn His Arg Val Tyr															
745 750 755															
cga ctg gcc ctc gcc aag ctc tcc cct cct gtc atc ccc ttc atg ccc	2537														
Arg Leu Ala Leu Ala Lys Leu Ser Pro Pro Val Ile Pro Phe Met Pro															
760 765 770															
ctt ctt ctc aaa gac atg acc ttc att cat gag gga aac cac aca cta	2585														
Leu Leu Leu Lys Asp Met Thr Phe Ile His Glu Gly Asn His Thr Leu															
775 780 785 790															
gtg gag aat ctc atc aac ttt gag aag atg aga atg atg gcc aga gcc	2633														
Val Glu Asn Leu Ile Asn Phe Glu Lys Met Arg Met Met Ala Arg Ala															
795 800 805															
gcg cgg atg ctg cac cac tgc cga agc cac aac cct gtg cct ctc tca	2681														
Ala Arg Met Leu His His Cys Arg Ser His Asn Pro Val Pro Leu Ser															
810 815 820															
cca ctc aga agc cga gtt tcc cac ctc cac gag gac agc cag gtg gcg	2729														
Pro Leu Arg Ser Arg Val Ser His Leu His Glu Asp Ser Gln Val Ala															
825 830 835															
agg att tcc aca tgc tcg gag cag tcc ctg agc acc cgg agt cca gcc	2777														
Arg Ile Ser Thr Cys Ser Glu Gln Ser Leu Ser Thr Arg Ser Pro Ala															
840 845 850															
agc acc tgg gct tat gtc cag cag ctg aag gtc att gac aac cag cgg	2825														
Ser Thr Trp Ala Tyr Val Gln Gln Leu Lys Val Ile Asp Asn Gln Arg															
855 860 865 870															
gaa ctc tcc cgc ctg tcc cga gag ctg gag cca tgaggagggg ctgggaactgg	2878														

- 30 -

Glu Leu Ser Arg Leu Ser Arg Glu Leu Glu Pro
875 880

agctggagca ggcacttgca gccgggaaag ccagggtgtg ccgggccaaag atactcacag 2938
gctggccaca gctgggcaag gctctccgtg gactggactc gactccctgg agcaggcagt 2998
gtggaggcag ccatcccctg tgatgactgg cagctaagga ggacctcgga gtggaccaa 3058
gccaggaata acgaatgacc aaggggcaag gaaggaggga cagagaggcc ccaggagtgg 3118
gtggagagtg gactgcgctg ggacgttgtg tgcaatagag aggtctccac accagatgtc 3178
ttccagattc tgtgcctctg gctttgttgt ccagccaggc ctgcagttta ttttcacagt 3238
ggacagagag agagagagag gctgcatgtg tgtaccgtgt gtggcaaggg cagggccttg 3298
gcctgggggca gggggcccctg ctttctttcc acagctttct tccaacagca ggcagtgggg 3358
ctgcgggcct gaaaaaaaaa aaaaaaaaaa aaaaaa 3394

<210> 12
<211> 881
<212> PRT
<213> Homo sapiens

<400> 12
Met Val Leu Arg Arg Met His Arg Pro Arg Ser Cys Ser Tyr Gln Leu
1 5 10 15
Leu Leu Glu His Gln His Pro Ser Cys Ile Gln Gly Leu Arg Trp Thr
20 25 30
Pro Leu Thr Asn Ser Glu Glu Ser Leu Asp Phe Ser Glu Ser Leu Glu
35 40 45
Gln Ala Ser Thr Glu Arg Val Leu Arg Ala Gly Arg Gln Leu His Gln
50 55 60
His Leu Leu Ala Thr Cys Pro Asn Leu Ile Arg Asp Arg Lys Tyr His
65 70 75 80
Leu Arg Leu Tyr Arg Gln Cys Cys Ser Gly Arg Glu Leu Val Asp Gly
85 90 95
Ile Leu Ala Leu Gly Leu Gly Val His Ser Arg Ser Gln Val Val Gly
100 105 110
Ile Cys Gln Val Leu Leu Asp Glu Gly Ala Leu Cys His Val Lys His
115 120 125
Asp Trp Ala Phe Gln Asp Arg Asp Ala Gln Phe Tyr Arg Phe Pro Gly
130 135 140
Pro Glu Pro Glu Pro Val Gly Thr His Glu Met Glu Glu Glu Leu Ala
145 150 155 160
Glu Ala Val Ala Leu Leu Ser Gln Arg Gly Pro Asp Ala Leu Leu Thr
165 170 175
Val Ala Leu Arg Lys Pro Pro Gly Gln Arg Thr Asp Glu Glu Leu Asp
180 185 190
Leu Ile Phe Glu Glu Leu Leu His Ile Lys Ala Val Ala His Leu Ser
195 200 205
Asn Ser Val Lys Arg Glu Leu Ala Ala Val Leu Leu Phe Glu Pro His
210 215 220
Ser Lys Ala Gly Thr Val Leu Phe Ser Gln Gly Asp Lys Gly Thr Ser
225 230 235 240
Trp Tyr Ile Ile Trp Lys Gly Ser Val Asn Val Val Thr His Gly Lys
245 250 255

- 31 -

Gly Leu Val Thr Thr Leu His Glu Gly Asp Asp Phe Gly Gln Leu Ala
 260 265 270
 Leu Val Asn Asp Ala Pro Arg Ala Ala Thr Ile Ile Leu Arg Glu Tyr
 275 280 285
 Asn Cys His Phe Leu Arg Val Asp Lys Gln Asp Phe Asn Arg Ile Ile
 290 295 300
 Lys Asp Val Glu Ala Lys Thr Met Arg Leu Glu Glu His Gly Lys Val
 305 310 315 320
 Val Leu Val Leu Glu Arg Ala Ser Gln Gly Ala Gly Pro Ser Arg Pro
 325 330 335
 Pro Thr Pro Gly Arg Asn Arg Tyr Thr Val Met Ser Gly Thr Pro Asp
 340 345 350
 Lys Ile Leu Glu Leu Leu Leu Glu Ala Met Gly Leu Asp Ser Ser Ala
 355 360 365
 His Asp Pro Lys Glu Thr Phe Leu Ser Asp Phe Leu Leu Thr His Arg
 370 375 380
 Val Phe Met Pro Ser Ala Gln Leu Cys Ala Ala Leu Leu His His Phe
 385 390 395 400
 His Val Glu Pro Ala Gly Gly Ser Glu Gln Glu Arg Ser Thr Tyr Val
 405 410 415
 Cys Asn Lys Arg Gln Gln Ile Leu Arg Leu Val Ser Gln Trp Val Ala
 420 425 430
 Leu Tyr Gly Ser Met Leu His Thr Asp Pro Val Ala Thr Ser Phe Leu
 435 440 445
 His Lys Leu Ser Asp Leu Val Gly Arg Asp Thr Arg Leu Ser Asn Leu
 450 455 460
 Leu Arg Glu Gln Trp Pro Glu Arg Arg Arg Cys His Arg Leu Glu Asn
 465 470 475 480
 Gly Cys Gly Asn Ala Ser Pro Gln Met Lys Ala Arg Asn Leu Pro Val
 485 490 495
 Trp Leu Pro Asn Gln Asp Glu Pro Leu Pro Gly Ser Ser Cys Ala Ile
 500 505 510
 Gln Val Gly Asp Lys Val Pro Tyr Asp Ile Cys Arg Pro Asp His Ser
 515 520 525
 Val Leu Thr Leu Gln Leu Pro Val Thr Ala Ser Val Arg Glu Val Met
 530 535 540
 Ala Ala Leu Ala Gln Glu Asp Gly Trp Thr Lys Gly Gln Val Leu Val
 545 550 555 560
 Lys Val Asn Ser Ala Gly Asp Ala Ile Gly Leu Gln Pro Asp Ala Arg
 565 570 575
 Gly Val Ala Thr Ser Leu Gly Leu Asn Glu Arg Leu Phe Val Val Asn
 580 585 590
 Pro Gln Glu Val His Glu Leu Ile Pro His Pro Asp Gln Leu Gly Pro
 595 600 605
 Thr Val Gly Ser Ala Glu Gly Leu Asp Leu Val Ser Ala Lys Asp Leu
 610 615 620
 Ala Gly Gln Leu Thr Asp His Asp Trp Ser Leu Phe Asn Ser Ile His
 625 630 635 640
 Gln Val Glu Leu Ile His Tyr Val Leu Gly Pro Gln His Leu Arg Asp
 645 650 655

- 32 -

Val Thr Thr Ala Asn Leu Glu Arg Phe Met Arg Arg Phe Asn Glu Leu
660 665 670
Gln Tyr Trp Val Ala Thr Glu Leu Cys Leu Cys Pro Val Pro Gly Pro
675 680 685
Arg Ala Gln Leu Leu Lys Lys Phe Ile Lys Leu Ala Ala His Leu Lys
690 695 700
Glu Gln Lys Asn Val Asn Ser Phe Phe Ala Val Met Phe Gly Leu Ser
705 710 715 720
Asn Ser Pro Ile Ser Arg Leu Ala His Thr Trp Glu Arg Leu Pro His
725 730 735
Lys Val Arg Lys Leu Tyr Ser Ala Leu Glu Arg Leu Leu Asp Pro Ser
740 745 750
Trp Asn His Arg Val Tyr Arg Leu Ala Leu Ala Lys Leu Ser Pro Pro
755 760 765
Val Ile Pro Phe Met Pro Leu Leu Leu Lys Asp Met Thr Phe Ile His
770 775 780
Glu Gly Asn His Thr Leu Val Glu Asn Leu Ile Asn Phe Glu Lys Met
785 790 795 800
Arg Met Met Ala Arg Ala Ala Arg Met Leu His His Cys Arg Ser His
805 810 815
Asn Pro Val Pro Leu Ser Pro Leu Arg Ser Arg Val Ser His Leu His
820 825 830
Glu Asp Ser Gln Val Ala Arg Ile Ser Thr Cys Ser Glu Gln Ser Leu
835 840 845
Ser Thr Arg Ser Pro Ala Ser Thr Trp Ala Tyr Val Gln Gln Leu Lys
850 855 860
Val Ile Asp Asn Gln Arg Glu Leu Ser Arg Leu Ser Arg Glu Leu Glu
865 870 875 880
Pro

<210> 13
<211> 4109
<212> DNA
<213> Homo sapiens

<220>
<221> CDS
<222> (216)..(1883)
<223> Alternate splicing of cAMP--GEFI

<400> 13
ggatccccctt atcaaagctg atgggggtcg ctggggaccc cccagccttt tgctagagcc 60
tgcacggctg tgggagcttg aaaagaaaca tgaaggtggg ctggccaggt gagagctgct 120
ggcaggtggg cctggctgtg gaggatagcc cagctctggg agcaccgctg gtgggagccc 180
tccctgacgt ggtgccggag gggacactac tcaac atg gtg ttg aga agg atg 233
Met Val Leu Arg Arg Met
1 5
cac cgg ccc cga agc tgc tcc tac cag ctg ctg ctg gag cac cag cat 281
His Arg Pro Arg Ser Cys Ser Tyr Gln Leu Leu Leu Glu His Gln His
10 15 20
ccg agc tgc atc cag ggg ctg cgc tgg aca cca ctc acc aac agc gag 329
Pro Ser Cys Ile Gln Gly Leu Arg Trp Thr Pro Leu Thr Asn Ser Glu
25 30 35

- 33 -

gag tcc ctg gat ttc agc gag agc ctg gag cag gcc tcc aca gag cgg Glu Ser Leu Asp Phe Ser Glu Ser Leu Glu Gln Ala Ser Thr Glu Arg	377
40 45 50	
gtg ctc agg gct ggg agg cag ctg cat cag cat cta ctg gcc acc tgc Val Leu Arg Ala Gly Gln Leu His Gln His Leu Leu Ala Thr Cys	425
55 60 65 70	
cca aac ctc atc cga gac cgg aag tac cac ctt agg ctc tat cgg cag Pro Asn Leu Ile Arg Asp Arg Lys Tyr His Leu Arg Leu Tyr Arg Gln	473
75 80 85	
tgc tgc tct ggc cgg gag ctg gtg gat ggg atc ttg gcc ctg gga ctt Cys Cys Ser Gly Arg Glu Leu Val Asp Gly Ile Leu Ala Leu Gly Leu	521
90 95 100	
ggg gtc cat tcc cgg agc caa gtt gtg gga atc tgc cag gtg ctg ctg Gly Val His Ser Arg Ser Gln Val Val Gly Ile Cys Gln Val Leu Leu	569
105 110 115	
gat gaa ggt gcc ctc tgc cat gtg aaa cac gac tgg gcc ttc cag gac Asp Glu Gly Ala Leu Cys His Val Lys His Asp Trp Ala Phe Gln Asp	617
120 125 130	
cga gat gcc caa ttc tac cgg ttc ccc ggg ccc gag ccc gag ccc gtg Arg Asp Ala Gln Phe Tyr Arg Phe Pro Gly Pro Glu Pro Glu Pro Val	665
135 140 145 150	
gga act cat gag atg gag gag gag ttg gcc gaa gct gtg gcc ctg ctc Gly Thr His Glu Met Glu Glu Glu Leu Ala Glu Ala Val Ala Leu Leu	713
155 160 165	
tcc cag cgg ggg cct gac gcc ctg ctc act gtg gca ctt cga aag ccc Ser Gln Arg Gly Pro Asp Ala Leu Leu Thr Val Ala Leu Arg Lys Pro	761
170 175 180	
cca ggt cag cgc acg gat gaa gag ctg gac ctc atc ttt gag gag ctg Pro Gly Gln Arg Thr Asp Glu Glu Leu Asp Leu Ile Phe Glu Glu Leu	809
185 190 195	
ctg cac atc aag gct gtg gcc cac ctc tcc aac tgc gtg aag cga gaa Leu His Ile Lys Ala Val Ala His Leu Ser Asn Ser Val Lys Arg Glu	857
200 205 210	
tta gcg gct gtt ctg ctc ttt gaa cca cac agc aag gca ggg acc gtg Leu Ala Ala Val Leu Phe Glu Pro His Ser Lys Ala Gly Thr Val	905
215 220 225 230	
ttg ttc agc cag ggg gac aag ggc act tgc tgg tac att atc tgg aag Leu Phe Ser Gln Gly Asp Lys Gly Thr Ser Trp Tyr Ile Ile Trp Lys	953
235 240 245	
gga tct gtc aac gtg gtg acc cat ggc aag ggg ctg gtg acc acc ctg Gly Ser Val Asn Val Val Thr His Gly Lys Gly Leu Val Thr Thr Leu	1001
250 255 260	
cat gag gga gat gat ttt gga cag ctg gct ctg gtg aat gat gca ccc His Glu Gly Asp Asp Phe Gly Gln Leu Ala Leu Val Asn Asp Ala Pro	1049
265 270 275	
cgg gca gcc acc atc atc ctg cga gaa tac aac tgt cat ttc ctg cgt Arg Ala Ala Thr Ile Ile Leu Arg Glu Tyr Asn Cys His Phe Leu Arg	1097
280 285 290	
gtg gac aag cag gac ttc aac cgt atc atc aag gat gtg gag gca aag Val Asp Lys Gln Asp Phe Asn Arg Ile Ile Lys Asp Val Glu Ala Lys	1145
295 300 305 310	
acc atg cgg ctg gaa gaa cat ggc aaa gtg gtg ctg gtg ctg gag aga Thr Met Arg Leu Glu Glu His Gly Lys Val Val Leu Val Leu Glu Arg	1193
315 320 325	
gcc tct cag ggc gcc ggc cct tcc cga ccc cca acc cca ggc agg aac Ala Ser Gln Gly Ala Gly Pro Ser Arg Pro Pro Thr Pro Gly Arg Asn	1241

- 34 -

330	335	340	
cggtat	acagtgt	tctggc	actcca
ArgTyr	ThrVal	MetSer	GlyThr
345		350	
cggtat	acagtgt	tctggc	actcca
ArgTyr	ThrVal	MetSer	GlyThr
345		350	
ttggag	gccatg	gga	cta
LeuGlu	AlaMet	GlyLeu	AspThr
360		365	
ttggag	gccatg	gga	cta
LeuGlu	AlaMet	GlyLeu	AspThr
360		365	
ttcttc	agcgac	ttcttc	ctc
PheLeu	SerAsp	PheLeu	Thr
375		380	
ttcttc	agcgac	ttcttc	ctc
PheLeu	SerAsp	PheLeu	Thr
375		380	
caactc	tgcgct	gcc	ctt
GlnLeu	CysAla	AlaLeu	His
		395	
caactc	tgcgct	gcc	ctt
GlnLeu	CysAla	AlaLeu	His
		395	
ggcagc	gagcag	gagcgc	agc
GlySer	GluGln	GluArg	Ser
	410		
ggcagc	gagcag	gagcgc	agc
GlySer	GluGln	GluArg	Ser
	410		
atcttg	cggtct	gtcagc	cagtgg
IleLeu	ArgLeu	ValSer	GlnTrp
	425		430
atcttg	cggtct	gtcagc	cagtgg
IleLeu	ArgLeu	ValSer	GlnTrp
	425		430
cacact	gac	cctgtg	gcc
HisThr	AspPro	ValAla	Thr
	440		445
cacact	gac	cctgtg	gcc
HisThr	AspPro	ValAla	Thr
	440		445
gtgggc	agggac	acc	cga
ValGly	ArgAsp	ThrArg	Leu
455		460	
gtgggc	agggac	acc	cga
ValGly	ArgAsp	ThrArg	Leu
455		460	
gagagg	cggtgc	cacagg	ttggag
GluArg	ArgArg	CysHis	ArgLeu
	475		480
gagagg	cggtgc	cacagg	ttggag
GluArg	ArgArg	CysHis	ArgLeu
	475		480
cctcag	atgaag	gtgtct	gcc
ProGln	MetLys	ValSer	AlaTrp
	490		495
cctcag	atgaag	gtgtct	gcc
ProGln	MetLys	ValSer	AlaTrp
	490		495
cctgga	ctgcag	gca	cct
ProGly	LeuGln	AlaPro	Pro
	505		510
cctgga	ctgcag	gca	cct
ProGly	LeuGln	AlaPro	Pro
	505		510
ggcggt	gggaag	ctctcc	cac
GlyArg	GlyLys	LeuSer	His
	520		525
ggcggt	gggaag	ctctcc	cac
GlyArg	GlyLys	LeuSer	His
	520		525
gggtgt	cacggg	gcc	ctt
GlyVal	HisGly	AlaAla	Leu
535		540	
gggtgt	cacggg	gcc	ctt
GlyVal	HisGly	AlaAla	Leu
535		540	
cca	gga	ggc	gca
ProGly	GlyAla	GluAla	
		555	
cca	gga	ggc	gca
ProGly	GlyAla	GluAla	
		555	
gactcaatgg gccttttatt ccttttggga ggtaattcat gcccacaggt agagcctggg 1973			
agatgaggaa tgggtggctgg agttggcccc tgaggccac ggggctcctg gtggccagt 2033			
ctgtgggagc tcagaggaag gcgaggcccc tccctggaaa gtcagagggg gcccagggc 2093			
attctcagct ggggcttaaa gggaagcaaa gaggggatga gaagatgtga ctgcagccag 2153			
gatttgggtg aggaatgggt ggagaaagaa ctggaaaggt agattaggtt ggtgggtggg 2213			
agggctgcac tgggaataaa atccaggggc ctcaccatcc agcaggetcc ccatgggtccc 2273			
acctactgt gtgtcccagc cctgctgc tggagactct tactggttc ttctcctccg 2333			
caggcgaagc ccgttctccc ctcagggcct ttgcacttat cccttagcct ggaagggctc 2393			
ttctctggtc ctgcctggca ggatccttct cgttctcctc tcagccaaat gctgggttcc 2453			

- 35 -

cagacaggcc ccctcccgtt tccctatccg gagcaccctc ccctttctcc caccacgctc 2513
 tgccttggtg tcctctgtga tttttttcaa agcactctcg ccatgtggag ttgttttctt 2573
 tgcctacttg ctttctgtcc tttggtatct ctcccatccc actggagagt tgacttcctc 2633
 cctgcggggc cccacctccg cttctccttg ctcacatgct gcctggctct tggctgacgc 2693
 tcagaaactg atgtgggctc accctcgcag gctgggcact gcgctagggt ctgatgctgc 2753
 acctcattca accctggcct gtggtggctg cagtcaggca aggggcagag cagttcaaca 2813
 acccgtgatg tgcagtgaca cgggacactt tccctgaatg tgcactggag tcggctcttc 2873
 caactgtagg tccgctgcct taaaccagga aggggaagag ccaactccag tgcagattca 2933
 gggaaagtgt cccgtgtaag cgactactgg agttgagagc tgcagggggc aagggtggag 2993
 gaagaagagg attcctggtg ggcaaacca ggaggaaagg gaaccagga ggaggatccc 3053
 tggaggagcg agagtaagag taggaccca gagcaggcag gaggtgggca gggccccctg 3113
 ggcaaggaca ggggtctgat ctttgtccta agagcagtgc aaaccattaa aggggtttga 3173
 gccaaagttg gcaaagtctg atttgtgttt tgaataggtc agctaaagag cctgggcttt 3233
 ctccagcaga caatggacac cactgaagtc gagtgcggga gcttcacagg gctctggtca 3293
 ggcagattgg aggaggaggg ctggaggcac ctgggaggga ggaggaggag catagtggg 3353
 ccattggagt agtaaggagt gaggcctcgt gggcagggtg aactggagtt agcctgggca 3413
 ggggaggagg gggcacagaa gaccacttc aaagaagaat cttcaaagca agatgacaag 3473
 ctacggaagt gtggaggagc ctggagctgg ggagaatggc tggggaacag agtggctctg 3533
 gaggcaggat gcaaggatcc gatgggtata tggagtgtga ggtaatggtt cattcatgtg 3593
 gaaggatgca gggggttttt gagaccaggg tttggaagag agttcagcac tgctggtagt 3653
 tttgggaatc acccatgtgc aggcgacaca tgaggcagta aggaactctg caggggtccc 3713
 tgagatttgg aaatgtaggg aagagcaatg gattgaggtc cgaacctgga ggatctgcta 3773
 tacgcagagc tgggaggagg gacagagtca gtaccagagt cggaaaaaag caggggtggga 3833
 aggggaacct gagtccaggag acttgccctg caggcctgc cctgccagca gaggcctgac 3893
 agtggtttcc atgaactgca tccctgctgt gggctgggac agggccactg acacagtatc 3953
 ggagcacaga aggggaaagg agcaggaggg attccaactc tgccagttag cagctgtgtg 4013
 gctttgggca tgttacttaa cctctctgag cctcatttat ttcattcata aaatggaaat 4073
 aaaaataata cttttgtcaa aaaaaaaaaa aaaaaa 4109

<210> 14
 <211> 556
 <212> PRT
 <213> Homo sapiens

<400> 14
 Met Val Leu Arg Arg Met His Arg Pro Arg Ser Cys Ser Tyr Gln Leu
 1 5 10 15
 Leu Leu Glu His Gln His Pro Ser Cys Ile Gln Gly Leu Arg Trp Thr
 20 25 30
 Pro Leu Thr Asn Ser Glu Glu Ser Leu Asp Phe Ser Glu Ser Leu Glu
 35 40 45
 Gln Ala Ser Thr Glu Arg Val Leu Arg Ala Gly Arg Gln Leu His Gln
 50 55 60

- 36 -

His 65 Leu Leu Ala Thr Cys 70 Pro Asn Leu Ile Arg 75 Asp Arg Lys Tyr His 80
 Leu Arg Leu Tyr Arg 85 Gln Cys Cys Ser Gly 90 Arg Glu Leu Val Asp 95 Gly
 Ile Leu Ala Leu Gly 100 Leu Gly Val His 105 Ser Arg Ser Gln Val Val Gly 110
 Ile Cys Gln Val Leu Leu Asp 115 Glu Gly Ala Leu Cys His 125 Val Lys His
 Asp Trp 130 Ala Phe Gln Asp 135 Arg Asp Ala Gln Phe Tyr 140 Arg Phe Pro Gly
 Pro Glu 145 Pro Glu Pro Val 150 Gly Thr His Glu Met 155 Glu Glu Glu Leu Ala 160
 Glu Ala Val Ala Leu 165 Leu Ser Gln Arg Gly 170 Pro Asp Ala Leu Leu Thr 175
 Val Ala Leu Arg 180 Lys Pro Pro Gly Gln 185 Arg Thr Asp Glu Glu 190 Leu Asp
 Leu Ile Phe 195 Glu Glu Leu Leu His 200 Ile Lys Ala Val Ala His 205 Leu Ser
 Asn Ser 210 Val Lys Arg Glu Leu 215 Ala Ala Val Leu Leu Phe Glu Pro His 220
 Ser 225 Lys Ala Gly Thr Val 230 Leu Phe Ser Gln Gly 235 Asp Lys Gly Thr Ser 240
 Trp Tyr Ile Ile Trp 245 Lys Gly Ser Val Asn 250 Val Val Thr His Gly 255 Lys
 Gly Leu Val Thr 260 Thr Leu His Glu Gly 265 Asp Asp Phe Gly Gln 270 Leu Ala
 Leu Val Asn 275 Asp Ala Pro Arg Ala 280 Ala Thr Ile Ile Leu Arg Glu Tyr 285
 Asn Cys 290 His Phe Leu Arg Val 295 Asp Lys Gln Asp Phe Asn Arg Ile Ile 300
 Lys 305 Asp Val Glu Ala Lys 310 Thr Met Arg Leu Glu Glu His Gly Lys Val 320
 Val Leu Val Leu Glu 325 Arg Ala Ser Gln Gly 330 Ala Gly Pro Ser Arg Pro 335
 Pro Thr Pro Gly 340 Arg Asn Arg Tyr Thr 345 Val Met Ser Gly Thr Pro Asp 350
 Lys Ile Leu Glu Leu Leu Leu Glu 360 Ala Met Gly Leu Asp 365 Ser Ser Ala
 His Asp 370 Pro Lys Glu Thr Phe 375 Leu Ser Asp Phe Leu Leu Thr His Arg 380
 Val Phe Met Pro Ser Ala 390 Gln Leu Cys Ala Ala 395 Leu Leu His His Phe 400
 His Val Glu Pro Ala 405 Gly Gly Ser Glu Gln 410 Glu Arg Ser Thr Tyr Val 415
 Cys Asn Lys Arg 420 Gln Gln Ile Leu Arg 425 Leu Val Ser Gln Trp Val Ala 430
 Leu Tyr Gly 435 Ser Met Leu His Thr 440 Asp Pro Val Ala Thr Ser Phe Leu 445
 Gln Lys 450 Leu Ser Asp Leu Val 455 Gly Arg Asp Thr Arg 460 Leu Ser Asn Leu

- 37 -

Leu Arg Glu Gln Trp Pro Glu Arg Arg Arg Cys His Arg Leu Glu Asn
 465 470 475 480
 Gly Cys Gly Asn Ala Ser Pro Gln Met Lys Val Ser Ala Trp Pro Gln
 485 490 495
 Phe Leu Ser Ser Ala Pro Pro Gly Leu Gln Ala Pro Pro Ser Pro Pro
 500 505 510
 Asp Pro Glu Gly Leu Cys Gly Arg Gly Lys Leu Ser Ser His Arg His
 515 520 525
 Thr Leu Gly Ser Leu Ile Gly Val His Gly Ala Leu Ala Ala Cys Gly
 530 535 540
 Ala Leu Gly Gln Ala Val Pro Gly Gly Ala Glu Ala
 545 550 555

<210> 15
 <211> 1966
 <212> DNA
 <213> Rattus norvegicus

<220>
 <221> CDS
 <222> (3)..(875)
 <223> cAMP-GEFII

<400> 15
 aa ggt gtg ctc aaa cct aat gat gtt tca gta ttt acg acg ctc acc 47
 Gly Val Leu Lys Pro Asn Asp Val Ser Val Phe Thr Thr Leu Thr 15
 1 5 10
 att aat gga cgc ctg ttt gcc tgc ccg cga gag caa ttc gac tca ctg 95
 Ile Asn Gly Arg Leu Phe Ala Cys Pro Arg Glu Gln Phe Asp Ser Leu 20 25 30
 act ccc ttg cca gaa cag gag ggc ccg acc act ggg aca gtg ggg acg 143
 Thr Pro Leu Pro Glu Gln Glu Gly Pro Thr Thr Gly Thr Val Gly Thr 35 40 45
 ttt gaa ctg atg agc tcg aaa gac ttg gcg tac cag atg aca acg tat 191
 Phe Glu Leu Met Ser Ser Lys Asp Leu Ala Tyr Gln Met Thr Thr Tyr 50 55 60
 gac tgg gaa ctc ttc aac tgt gtg ctc gag ctg gag cta atc tac cac 239
 Asp Trp Glu Leu Phe Asn Cys Val Leu Glu Leu Glu Leu Ile Tyr His 65 70 75
 aca ttt gga agg cat aat ttt aaa aag acc aca gca aac ttg gat ttg 287
 Thr Phe Gly Arg His Asn Phe Lys Lys Thr Thr Ala Asn Leu Asp Leu 80 85 90 95
 ttc ctg agg aga ttt aat gaa att cag ttt tgg gtt gtc act gag atc 335
 Phe Leu Arg Arg Phe Asn Glu Ile Gln Phe Trp Val Val Thr Thr Glu Ile 100 105 110
 tgc ctt tgt tcc cag ctc agc aag cgt gtt cag ctt ttg aaa aaa tgt 383
 Cys Leu Cys Ser Gln Leu Ser Lys Arg Val Gln Leu Leu Lys Lys Cys 115 120 125
 atc aag ata gcg gct cac tgc aag gag tac aaa aac ttg aat tcc ttc 431
 Ile Lys Ile Ala Ala His Cys Lys Glu Tyr Lys Asn Leu Asn Ser Phe 130 135 140
 ttc ggc atc gtc atg ggg ctc agt aac gtg gct gag agc cgc ctg gca 479
 Phe Gly Ile Val Met Gly Leu Ser Asn Val Ala Glu Ser Arg Leu Ala 145 150 155
 tta aca tgg gag aaa ctg ccg agc aag ttt aag aag ttc tat gcg gag 527
 Leu Thr Trp Glu Lys Leu Pro Ser Lys Phe Lys Lys Phe Tyr Ala Glu 160 165 170 175

[illegible]

ccacaatctt	tcaaaagatg	ctgtgtatgc	tactactgac	tgtgttgcta	ctagagaatt	975
ccccagaatg	agcaagagac	acctcctgag	agccccctcg	ggccacatcc	tgctttccga	1035
ccacacagga	gaaggatctg	tcttgtctaa	cgcggaacatg	ccgtagctta	ggaaccatca	1095
gctgtagtca	tcttcttcac	ggttggcaca	ccaccgcagg	ctcacgtgaa	ggcataacct	1155
ggcgaggcta	caccaggccc	ctgacatccc	ttcccaggct	gttgacagcat	gagactgtcc	1215
cgtggatagg	tttgacttgg	aatcgctgca	atgatataat	tgaatgattt	gtttacttag	1275
caccttattt	ggggtctggg	ttctggggag	ggtgttgacc	ataaaagtcc	aaattatcca	1335
tcatgttctt	ccatcgctcat	aatcttacct	ctgaaggaat	ggaacctcat	cacaacacta	1395
tgaaacatta	tgactgttca	gtctgtaatt	tcggaatgat	ctatagaata	atatgtttac	1455
attgtaactt	ttaaaaactt	acaaatcagg	attacacaca	tgagaattcc	actaagaaac	1515
accaagggtt	ttaatatcgc	cagcgttaag	atagaaagta	acatttcaga	agagcacaaat	1575
atacacaaaa	cattttttca	aattgaaata	ttttcctggg	cattaaaaaa	cctttccact	1635
acaaatttat	tgttactgat	gaaaaaaaaa	gcatattttc	tggacttaaa	tgttattaca	1695
aaaatcttaa	ttttcagcaa	ttgttttgca	ctttcagata	gattgtaaat	aggttatgca	1755
gtcaatggta	tagaattatt	tatttgtctac	ataatagaca	ttgtgccaaa	taattccttt	1815
ttattttatt	tattcagtat	gaaattttgg	agtacatttt	ttctgttttc	ttaattagac	1875
tacatttaat	gtataggaat	tgtatgtaca	tatctcttct	gtaaataaca	gccagtatct	1935
tcattaaata	tacttgacaa	gaaaaaaaaa	a			1966

```
<210> 16
<211> 291
<212> PRT
<213> Rattus norvegicus
<400> 16
```

- 39 -

Gly Val Leu Lys Pro Asn Asp Val Ser Val Phe Thr Thr Leu Thr Ile
 1 5 10 15
 Asn Gly Arg Leu Phe Ala Cys Pro Arg Glu Gln Phe Asp Ser Leu Thr
 20 25 30
 Pro Leu Pro Glu Gln Glu Gly Pro Thr Thr Gly Thr Val Gly Thr Phe
 35 40 45
 Glu Leu Met Ser Ser Lys Asp Leu Ala Tyr Gln Met Thr Thr Tyr Asp
 50 55 60
 Trp Glu Leu Phe Asn Cys Val Leu Glu Leu Glu Leu Ile Tyr His Thr
 65 70 75 80
 Phe Gly Arg His Asn Phe Lys Lys Thr Thr Ala Asn Leu Asp Leu Phe
 85 90 95
 Leu Arg Arg Phe Asn Glu Ile Gln Phe Trp Val Val Thr Glu Ile Cys
 100 105 110
 Leu Cys Ser Gln Leu Ser Lys Arg Val Gln Leu Leu Lys Lys Cys Ile
 115 120 125
 Lys Ile Ala Ala His Cys Lys Glu Tyr Lys Asn Leu Asn Ser Phe Phe
 130 135 140
 Gly Ile Val Met Gly Leu Ser Asn Val Ala Glu Ser Arg Leu Ala Leu
 145 150 155 160
 Thr Trp Glu Lys Leu Pro Ser Lys Phe Lys Lys Phe Tyr Ala Glu Phe
 165 170 175
 Glu Ser Leu Met Asp Pro Ser Arg Asn His Lys Ala Tyr Arg Leu Thr
 180 185 190
 Ala Ala Lys Leu Glu Pro Pro Leu Ile Pro Phe Met Pro Leu Leu Ile
 195 200 205
 Lys Asp Met Thr Phe Thr His Glu Gly Asn Lys Thr Phe Ile Asp Asn
 210 215 220
 Leu Val Asn Phe Glu Lys Met Arg Met Ile Ala Asn Thr Ala Arg Thr
 225 230 235 240
 Val Arg Tyr Tyr Arg Ser Gln Pro Phe Asn Pro Asp Ala Ala Gln Ala
 245 250 255
 Asn Lys Asn His Gln Asp Val Arg Ser Tyr Val Arg Gln Leu Asn Val
 260 265 270
 Ile Asp Asn Gln Arg Thr Leu Ser Gln Met Ser His Arg Leu Glu Pro
 275 280 285
 Arg Arg Pro
 290

<210> 17
 <211> 3013
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (407)..(2953)
 <223> CAMP-GEFII

<400> 17
 gatccagcga agatgtggat ataatcttca ctgcactgaa agaagttaaa gcttttgaga 60
 aatttcaccc aaatctcctt catcagattt gcttatgtgg ttattatgag aatctggaaa 120
 aggaataaac attatttcgc caggggtgata tggaacaaac tggatatgctg cctggcaggg 180

ggc aaq tta qca cta gtg aat gat gcc cca cga gct gcc tct atc gtc 1231

- 41 -

Gly 260	Lys	Leu	Ala	Leu	Val 265	Asn	Asp	Ala	Pro	Arg 270	Ala	Ala	Ser	Ile	Val 275	
tta	cga	gaa	gat	aac	tgc	cat	ttc	tta	aga	gta	gac	aag	gag	gat	ttc	1279
Leu	Arg	Glu	Asp	Asn 280	Cys	His	Phe	Leu	Arg 285	Val	Asp	Lys	Glu	Asp 290	Phe	
aac	cgg	atc	cta	agg	gac	gtg	gag	gcg	aat	aca	gtc	aga	ctt	aaa	gaa	1327
Asn	Arg	Ile	Leu 295	Arg	Asp	Val	Glu	Ala 300	Asn	Thr	Val	Arg	Leu 305	Lys	Glu	
cat	gac	caa	gat	gtc	ttg	gtg	ctg	gag	aag	gtc	cca	gca	ggg	aac	aga	1375
His	Asp	Gln 310	Asp	Val	Leu	Val	Leu 315	Glu	Lys	Val	Pro	Ala 320	Gly	Asn	Arg	
gct	tct	aat	caa	gga	aac	tca	cag	cct	cag	caa	aag	tat	act	gtg	atg	1423
Ala	Ser	Asn	Gln	Gly	Asn	Ser 330	Gln	Pro	Gln	Gln	Lys 335	Tyr	Thr	Val	Met	
tca	gga	aca	cct	gaa	aaa	att	tta	gag	cat	ttt	cta	gaa	aca	ata	cgc	1471
Ser	Gly	Thr	Pro	Glu	Lys 345	Ile	Leu	Glu	His	Phe 350	Leu	Glu	Thr	Ile	Arg 355	
ctt	gag	gca	act	tta	aat	gaa	gca	aca	gat	tct	ggt	tta	aat	gac	ttt	1519
Leu	Glu	Ala	Thr	Leu 360	Asn	Glu	Ala	Thr	Asp 365	Ser	Val	Leu	Asn	Asp 370	Phe	
att	atg	atg	cac	tgt	gtt	ttt	atg	cca	aat	acc	cag	ctt	tgc	ccg	gca	1567
Ile	Met	Met	His 375	Cys	Val	Phe	Met	Pro 380	Asn	Thr	Gln	Leu	Cys 385	Pro	Ala	
ctg	gtg	gcc	cac	tac	cac	gca	cag	cct	tca	caa	ggt	aca	gaa	cag	gag	1615
Leu	Val	Ala 390	His	Tyr	His	Ala	Gln 395	Pro	Ser	Gln	Gly	Thr 400	Glu	Gln	Glu	
aaa	atg	gat	tat	gcc	ctc	aac	aat	aag	agg	cga	gtc	atc	cgc	ctg	gtt	1663
Lys	Met	Asp	Tyr	Ala	Leu 410	Asn	Asn	Lys	Arg	Arg	Val 415	Ile	Arg	Leu	Val	
cta	cag	tggt	gct	gcc	atg	tat	gga	gac	ctc	ctg	caa	gag	gat	gac	gta	1711
Leu	Gln	Trp	Ala	Ala	Met 425	Tyr	Gly	Asp	Leu	Leu 430	Gln	Glu	Asp	Asp	Val 435	
tct	atg	gcc	ttc	ctg	gag	gag	ttt	tat	gta	tct	gta	tca	gat	gat	gcc	1759
Ser	Met	Ala	Phe 440	Leu	Glu	Glu	Phe	Tyr	Val 445	Ser	Val	Ser	Asp	Asp 450	Ala	
cgg	atg	att	gct	gcc	ctc	aag	gag	caa	ctg	cca	gag	ttg	gag	aag	att	1807
Arg	Met	Ile	Ala 455	Ala	Leu	Lys	Glu	Gln 460	Leu	Pro	Glu	Leu	Glu 465	Lys	Ile	
gtc	aag	caa	atc	tca	gaa	gat	gca	aag	gca	cca	caa	aag	aag	cac	aag	1855
Val	Lys	Gln 470	Ile	Ser	Glu	Asp	Ala 475	Lys	Ala	Pro	Gln	Lys 480	Lys	His	Lys	
gtt	ctt	ttg	caa	cag	ttc	aat	acg	ggc	gat	gag	aga	gcc	cag	aag	cgc	1903
Val	Leu	Leu	Gln	Gln	Phe	Asn 490	Thr	Gly	Asp	Glu	Arg 495	Ala	Gln	Lys	Arg	
cag	cct	atc	cgc	ggc	tct	gat	gaa	gtt	ctg	ttt	aag	gtc	tat	tgc	atg	1951
Gln	Pro	Ile	Arg	Gly	Ser 505	Asp	Glu	Val	Leu	Phe 510	Lys	Val	Tyr	Cys	Met 515	
gac	cac	acc	tac	aca	acc	att	cgg	gtg	cca	gtg	gcc	act	tgc	gtg	aag	1999
Asp	His	Thr	Tyr	Thr 520	Thr	Ile	Arg	Val	Pro 525	Val	Ala	Thr	Ser	Val 530	Lys	
gaa	gtc	atc	agt	gca	gtt	gcc	gac	aag	ctg	ggc	tcc	ggg	gag	ggc	ctg	2047
Glu	Val	Ile	Ser 535	Ala	Val	Ala	Asp	Lys 540	Leu	Gly	Ser	Gly	Glu 545	Gly	Leu	
atc	ata	gtc	aag	atg	agt	tcc	gga	gga	gaa	aag	gtg	gtg	ctc	aaa	cct	2095
Ile	Ile	Val 550	Lys	Met	Ser	Ser	Gly 555	Gly	Glu	Lys	Val	Val 560	Leu	Lys	Pro	

- 42 -

aat gat gtt tca gta ttt acg acg ctc acc att aat gga cgc ctg ttt	2143
Asn Asp Val Ser Val Phe Thr Thr Leu Thr Ile Asn Gly Arg Leu Phe	
565 570 575	
gct tgc ccg cga gag caa ttc gat tca ctg act ccc tta cca gaa cag	2191
Ala Cys Pro Arg Glu Gln Phe Asp Ser Leu Thr Pro Leu Pro Glu Gln	
580 585 590 595	
gaa ggc cca act gtt gga aca gtg gga act ttt gaa ctg atg agc tcc	2239
Glu Gly Pro Thr Val Gly Thr Val Gly Thr Phe Glu Leu Met Ser Ser	
600 605 610	
aaa gat tta gca tac cag atg aca att tat gat tgg gaa ctc ttc aac	2287
Lys Asp Leu Ala Tyr Gln Met Thr Ile Tyr Asp Trp Glu Leu Phe Asn	
615 620 625	
tgc gtg cat gag ctg gag cta atc tat cac aca ttt gga agg cat aat	2335
Cys Val His Glu Leu Glu Leu Ile Tyr His Thr Phe Gly Arg His Asn	
630 635 640	
ttt aaa aag acc aca gca aac ttg gat ttg ttc ctg agg aga ttt aat	2383
Phe Lys Lys Thr Thr Ala Asn Leu Asp Leu Phe Leu Arg Arg Phe Asn	
645 650 655	
gaa att cag ttt tgg gtc gtc act gag atc tgc ctt tgt tct cag ctc	2431
Glu Ile Gln Phe Trp Val Val Thr Glu Ile Cys Leu Cys Ser Gln Leu	
660 665 670 675	
agc aag cgt gtt cag cta tta aaa aaa ttt att aag ata gca gcc cac	2479
Ser Lys Arg Val Gln Leu Leu Lys Lys Phe Ile Lys Ile Ala Ala His	
680 685 690	
tgt aag gag tat aaa aat ctg aat tcc ttt ttt gcc atc gtc atg gga	2527
Cys Lys Glu Tyr Lys Asn Leu Asn Ser Phe Phe Ala Ile Val Met Gly	
695 700 705	
cta agt aac att gct gtg agc cgc ttg gca cta acg tgg gag aaa ctg	2575
Leu Ser Asn Ile Ala Val Ser Arg Leu Ala Leu Thr Trp Glu Lys Leu	
710 715 720	
cca agc aag ttc aag aag ttc tat gcg gag ttt gaa agt tta atg gac	2623
Pro Ser Lys Phe Lys Lys Phe Tyr Ala Glu Phe Glu Ser Leu Met Asp	
725 730 735	
cct tca agg aac cac agg gcc tac agg ctg aca gta gct aag ctg gaa	2671
Pro Ser Arg Asn His Arg Ala Tyr Arg Leu Thr Val Ala Lys Leu Glu	
740 745 750 755	
cct cct ctc atc ccc ttc atg cct ttg ctc att aaa gat atg aca ttt	2719
Pro Pro Leu Ile Pro Phe Met Pro Leu Leu Ile Lys Asp Met Thr Phe	
760 765 770	
act cat gag ggg aac aag acg ttc att gac aat cta gta aac ttt gaa	2767
Thr His Glu Gly Asn Lys Thr Phe Ile Asp Asn Leu Val Asn Phe Glu	
775 780 785	
aaa atg cgc atg att gca aat acg gcc aga aca gtg aga tac tac agg	2815
Lys Met Arg Met Ile Ala Asn Thr Ala Arg Thr Val Arg Tyr Tyr Arg	
790 795 800	
agc caa ccc ttc aat cct gat gca gct caa gct aat aag aac cat cag	2863
Ser Gln Pro Phe Asn Pro Asp Ala Ala Gln Ala Asn Lys Asn His Gln	
805 810 815	
gat gtc cgg agt tat gta cgg caa tta aat gtg att gac aac cag aga	2911
Asp Val Arg Ser Tyr Val Arg Gln Leu Asn Val Ile Asp Asn Gln Arg	
820 825 830 835	
act tta tca cag atg tca cac aga tta gag cct cgt cga cca	2953
Thr Leu Ser Gln Met Ser His Arg Leu Glu Pro Arg Arg Pro	
840 845	
tagacatttc aaatgcccaa agcaacagtt tgcctccagt ccacaatttt caaaaatgcc	3013

- 43 -

<210> 18

<211> 849

<212> PRT

<213> Homo sapiens

<400> 18

```

Met Val Leu Trp Lys Arg Ala Leu Thr Met Thr Gly Phe Leu Thr Arg
 1          5          10          15
Arg Thr His Leu Ile Glu Pro His Val Pro Leu Arg Pro Ala Asn Thr
          20          25          30
Ile Thr Lys Val Pro Ser Glu Lys Ile Leu Arg Ala Gly Lys Ile Leu
 35          40          45
Arg Asn Ala Ile Leu Ser Arg Ala Pro His Met Ile Arg Asp Arg Lys
 50          55          60
Tyr His Leu Lys Thr Tyr Arg Gln Cys Cys Val Gly Thr Glu Leu Val
 65          70          75          80
Asp Trp Met Ile Asp Glu Thr Pro Cys Val His Ser Arg Thr Gln Ala
 85          90          95
Val Gly Met Trp Gln Val Leu Leu Glu Asp Gly Val Leu Asn His Val
100          105          110
Asp Gln Glu His His Phe Gln Asp Phe Tyr Leu Phe Tyr Arg Phe Leu
115          120          125
Asp Asp Glu His Glu Asp Ala Pro Leu Pro Thr Glu Glu Glu Lys Lys
130          135          140
Glu Cys Asp Glu Glu Leu Gln Asp Thr Met Leu Leu Leu Ser Gln Met
145          150          155          160
Gly Pro Asp Ala His Met Arg Met Ile Leu Arg Lys Pro Pro Gly Gln
165          170          175
Arg Thr Val Asp Asp Leu Glu Ile Ile Tyr Glu Glu Leu Leu His Ile
180          185          190
Lys Ala Leu Ser His Leu Ser Thr Thr Val Lys Arg Glu Leu Ala Gly
195          200          205
Val Leu Ile Phe Glu Ser His Ala Lys Gly Gly Thr Val Leu Phe Asn
210          215          220
Gln Gly Glu Glu Gly Thr Ser Trp Tyr Ile Ile Leu Lys Gly Ser Val
225          230          235          240
Asn Val Val Ile Tyr Gly Lys Gly Val Val Cys Thr Leu His Glu Gly
245          250          255
Asp Asp Phe Gly Lys Leu Ala Leu Val Asn Asp Ala Pro Arg Ala Ala
260          265          270
Ser Ile Val Leu Arg Glu Asp Asn Cys His Phe Leu Arg Val Asp Lys
275          280          285
Glu Asp Phe Asn Arg Ile Leu Arg Asp Val Glu Ala Asn Thr Val Arg
290          295          300
Leu Lys Glu His Asp Gln Asp Val Leu Val Leu Glu Lys Val Pro Ala
305          310          315          320
Gly Asn Arg Ala Ser Asn Gln Gly Asn Ser Gln Pro Gln Gln Lys Tyr
325          330          335
Thr Val Met Ser Gly Thr Pro Glu Lys Ile Leu Glu His Phe Leu Glu
340          345          350
Thr Ile Arg Leu Glu Ala Thr Leu Asn Glu Ala Thr Asp Ser Val Leu
355          360          365

```

- 44 -

Asn Asp Phe Ile Met Met His Cys Val Phe Met Pro Asn Thr Gln Leu
 370 375 380
 Cys Pro Ala Leu Val Ala His Tyr His Ala Gln Pro Ser Gln Gly Thr
 385 390 395 400
 Glu Gln Glu Lys Met Asp Tyr Ala Leu Asn Asn Lys Arg Arg Val Ile
 405 410 415
 Arg Leu Val Leu Gln Trp Ala Ala Met Tyr Gly Asp Leu Leu Gln Glu
 420 425 430
 Asp Asp Val Ser Met Ala Phe Leu Glu Glu Phe Tyr Val Ser Val Ser
 435 440 445
 Asp Asp Ala Arg Met Ile Ala Ala Leu Lys Glu Gln Leu Pro Glu Leu
 450 455 460
 Glu Lys Ile Val Lys Gln Ile Ser Glu Asp Ala Lys Ala Pro Gln Lys
 465 470 475 480
 Lys His Lys Val Leu Leu Gln Gln Phe Asn Thr Gly Asp Glu Arg Ala
 485 490 495
 Gln Lys Arg Gln Pro Ile Arg Gly Ser Asp Glu Val Leu Phe Lys Val
 500 505 510
 Tyr Cys Met Asp His Thr Tyr Thr Thr Ile Arg Val Pro Val Ala Thr
 515 520 525
 Ser Val Lys Glu Val Ile Ser Ala Val Ala Asp Lys Leu Gly Ser Gly
 530 535 540
 Glu Gly Leu Ile Ile Val Lys Met Ser Ser Gly Gly Glu Lys Val Val
 545 550 555 560
 Leu Lys Pro Asn Asp Val Ser Val Phe Thr Thr Leu Thr Ile Asn Gly
 565 570 575
 Arg Leu Phe Ala Cys Pro Arg Glu Gln Phe Asp Ser Leu Thr Pro Leu
 580 585 590
 Pro Glu Gln Glu Gly Pro Thr Val Gly Thr Val Gly Thr Phe Glu Leu
 595 600 605
 Met Ser Ser Lys Asp Leu Ala Tyr Gln Met Thr Ile Tyr Asp Trp Glu
 610 615 620
 Leu Phe Asn Cys Val His Glu Leu Glu Leu Ile Tyr His Thr Phe Gly
 625 630 635 640
 Arg His Asn Phe Lys Lys Thr Thr Ala Asn Leu Asp Leu Phe Leu Arg
 645 650 655
 Arg Phe Asn Glu Ile Gln Phe Trp Val Val Thr Glu Ile Cys Leu Cys
 660 665 670
 Ser Gln Leu Ser Lys Arg Val Gln Leu Leu Lys Lys Phe Ile Lys Ile
 675 680 685
 Ala Ala His Cys Lys Glu Tyr Lys Asn Leu Asn Ser Phe Phe Ala Ile
 690 695 700
 Val Met Gly Leu Ser Asn Ile Ala Val Ser Arg Leu Ala Leu Thr Trp
 705 710 715 720
 Glu Lys Leu Pro Ser Lys Phe Lys Lys Phe Tyr Ala Glu Phe Glu Ser
 725 730 735
 Leu Met Asp Pro Ser Arg Asn His Arg Ala Tyr Arg Leu Thr Val Ala
 740 745 750
 Lys Leu Glu Pro Pro Leu Ile Pro Phe Met Pro Leu Leu Ile Lys Asp
 755 760 765

- 45 -

Met Thr Phe Thr His Glu Gly Asn Lys Thr Phe Ile Asp Asn Leu Val
 770 775 780

Asn Phe Glu Lys Met Arg Met Ile Ala Asn Thr Ala Arg Thr Val Arg
 785 790 795 800

Tyr Tyr Arg Ser Gln Pro Phe Asn Pro Asp Ala Ala Gln Ala Asn Lys
 805 810 815

Asn His Gln Asp Val Arg Ser Tyr Val Arg Gln Leu Asn Val Ile Asp
 820 825 830

Asn Gln Arg Thr Leu Ser Gln Met Ser His Arg Leu Glu Pro Arg Arg
 835 840 845

Pro



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7 : C12N 15/12, C07K 14/82, 16/32, 14/47, C12Q 1/68, C12N 5/10, A01K 67/027, G01N 33/53, 33/574		A3	(11) International Publication Number: WO 00/24768
			(43) International Publication Date: 4 May 2000 (04.05.00)
(21) International Application Number: PCT/US99/24826		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 22 October 1999 (22.10.99)		Published With international search report.	
(30) Priority Data: 60/105,507 23 October 1998 (23.10.98) US 60/108,685 16 November 1998 (16.11.98) US		(88) Date of publication of the international search report: 9 November 2000 (09.11.00)	
(71) Applicant (for all designated States except US): MASSACHUSETTS INSTITUTE OF TECHNOLOGY [US/US]; 77 Massachusetts Avenue, Cambridge, MA 02138 (US).			
(72) Inventors; and			
(75) Inventors/Applicants (for US only): KAWASAKI, Hiroaki [JP/JP]; 3-20-2, Aoba, Higashi-ku, Fukuoka 813-0025, (JP). GRAYBIEL, Ann [US/US]; Boyce Farm Road, Lincoln, MA 01773 (US). HOUSMAN, David [US/US]; 64 Homer Street, Newton, MA 02158 (US).			
(74) Agent: CAMACHO, Jennifer, A.; Testa, Hurwitz & Thibault, LLP, High Street Tower, 125 High Street, Boston, MA 02110 (US).			
(54) Title: GENES INTEGRATING SIGNAL TRANSDUCTION PATHWAYS			
(57) Abstract <p>The present invention describes the identification, isolation, sequencing and characterization of two human CalDAG-GEF, and two human cAMP-GEF genes, which are associated with the Ras pathway. Also identified are CalDAG-GEF gene homologues in mice and cAMP-GEF gene homologues in rats. Nucleic acids and proteins comprising or derived from the CalDAG-GEFs and/or cAMP-GEFs are useful in screening and diagnosing certain Ras-associated cancers, in identifying and developing therapeutics for treatment of certain Ras-associated cancers, and in producing cell lines and transgenic animals useful as models of Ras-associated cancers.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

INTERNATIONAL SEARCH REPORT

Internat Application No
PCT/US /24826

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/12 C07K14/82 C07K16/32 C07K14/47 C12Q1/68
C12N5/10 A01K67/027 G01N33/53 G01N33/574

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K C12Q A01K G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EBINU, J.O. ET AL.: "RasGRP, a Ras guanyl nucleotide-releasing protein with calcium- and diacylglycerol-binding motifs." SCIENCE, (1998 MAY 15) 280 (5366) 1082-6, XP000882708 the whole document	1
A	--- GOTOH, T. ET AL.: "Identification of Rap1 as a target for the Crk SH3 domain-binding guanine nucleotide-releasing factor C3G." MOLECULAR AND CELLULAR BIOLOGY, (1995) 15 (12) 6746-53, XP000881340 the whole document --- -/-	1

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

Z document member of the same patent family

Date of the actual completion of the international search

5 April 2000

Date of mailing of the international search report

3 July 2000 (03.07.00)

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl
Fax: (+31-70) 340-3016

Authorized officer

Nichogiannopoulou, A

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>KAIBUCHI, K. ET AL.: "Molecular cloning of the cDNA for stimulatory GDP/GTP exchange protein for smg p21s (ras p21-like small GTP-binding proteins) and characterization of stimulatory GDP/GTP exchange protein." MOLECULAR AND CELLULAR BIOLOGY, (1991) 11 (5) 2873-80, XP000881341 the whole document</p>	1
X	<p>KEDRA D ET AL: "THE GERMINAL CENTER KINASE GENE AND A NOVEL CDC25-LIKE GENE ARE LOCATED IN THE VICINITY OF THE PYGM GENE ON 11Q13" HUM. GENET., vol. 100, 1 October 1997 (1997-10-01), pages 611-619, XP002069545 page 613, last paragraph -page 615, paragraph FIRST; figure 2 & DATABASE EMBL [Online] AC Y12336, 19 June 1997 (1997-06-19) KEDRA D ET AL: "H. sapiens mRNA for F25B3.3 kinase like protein from C. elegans" Protein with 96.9% identity to SEQ ID No:2 and 100% identity to SEQ ID No:4 the whole document</p>	1,3,5-8, 10, 38-54, 63-69
P,X	<p>WO 98 53061 A (QUEENSLAND INST MED RES; HANCOCK JOHN (AU); SILINS GINTERS (AU)) 26 November 1998 (1998-11-26)</p> <p>MCG7, a hu protein with 100% identity in 609 aa overlap with SEQ ID No:4 claim 5; figure 13B</p>	1,3,5-8, 10, 38-54, 63-69
P,X	<p>KAWASAKI H ET AL: "A Rap guanine nucleotide exchange factor enriched highly in the basal ganglia" PROC. NATL. ACAD. SCI. USA, vol. 95, October 1998 (1998-10), pages 13278-13283, XP000882748 the whole document</p>	1,3,5-8, 10, 38-54, 63-69

INTERNATIONAL SEARCH REPORT

Intern. application No.
PCT/US 99/24826

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 76-81, 86-109 and 127 -in as far as they concern in vivo methods- are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

See additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

1, 3-8, 10, 12-19, 28-32, 38-70, 72, 73-75, 76, 78-82, 84, 86, 88-98, 110-112, 116, 118, 119, 121-129 (all partially)

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

1. Claims: 1, 3-8, 10, 12-19, 28-32, 38-70, 72, 73-75, 76, 78-82, 84, 86, 88-98, 110-112, 116, 118, 119, 121-129 (all partially)

Claims relating to CalDAG-GEFI proteins and nucleic acids encoding them, mutants, variants, species orthologs and homologues, functional domains and antigenic determinants thereof. Methods for identifying variants or homologues of such proteins. Recombinant expression vectors, host cells and animal models for cancer expressing such recombinant modified proteins. Methods for producing said proteins and pure preparations of such. Antibodies selectively binding to such proteins, methods and cell lines for producing them. Methods for identifying compounds that modulate the expression of such proteins, and for identifying compounds that can selectively bind to them. Diagnostic methods for detecting mutations and pharmaceutical preparations comprising pure protein, expression vectors encoding the protein or antisense sequences. Pharmaceutical preparations comprising antibodies or antigenic determinants and methods of treatment.

2. Claims: 1, 3-8, 10, 12-19, 28-32, 38-70, 72, 73-75, 76, 78-82, 84, 86, 88-98, 110-112, 116, 118, 119, 121-129 (all partially)

As in subject 1, the proteins being CalDAG-GEFII proteins.

3. Claims: 2, 4-7, 9, 11, 20-27, 33-69, 71-75, 77-81, 83, 85, 87, 99-109, 113-115, 117, 118, 120-126, 130 (all partially)

As in subject 1, the proteins being cAMP-GEFI proteins.

4. Claims: 2, 4-7, 9, 11, 20-27, 33-69, 71-75, 77-81, 83, 85, 87, 99-109, 113-115, 117, 118, 120-126, 130 (all partially)

As in subject 1, the proteins being cAMP-GEFII proteins.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/24826

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9853061 A	26-11-1998	AU 7512998 A	11-12-1998

THIS PAGE BLANK (USPTO)